



Polya Trees and Random Distributions

R. Daniel Mauldin; William D. Sudderth; S. C. Williams

The Annals of Statistics, Vol. 20, No. 3 (Sep., 1992), 1203-1221.

Stable URL:

<http://links.jstor.org/sici?sici=0090-5364%28199209%2920%3A3%3C1203%3APTARD%3E2.0.CO%3B2-G>

The Annals of Statistics is currently published by Institute of Mathematical Statistics.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ims.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact jstor-info@umich.edu.

POLYA TREES AND RANDOM DISTRIBUTIONS

BY R. DANIEL MAULDIN,¹ WILLIAM D. SUDDERTH² AND S. C. WILLIAMS

*University of North Texas, University of Minnesota
and Utah State University*

Trees of Polya urns are used to generate sequences of exchangeable random variables. By a theorem of de Finetti each such sequence is a mixture of independent, identically distributed variables and the mixing measure can be viewed as a prior on distribution functions. The collection of these *Polya tree priors* forms a convenient conjugate family which was mentioned by Ferguson and includes the Dirichlet processes of Ferguson. Unlike Dirichlet processes, Polya tree priors can assign probability 1 to the class of continuous distributions. This property and a few others are investigated.

1. Introduction. The Polya urn scheme is, perhaps, the simplest and most concrete way to generate a sequence X_1, X_2, \dots of exchangeable random variables having values in a finite set $E = \{0, \dots, k\}$. Suppose that the urn u has initially u_i balls of color i for $i \in E$ and, that, at each stage, a ball is drawn at random and replaced by two of the same color. Let $X_n = i$ if the n th ball selected is of color i . It is well known that the X_n are exchangeable and that the sample distribution of X_1, \dots, X_n converges almost surely to a random probability vector $\Theta = (\Theta_0, \dots, \Theta_k)$ which has a Dirichlet distribution with parameters (u_0, \dots, u_k) . Furthermore, given $\Theta = \theta$, the variables X_1, X_2, \dots are independent with $P[X_n = i] = \theta_i$ for all n and i . (These facts are reviewed in the next section.)

All of these results were generalized by Blackwell and MacQueen (1973) who showed that the random distributions constructed by Ferguson (1973) can be viewed as the limit of the sample distributions of variables which are obtained from a Polya urn scheme based on a continuum of colors. The urn scheme makes many properties of Ferguson distributions intuitively clear. For example, the Ferguson distributions form a conjugate family of prior distributions for nonparametric problems just as the Dirichlet distributions form a conjugate family for multinomial sampling. [By the way, Ferguson (1973) called his distributions “Dirichlet processes.” To avoid confusion with the processes constructed here, we will continue to call them Ferguson distributions.]

This paper shows how to construct another conjugate family of prior distributions from trees of Polya urns. Mauldin and Williams (1990) gave the

Received August 1990; revised May 1991.

¹Research supported by NSF Grant DMS-88-03361.

²Research supported by NSF Grant DMS-89-11548.

AMS 1980 subject classifications. Primary 62A15; secondary 62G99, 60G09, 60G57.

Key words and phrases. Prior distributions, random measures, Polya urns, Dirichlet distributions.

first such construction and the following is a natural generalization. Let E^* be the set of all finite sequences of elements of $E = \{0, \dots, k\}$ including the empty sequence ϕ . A *Polya tree* is a function that assigns to every p in E^* an urn $u(p)$ which contains $u(p)_i$ balls of color i for each $i \in E$. The Polya tree u can be used to generate a sequence of random variables X_{11}, X_{12}, \dots and a new tree $u^{(1)}$ as follows: Draw a ball at random from $u(\phi)$ and replace it by two of the same color. Set $X_{11} = i_1$ if the ball is of color i_1 . Next draw a ball from $u(i_1)$, replace it by two of the same color, and set $X_{12} = i_2$ if the ball is of color i_2 . Go on to $u(i_1, i_2)$ and continue in this fashion. Set $X_1 = (X_{11}, X_{12}, \dots)$ and let $u^{(1)}$ be the Polya tree which was obtained in the construction. Iterate the entire process to obtain X_1, X_2, \dots and $u^{(1)}, u^{(2)}, \dots$. It is shown in Theorem 4.1 that the X_n are exchangeable. So, by a theorem of de Finetti, there is a measure $Q = Q_u$ defined on the space of probability measures on $E^N = E \times E \times \dots$ such that the distribution of X_1, X_2, \dots can be obtained by first choosing Θ with distribution Q and then choosing X_1, X_2, \dots to be independent with distribution θ given $\Theta = \theta$. This de Finetti measure Q will be calculated explicitly (Theorem 4.2).

The measure Q can be regarded as a prior which picks a random distribution on E^N and the collection of these priors based on Polya trees forms a conjugate family. (If Θ is a random probability with distribution Q_u and X_1, \dots, X_n is a random sample from Θ , then, by Theorem 4.3, the posterior distribution of Θ is $Q_{u^{(n)}}$ where $u^{(n)}$ is the Polya tree after the n th stage of the construction above.)

If E^N is mapped measurably into another space, then the structure of the Polya priors can be carried over to the new space to obtain random distributions there. For example, let I be the unit interval and define $\psi: E^N \rightarrow I$ by

$$(1.1) \quad \psi(x_1, x_2, \dots) = \sum_{n=1}^{\infty} \frac{x_n}{(k+1)^n}.$$

Let X_1, X_2, \dots be the E^N -valued variables constructed above and set $Y_n = \psi(X_n)$ for $n = 1, 2, \dots$. The Y_n are exchangeable because the X_n are and the de Finetti measure $\bar{Q} = \bar{Q}_u$ for the Y_n is easily calculated from Q . We will use this example throughout the paper, but there are other mappings which may be of interest also. Ferguson has suggested mapping E^N to the probability simplex in dimension $k+1$ by the rule

$$(x_1, x_2, x_3, \dots) \rightarrow \frac{1}{k+1} \left(e_{x_1} + \frac{k}{k+1} \left(e_{x_2} + \frac{k}{k+1} (e_{x_3} + \dots) \right) \right),$$

where $e_0 = (1, 0, \dots, 0)$, $e_1 = (0, 1, \dots, 0)$, \dots , $e_k = (0, 0, \dots, 1)$ are unit vectors. Lavine (1990) maps E^N to other spaces by using trees of nested partitions.

The priors \bar{Q}_u based on Polya trees are tailfree processes in the sense of Freedman (1963) and Fabius (1964). A nice discussion of some of their properties is given by Ferguson (1974). He points out that the family of Polya

tree priors includes the Ferguson distributions of his 1973 paper, but that, unlike Ferguson distributions, Polya tree priors can assign probability 1 to the set of continuous probability measures. Simple conditions are given in Theorem 5.2 and its corollary to guarantee that \bar{Q}_u is supported by the continuous measures. Ferguson (1974) also points out that results of Kraft (1964) can be used to give conditions on u which guarantee that \bar{Q}_u assigns probability 1 to the set of distributions which have densities with respect to Lebesgue measure.

The priors \bar{Q}_u defined on probability measures on $[0, 1]$ are also related to the distributions constructed by Dubins and Freedman (1967) in a different fashion. In fact, if $k = 1$ and if $u(p)$ contains exactly one ball of color 0 and one ball of color 1 for every p , then \bar{Q}_u was shown by Mauldin and Williams (1991) to be one of the Dubins and Freedman measures. The family of all \bar{Q}_u is the natural conjugate family of priors which contains this basic measure. [See Graf, Mauldin and Williams (1986) for another generalization of the Dubins and Freedman measures.]

The next section is a review of classical Polya urn schemes, Dirichlet distributions and their connection by way of de Finetti's theorem. After these preliminaries a study is made in Section 3 of Polya tree processes which end after a finite number of stages. This is in preparation for the study of infinite trees and also provides a conjugate family of priors for finite, sequential sampling schemes. Section 4 treats infinite Polya trees u and characterizes the de Finetti measures Q_u . Section 5 gives conditions under which Q_u concentrates on continuous distributions and Section 6 gives conditions under which the support of Q_u is large. The expected distribution function and the expected mean of a Polya tree prior are calculated in Sections 7 and 8.

Applications of Polya trees are discussed by Lavine (1992). He shows how to construct a Polya tree prior with a given predictive density and how to use mixtures of them to model uncertainty about a parametric model. Examples with data are presented.

2. Polya urns and Dirichlet distributions. The initial data needed for the classical Polya urn scheme described in the introduction are the vector $u = (u_0, \dots, u_k)$, where each u_i represents the initial number of balls of color i . Such a vector will be called an *urn vector* if u_i is nonnegative for every $i \in E$ and the quantity $|u| = u_0 + \dots + u_k$ is strictly positive. The probability of drawing a ball of color i from u is, by definition, $u_i/|u|$ for each i . After drawing a ball of color i , the urn at the next stage has $u_i + 1$ balls of color i . With these conventions, the Polya urn scheme, as described in the introduction, generates a sequence

$$X = (X_1, X_2, \dots)$$

of random variables with values in E . (As before, $X_n = i$ if the n th ball drawn has color i .) The sequence X is said to have a *Polya distribution with parameter u* or, more briefly, X is $\mathcal{P}(u)$.

In order to calculate the joint distribution of X_1, \dots, X_n , let $j = (j_1, \dots, j_n)$ be a sequence of n elements of E and set

$$(2.1) \quad P(j; u) = P[X_1 = j_1, \dots, X_n = j_n].$$

For each $i \in E$, let $c(i)$ be the number of i 's occurring in the sequence j and define $s(i) = u_i + c(i)$. Notice that, if $X_1 = j_1, \dots, X_n = j_n$, then $s(i)$ corresponds to the number of balls of color i in the urn after the n th stage.

A simple counting argument shows that

$$(2.2) \quad P(j; u) = \frac{(s(0) - 1)_{c(0)} \cdots (s(k) - 1)_{c(k)}}{(|u| + n - 1)_n},$$

where, as usual,

$$(a)_0 = 1 \quad \text{and} \quad (a)_b = a(a - 1) \cdots (a - b + 1), \quad b = 1, 2, \dots$$

It is clear from (2.2) that $P(j; u)$ is unchanged when the coordinates of j are permuted. We record this in a lemma.

LEMMA 2.1. *If (X_1, X_2, \dots) has a Polya distribution, then X_1, X_2, \dots are exchangeable.*

The next result is a version of de Finetti's theorem on exchangeable variables. It is a special case of results in Meyer (1966) or in Aldous (1983).

THEOREM 2.1. *Let X_1, X_2, \dots be exchangeable variables with values in $E = \{0, \dots, k\}$. For $n = 1, 2, \dots$ and $i \in E$, let $C_n(i) = \#\{j \leq n: X_j = i\}$. Then:*

(a) *$(C_n(0), \dots, C_n(k))/n$ converges almost surely to a random probability vector $\Theta = (\Theta_0, \dots, \Theta_k)$.*

(b) *Additionally, given Θ , the variables X_1, X_2, \dots are independent and each has distribution Θ .*

The distribution of the random probability vector Θ in Theorem 2.1 is called the *de Finetti measure* for the sequence $X = (X_1, X_2, \dots)$. [It is called the *directing measure* by Aldous (1983).]

Let

$$S_k = \left\{ (\theta_0, \dots, \theta_k) : \theta_0 \geq 0, \dots, \theta_k \geq 0, \sum_{i=0}^k \theta_i = 1 \right\}.$$

Here is a simple characterization of the de Finetti measure.

THEOREM 2.2. *A probability measure μ defined on the Borel subsets of S_k is the de Finetti measure for the exchangeable sequence X if and only if, for every finite sequence i_1, \dots, i_m of elements of E*

$$(2.3) \quad P[X_i = i_1, \dots, X_m = i_m] = \int \theta_0^{c_0} \cdots \theta_k^{c_k} d\mu(\theta_0, \dots, \theta_k),$$

where c_j is the number of elements in i_1, \dots, i_m which are equal to j for each j .

PROOF. If μ is the de Finetti measure for X , formula (2.3) follows from Theorem 2.1(b). Conversely, μ is uniquely determined by its moments since it has compact support. \square

Notice that (2.3) is equivalent to

$$(2.4) \quad P[X_1 = i_1, \dots, X_m = i_m] = E \left[\prod_{a=1}^m \Theta_{i_a} \right],$$

where Θ is a random probability vector with distribution μ .

Our next task is to identify the de Finetti measure when X is a Polya sequence. First we recall the definition of a Dirichlet distribution.

The Dirichlet distribution defined here will be slightly more general than usual in that some of the variables can be degenerate at zero. Our definition is consistent with that of Ferguson (1973). Let $\Theta = (\Theta_0, \dots, \Theta_k)$ be a random vector with values in S_k and let $u = (u_0, \dots, u_k)$ be an urn vector. Suppose first that $u_i > 0$ for all i . Then we say Θ is *Dirichlet with parameter u* , written $\mathbf{D}(u)$, if $k = 0$ and $\Theta_0 = 1$ almost surely or if $k > 0$ and $(\Theta_0, \dots, \Theta_{k-1})$ has the density function

$$f(\theta_0, \dots, \theta_{k-1}) = \frac{\Gamma(|u|)}{\Gamma(u_0) \cdots \Gamma(u_k)} \theta_0^{u_0} \cdots \theta_k^{u_k}$$

for $(\theta_0, \dots, \theta_k) \in S_k$. [Notice $\theta_k = 1 - (\theta_0 + \cdots + \theta_{k-1})$.] In the general case, take F to be that subset of $\{0, \dots, k\}$ consisting of those i for which $u_i > 0$. Say $F = \{i_0, \dots, i_r\}$ where $0 \leq r \leq k$. Now define Θ to be $\mathbf{D}(u)$ if $(\Theta_{i_0}, \dots, \Theta_{i_r})$ is $\mathbf{D}((u_{i_0}, \dots, u_{i_r}))$ and $\Theta_j = 0$ almost surely for $j \notin F$.

The next result gives the nice connection between Polya urn schemes and the Dirichlet distribution. It is a special case of the theorem in Blackwell and MacQueen (1973). Much of it can also be found in Blackwell and Kendall (1964). An elementary proof can be based on Theorem 2.2.

THEOREM 2.3. *If X is $\mathcal{P}(u)$, then the de Finetti measure for X is $\mathbf{D}(u)$.*

The Dirichlet family of distributions is useful for Bayesian analysis largely because it is the natural family conjugate to the multinomial. Here is a statement of this well-known fact.

THEOREM 2.4. *Suppose Θ is $\mathbf{D}(u)$ and, given Θ , X_1 has distribution Θ . Then, given $X_1 = i$, Θ is $\mathbf{D}(u + \delta(i))$, where $\delta(i)$ is the probability vector which has 1 in the i th coordinate.*

An elementary proof can be given using Bayes formula. A more interesting proof in the present context is to embed X_1 in a sequence $X = (X_1, X_2, \dots)$ of variables which are independent with distribution Θ given Θ . Then X is $\mathcal{P}(u)$ by Theorems 2.1 and 2.3. So, clearly, (X_2, X_3, \dots) is $\mathcal{P}(u + \delta(i))$ given $X_1 = i$. Use Theorems 2.1 and 2.3 again to see that Θ is $\mathbf{D}(u + \delta(i))$ given $X_1 = i$.

3. Finite Polya trees. Let s be a positive integer and let E_{s-1} be the set of all finite paths or sequences $p = (i_1, \dots, i_m)$ of elements of E whose length m is less than or equal to $s - 1$, including the empty sequence ϕ . A *Polya tree of height s* is a mapping u which assigns to each $p \in E_{s-1}$ an urn vector $u(p) = (u(p)_0, \dots, u(p)_k)$. Given such a u , the procedure explained in the introduction generates a random vector

$$X_1 = (X_{11}, \dots, X_{1s})$$

and a new tree $u^{(1)}$. [As before $X_{11} = i$ if the ball drawn from $u(\phi)$ is of color i , etc. However, this procedure, unlike the one in the introduction, ends after s stages.] The function $u^{(1)}$ is given by

$$u^{(1)}(i_1, \dots, i_m)_j = \begin{cases} u(i_1, \dots, i_m)_j + 1, & \text{if } X_{11} = i_1, \dots, X_{1m} = i_m, \\ & X_{1, m+1} = j, \\ u(i_1, \dots, i_m)_j, & \text{if not,} \end{cases}$$

for each $p = (i_1, \dots, i_m) \in E_{s-1}$ and $j \in E$.

Iterate the procedure using $u^{(1)}$ to get $X_2 = (X_{21}, \dots, X_{2s})$ and $u^{(2)}$, and so on.

The sequence $X = (X_1, X_2, \dots)$ is called *s-stage Polya with parameter u* or $\mathcal{P}_s(u)$.

To calculate the joint distribution of the first n variables, let

$$j = (j^{(1)}, \dots, j^{(n)})$$

be a sequence of n vectors in E^s :

$$j^{(a)} = (j_{a1}, \dots, j_{as}), \quad a = 1, \dots, n.$$

For each such j and each path p of length $r - 1$ in E_{s-1} , let $j(p)$ be the vector corresponding to the colors of those balls drawn from $u(p)$ if $(X_1, \dots, X_n) = j$. That is, if i is the number of draws made from $u(p)$, then

$$(3.0) \quad j(p) = (j_{a_1 r}, \dots, j_{a_r r}),$$

where $j^{(a_1)}, \dots, j^{(a_r)}$ are those vectors occurring in j (taken in order) whose first $r - 1$ coordinates coincide with p . Set $j(p) = \phi$ if the path p is not traversed. Now the probability of drawing $j(p)$ from $u(p)$ is just $P(j(p); u(p))$ as in (2.2). Furthermore

$$(3.1) \quad P[(X_1, \dots, X_n) = j] = \prod_{p \in E_{s-1}} P(j(p); u(p)),$$

where the convention is made that $P(\phi; u(p)) = 1$. One can verify (3.1) by writing the probability on the left as a product of the probabilities of the ns draws made and then grouping together the terms corresponding to the draws made from each $u(p)$.

LEMMA 3.1. *If (X_1, X_2, \dots) is $\mathcal{P}_s(u)$, then the variables X_1, X_2, \dots are exchangeable.*

PROOF. The probability $P(j(p); u(p))$ is invariant under permutations of the coordinates of $j(p)$ and, hence, is invariant under permutations of $j^{(1)}, \dots, j^{(n)}$. \square

Now Theorem 2.1 applies to the sequence X_1, X_2, \dots of exchangeable variables taking values in E^s , and we would like to identify the de Finetti measure. First some additional notation and terminology are needed.

It is convenient to view probability measures on E^s as being “strategies” in the sense of Dubins and Savage (1965).

DEFINITION. An s -day strategy is a mapping θ which assigns to every $p \in E_{s-1}$ a probability measure $\theta(p)$ on E . The notation θ_0 will often be used for $\theta(\phi)$.

An s -day strategy θ naturally determines a probability measure $\mu = \mu(\theta)$ on E^s as follows: The marginal μ distribution on the first coordinate is θ_0 and, given that the first r coordinates are p where $p \in E^r$, $1 \leq r \leq s$, the conditional μ distribution of the $(r + 1)$ st coordinate is $\theta(p)$.

To simplify notation, the measure $\mu(\theta)$ associated with θ will be written as θ below. In particular, the product rule for calculating the probability of an intersection gives, for $(i_1, \dots, i_s) \in E^s$,

$$(3.2) \quad \theta\{(i_1, \dots, i_s)\} = \theta_0(\{i_1\})\theta(i_1)(\{i_2\}) \cdots \theta(i_1, \dots, i_{s-1})(\{i_s\}).$$

If $\Theta_0, \Theta(i_1), \dots, \Theta(i_1, \dots, i_{s-1})$ are random probabilities on E , then we can use (3.2) with capital thetas to define a random probability Θ on E^s .

Let u be a Polya tree of height s .

DEFINITION. A random probability measure Θ on E^s is a *Dirichlet strategy with parameter u* [written $\mathbf{D}_s(u)$] if the random probability measures $\{\Theta(p) : p \in E_{s-1}\}$ are independent and $\Theta(p)$ is $\mathbf{D}(u(p))$ for every $p \in E_{s-1}$.

THEOREM 3.1. If $X = (X_1, X_2, \dots)$ is $\mathcal{P}_s(u)$, then the de Finetti measure of X is $\mathbf{D}_s(u)$.

PROOF. The proof is an application of Theorem 2.2. For the application, E is replaced in that theorem by E^s and,

$$j = (j^{(1)}, \dots, j^{(n)})$$

is a finite sequence of elements of E^s . We must verify (2.4) which here becomes

$$(3.3) \quad P[(X_1, \dots, X_n) = j] = E\left(\prod_{\alpha=1}^n \Theta(\{j^{(\alpha)}\})\right)$$

under the assumption that Θ is $\mathbf{D}_s(u)$. The left-hand side of (3.3) is given by (3.1). So it remains to calculate the right-hand side.

By (3.2), if θ is a strategy and

$$j^{(a)} = (j_{a1}, \dots, j_{as}),$$

then

$$\theta(\{j^{(a)}\}) = \theta_0\{j_{a1}\}\theta(j_{a1})\{j_{a2}\} \cdots \theta(j_{a1}, \dots, j_{a,s-1})\{j_{as}\}.$$

Now substitute into the right-hand side of (3.3), collect terms and use the independence of the $\Theta(p)$ to obtain

$$(3.4) \quad E\left(\prod_{a=1}^n \Theta(\{j^{(a)}\})\right) = \prod_{p \in E_{s-1}} E\left(\prod_{i \in j(p)} \Theta(p)(\{i\})\right).$$

Here $j(p)$ is the same as in (3.0) and i varies over the coordinates of $j(p)$ taken with their multiplicities. It follows from Theorems 2.2 and 2.3, and our assumption that $\Theta(p)$ is $\mathbf{D}(u(p))$ that

$$E\left(\prod_{i \in j(p)} \Theta(p)(\{i\})\right) = P(j(p); u(p)).$$

By (3.1) and (3.4), the proof is complete. \square

Just as the Dirichlet family is conjugate to ordinary multinomial sampling, the family of Dirichlet strategies is conjugate for “strategic sampling” in which an experiment takes place in several stages each of which depends on the preceding outcomes. Even if the experiment is terminated (censored) before the last stage, the Dirichlet strategies remain conjugate as was pointed out by Dickey (1990).

THEOREM 3.2. *Suppose Θ is $\mathbf{D}_s(u)$ and, given Θ , $X_1 = (X_{11}, \dots, X_{1s})$ has distribution Θ . Then, given $(X_{11}, \dots, X_{1r}) = (i_1, \dots, i_r)$ where $1 \leq r \leq s$, Θ is $\mathbf{D}_s(u')$ where*

$$\begin{aligned} u'(\phi) &= u(\phi) + \delta(i_1), \\ u'(i_1, \dots, i_a) &= u(i_1, \dots, i_a) + \delta(i_{a+1}) \quad \text{for } a = 1, \dots, r - 1, \\ u'(p) &= u(p) \quad \text{for all other } p \in E_{s-1}. \end{aligned}$$

PROOF. Do an induction on r using Theorem 2.4 and the independence of the $\Theta(p)$'s. [Alternatively, use Bayes formula and the fact that the density for Θ is the product of the densities for the $\Theta(p)$.] \square

4. Infinite Polya trees. An infinite Polya tree is a mapping u which assigns to every $p \in E^*$ an urn vector $u(p)$. Given such a u , the scheme described in the introduction generates sequences X_1, X_2, \dots and $u^{(1)}, u^{(2)}, \dots$ where, for each n ,

$$X_n = (X_{n1}, X_{n2}, \dots)$$

is a random element of E^N and $u^{(n)}$ is an infinite Polya tree. The sequence $X = (X_1, X_2, \dots)$ is infinite stage Polya with parameter u or $\mathcal{P}_\infty(u)$. We can

also code each X_n using a $(k + 1)$ -ary expansion as in (1.1) to get a random variable Y_n with values in the unit interval I . We will also call the sequence $Y = (Y_1, Y_2, \dots)$ *infinite stage Polya on I* or $\mathcal{P}_I(u)$.

The basic properties of the infinite stage Polya sequences are easily derived from the finite case. To do this, let $X = (X_1, X_2, \dots)$ be $\mathcal{P}_\infty(u)$ and, for positive integers s and n , let

$$X_n^{(s)} = (X_{n1}, \dots, X_{ns})$$

be the first s coordinates of X_n , let $X^{(s)}$ be the sequence $(X_1^{(s)}, X_2^{(s)}, \dots)$ and let $u^{(s)}$ be the restriction of u to E_{s-1} . Here is an obvious but useful fact.

LEMMA 4.1. *If X is $\mathcal{P}_\infty(u)$, then, for every s , $X^{(s)}$ is $\mathcal{P}_s(u^{(s)})$.*

THEOREM 4.1. *If (X_1, X_2, \dots) is $\mathcal{P}_\infty(u)$ [(Y_1, Y_2, \dots) is $\mathcal{P}_I(u)$], then X_1, X_2, \dots [Y_1, Y_2, \dots] are exchangeable.*

PROOF. Let n be a positive integer and let A be a Borel subset of $(E^N)^n$. To prove exchangeability of the X_i 's, we need to check that $P[(X_1, \dots, X_n) \in A]$ is invariant under permutations of the indices. It suffices to do this for sets A of the form

$$A = A_1 \times \dots \times A_n,$$

where each A_i is a cylinder set in E^N of the form

$$A_i = B_i \times E^{r_i}, \quad B_i \subset E^{r_i}$$

for some positive integer r_i . Take s to be the maximum of the r_i 's so that each A_i depends on only the first s coordinates and is of the form

$$A_i = C_i \times E^{N-s}, \quad C_i \subset E^s.$$

Thus

$$P[X_1 \in A_1, \dots, X_n \in A_n] = P[X_1^{(s)} \in C_1, \dots, X_n^{(s)} \in C_n].$$

Exchangeability of the X_i 's now follows from that of the $X_i^{(s)}$'s. The exchangeability of the Y_i 's is an easy consequence of that of the X_i 's. \square

A more general form of Theorem 2.1 [cf. Hewitt and Savage (1955) or Aldous (1983)] can now be used to see that there is a de Finetti measure for a sequence which is $\mathcal{P}_\infty(u)$ or $\mathcal{P}_I(u)$. To describe these measures, let $M(E^N)$ [$M(I)$] be the space of probability measures defined on the Borel subsets of E^N [I] and give this space its usual topology of weak convergence. Let X be $\mathcal{P}_\infty(u)$. Then the *de Finetti measure* $Q = Q_u$ for X is a probability measure defined on the Borel subsets of $M(E^N)$ and satisfying

$$(4.1) \quad P[X_1 \in A_1, \dots, X_n \in A_n] = \int \left(\prod_{i=1}^n \theta(A_i) \right) Q(d\theta)$$

for all n and all Borel subsets A_1, \dots, A_n of E^N . The de Finetti measure $\bar{Q} = \bar{Q}_u$ for a Y which is $\mathcal{P}_I(u)$ can be defined similarly; just replace the X_i 's

by Y_i 's and take the sets A_i to be Borel subsets of the unit interval. The existence and uniqueness of Q and \bar{Q} are well known [cf. Hewitt and Savage (1955) or Aldous (1983)]. There is, of course, a simple relationship between Q and \bar{Q} .

Let $\psi: E^N \rightarrow I$ be the mapping defined by (1.1). Clearly, ψ is continuous and induces a continuous mapping $\theta \rightarrow \theta\psi^{-1}$ from $M(E^N)$ into $M(I)$ where $(\theta\psi^{-1})(B) = \theta(\psi^{-1}(B))$ for B a Borel subset of I . Now \bar{Q} can be thought of as the distribution of $\theta\psi^{-1}$ when Θ has distribution Q ; that is,

$$(4.2) \quad \bar{Q}(F) = Q\{\theta: \theta\psi^{-1} \in F\}$$

for Borel subsets F of $M(I)$.

To characterize Q , it is again useful to use the notion of a strategy.

DEFINITION. A strategy θ is a mapping from E^* to the collection of probability measures on E .

Each strategy θ naturally determines a measure $\mu = \mu(\theta) \in M(E^N)$ by the requirement that $\mu(\theta)\{x \in E^N: x_1 = i_1, \dots, x_s = i_s\}$ be equal to the right-hand side of (3.2) for every finite sequence i_1, \dots, i_s of elements of E . As in the s -day case, we will write θ for $\mu(\theta)$ and we can obtain a random measure Θ with values in $M(E^N)$ by putting a joint distribution on $\{\Theta(p): p \in E^*\}$.

Let u be an infinite Polya tree.

DEFINITION. A random probability measure Θ with values in $M(E^N)$ is a *Dirichlet strategy with parameter u* [written $\mathbf{D}_\infty(u)$] if the random probability measures $\{\Theta(p): p \in E^*\}$ are independent and $\Theta(p)$ is $\mathbf{D}(u(p))$ for every $p \in E^*$. The measure corresponding to the distribution of such a Θ is also said to be $\mathbf{D}_\infty(u)$.

THEOREM 4.2. If $X = (X_1, X_2, \dots)$ is $\mathcal{P}_\infty(u)$, then the de Finetti measure Q_u of X is $\mathbf{D}_\infty(u)$.

PROOF. If Θ is $\mathbf{D}_\infty(u)$, then, for every positive integer s , Θ restricted to E_{s-1} is $\mathbf{D}_s(u)$. So, by Theorem 3.1, equality (4.1) holds when every A_i depends only on the first s coordinates. \square

Again the Dirichlet strategies are conjugate for strategic sampling even if the experiment is terminated at some finite stage.

THEOREM 4.3. Suppose Θ is $\mathbf{D}_\infty(u)$ and, given Θ , $X_1 = (X_{11}, X_{12}, \dots)$ has distribution Θ . Then, given X_1 [or given (X_{11}, \dots, X_{1r})], Θ is $\mathbf{D}_\infty(u^{(1)})$ where

$$\begin{aligned} u^{(1)}(\phi) &= u(\phi) + \delta(X_{11}), \\ u^{(1)}(X_{11}, \dots, X_{1r}) &= u(X_{11}, \dots, X_{1r}) + \delta(X_{1,r+1}) \quad \text{for } r \leq s - 1, \\ u^{(1)}(p) &= u(p) \quad \text{for other } p \in E^*. \end{aligned}$$

PROOF. In checking the properties of a conditional distribution, one can restrict attention to Θ restricted to the finite sets E_s and to finitely many of the X_{1n} 's. But then the desired properties follow from Theorem 3.2. \square

Another way of expressing the result of Theorem 4.3 is that if Θ has the prior distribution Q_u on $M(E^N)$, then, given X_1 , the posterior distribution of Θ is $Q_{u^{(1)}}$ where $u^{(1)}$ is the tree obtained in the construction of Section 1. It follows that the posterior of Θ given X_1, \dots, X_n is $Q_{u^{(n)}}$. (Notice that $u^{(n)}$ is a random tree depending on the values of X_1, \dots, X_n .)

Suppose next that Θ is a random element of $M(I)$ with distribution \bar{Q}_u and we wish to calculate the posterior distribution of Θ given Y_1 where $Y_1 = \psi(X_1)$. It follows from (1.1) and (4.2) that this posterior distribution will be $\bar{Q}_{u^{(1)}}$ if we make the convention that $u^{(1)}$ should be the tree associated with an X_1 such that $\psi(X_1) = Y_1$ and having only finitely many 0's (say) and if the probability under \bar{Q}_u that Y_1 is a $(k + 1)$ -ary rational is zero. This last condition is certainly satisfied if Y_1 has a continuous distribution in the sense of the next section.

5. Continuity of predictive distributions and random measures. A prior Q [respectively \bar{Q}] on $M(E^N)$ [$M(I)$] is a probability measure defined on the Borel subsets of $M(E^N)$ [$M(I)$]. Suppose the random measure Θ has distribution Q [\bar{Q}] and, given Θ , the variables X_1, X_2, \dots are independent each having distribution Θ . The marginal distribution of X_1 is called the *predictive distribution* for the prior Q [\bar{Q}].

The collection of *continuous measures* on $E^N[I]$ is given by

$$C = \{ \theta \in M(E^N) : \theta\{x\} = 0 \text{ for all } x \in E^N \}$$

$$[\bar{C}] = \{ \theta \in M(I) : \theta\{x\} = 0 \text{ for all } x \in I \}.$$

We are interested in conditions on an infinite Polya tree u for the prior Q_u [\bar{Q}_u] to have a continuous predictive distribution and also for $Q_u(C)$ [$\bar{Q}_u(\bar{C})$] to be 1. For convenience, we will work on E^N , but the conditions given apply as well on I for the prior \bar{Q}_u .

Now, under Q_u , the probability that X_1 equals $x \in E^N$ is just the probability, in the Polya tree construction, of traversing the path $x = (x_1, x_2, \dots)$ and this probability is given by the infinite product

$$(5.1) \quad \Pi(x; u) = \frac{u(\phi)_{x_1}}{|u(\phi)|} \cdot \frac{u(x_1)_{x_2}}{|u(x_1)|} \cdot \frac{u(x_1, x_2)_{x_3}}{|u(x_1, x_2)|} \cdots$$

Here is an immediate consequence and an obvious corollary.

THEOREM 5.1. *The predictive distribution of Q_u (\bar{Q}_u) is continuous if and only if $\Pi(x; u) = 0$ for every $x \in E^N$.*

COROLLARY 5.1. *If the proportion of balls of each color is bounded away from 1 in the sense that*

$$(5.2) \quad \sup \left\{ \frac{u(p)_i}{|u(p)|} : p \in E^*, i \in E \right\} < 1,$$

then the predictive distribution of Q_u (\bar{Q}_u) is continuous.

In the special case when every $u(p)_i$ is an integer, condition (5.2) is satisfied if the total number of balls in each urn is uniformly bounded and if there are balls of different colors in every urn.

The next two lemmas hold for a general prior Q [\bar{Q}] on $M(E^N)$ [$M(I)$].

LEMMA 5.1. *If $Q(C)$ [$\bar{Q}(\bar{C})$] equals 1, then the predictive distribution of Q [\bar{Q}] is continuous.*

PROOF. If $Q(C) = 1$ and $x \in E^N$, then

$$P[X_1 = x] = \int P[X_1 = x | \Theta = \theta] dQ(\theta) = \int \theta(x) dQ(\theta) = 0. \quad \square$$

LEMMA 5.2. *A necessary and sufficient condition for $Q(C)$ [$\bar{Q}(\bar{C})$] to be 1 is that $P[X_1 = X_2]$ be zero.*

PROOF. Now X_1 and X_2 are independent with distribution θ given $\Theta = \theta$. Hence, $P[X_1 = X_2 | \Theta = \theta]$ is zero if and only if θ is continuous. Consequently,

$$P[X_1 = X_2] = \int_{C^c} P[X_1 = X_2 | \Theta = \theta] dQ(\theta) > 0$$

if and only if $Q(C^c) > 0$. \square

To apply Lemma 5.2 to the case where $Q = Q_u$ for a Polya tree u , notice that, given $X_1 = x$, the distribution of X_2 is the predictive distribution for $Q_{u^{(1)}}$ by Theorem 4.3. Thus

$$(5.3) \quad P[X_2 = x | X_1 = x] = \Pi(x; u^{(1)}).$$

In the expression on the right, the tree $u^{(1)}$ is from the construction in Section 1 and, in particular, depends on x . Also, if m is the distribution of X_1 , then

$$(5.4) \quad P[X_1 = X_2] = \int P[X_2 = x | X_1 = x] dm(x).$$

Now we are ready to give conditions for $Q_u(C)$ to be 1.

THEOREM 5.2. *If $Q_u(C)$ [$\bar{Q}_u(\bar{C})$] equals 1, then $\Pi(x; u) = 0$ for every $x \in E^N$. If $\Pi(x; u^{(1)}) = 0$ for m almost every $x \in E^N$, then $Q_u(C)$ [$\bar{Q}_u(\bar{C})$] equals 1.*

PROOF. The first assertion is immediate from Theorem 5.1 and Lemma 5.1. The second follows from (5.3), (5.4) and Lemma 5.2. \square

The trees $u^{(1)}$ obtained in the Polya construction of Section 1 all have the property that every urn $u^{(1)}(p)$ differs from $u(p)$ at most by the addition of one ball. More precisely, every $u^{(1)}(p)_i$ is either $u(p)_i + 1$ or $u(p)_i$ depending on whether $p = (x_1, \dots, x_n)$ and $x_{n+1} = i$ for some n or not. Similarly every $|u^{(1)}(p)|$ is either $|u(p)| + 1$ or $|u(p)|$ according to whether $p = (x_1, \dots, x_n)$ for some n or not. It follows that the terms occurring in the infinite product expression (5.1) for $\Pi(x; u^{(1)})$ will be bounded away from 1 if (5.2) holds and the $|u(p)|$ are bounded away from zero. This observation yields a corollary to the second assertion of Theorem 5.2.

COROLLARY 5.2. *If (5.2) holds and*

$$(5.5) \quad \inf\{|u(p)| : p \in E^*\} > 0,$$

then $Q_u(C) [\bar{Q}_u(\bar{C})]$ is 1.

Condition (5.5) obviously holds when every $u(p)_i$ is an integer because in that case $|u(p)| \geq 1$. (We are not allowing empty urns.)

6. Support and consistency. As Ferguson (1973) remarked, it is often desirable for a prior to have large support. The *support* (or *topological carrier*) of a probability measure μ defined on the Borel subsets of a compact Hausdorff space M is the least compact set $S(\mu)$ which has μ measure 1. Notice that μ has *full support* in the sense that $S(\mu) = M$ if and only if every nonempty, open subset of M has positive μ measure.

Here is a characterization of the Polya tree priors which have full support.

THEOREM 6.1. *The following are equivalent conditions on an infinite Polya tree u :*

- (a) *The prior Q_u has full support.*
- (b) *The prior \bar{Q}_u has full support.*
- (c) *For every $p \in E^*$, the Dirichlet measure $\mathbf{D}(u(p))$ has full support.*
- (d) *For every $p \in E^*$ and $i \in E$, $u(p)_i > 0$.*

PROOF. (a) \Rightarrow (b). Assume (a) and let F be a nonempty, open subset of $M(I)$. Then the set $G = \{\theta : \theta\psi^{-1} \in F\}$ is a nonempty, open subset of $M(E^N)$ and, by (4.3),

$$\bar{Q}(F) = Q(G) > 0.$$

(Here and below $\bar{Q} = \bar{Q}_u$ and $Q = Q_u$.)

(b) \Rightarrow (d). Suppose (d) is false. So there exist $p = (i_1, \dots, i_m) \in E^*$ and $i \in E$ such that $u(p)_i = 0$. Thus, if Θ is a random measure with distribution Q_u , then $\Theta(p)\{i\} = 0$ with probability 1.

Let l and r be the numbers whose expansions to base $k + 1$ are

$$l = .i_1 \dots i_m i 00 \dots,$$

$$r = .i_1 \dots i_m i k k \dots,$$

and let J be the closed interval $[l, r]$. Let $g: I \rightarrow [0, \infty)$ be a continuous, non-zero function which equals zero on the complement of J . Consider the nonempty, open set

$$U = \left\{ \mu \in M(I) : \int g d\mu > 0 \right\}.$$

To prove (b) is false, we need only show $\overline{Q}(U) = 0$.

Notice that U is a subset of the set

$$F = \{ \mu \in M(I) : \mu(J) > 0 \}.$$

So it suffices to show $\overline{Q}(F) = 0$. By (4.3),

$$\overline{Q}(F) = Q\{ \theta : \theta(\psi^{-1}(J)) > 0 \}.$$

Also,

$$\psi^{-1}(J) = \{ x \in E^N : x_1 = i_1, \dots, x_m = i_m, x_{m+1} = i \}$$

and

$$\begin{aligned} \theta(\psi^{-1}(J)) &= \theta_0\{i_1\}\theta(i_1)\{i_2\} \cdots \theta(i_1, \dots, i_m)\{i\} \\ &\leq \theta(i_1, \dots, i_m)\{i\} = \theta(p)\{i\}. \end{aligned}$$

Hence,

$$\overline{Q}(F) \leq Q\{ \theta : \theta(p)\{i\} > 0 \} = 0.$$

(c) \Leftrightarrow (d). This is a trivial consequence of our conventions about the Dirichlet distribution. [Notice that to say $\mathbf{D}(u(p))$ has full support means that the support of $\mathbf{D}(u(p))$ is S_k .]

(d) \Rightarrow (a). Assume (d). It suffices to show that each set in a base for the topology of $M(E^N)$ has positive Q measure. The usual base for the weak topology consists of sets of the form

$$\left\{ \theta \in M(E^N) : \left| \int g_i d\theta - \int g_i d\theta_0 \right| < \varepsilon_i, i = 1, \dots, n \right\},$$

where the g_i are continuous, real-valued functions on E^N , θ_0 is a fixed element of $M(E^N)$ and the ε_i are positive numbers [cf. Section II.6 of Parthasarathy (1967)]. However, every continuous g on E^N can be uniformly approximated by a finite, linear combination of indicator functions of sets of the form

$$(6.1) \quad C = \{ x \in E^N : x_1 = i_1, \dots, x_n = i_n \}.$$

(This follows from the Stone-Weierstrass theorem, for example.) Hence, another base for the topology of $M(E^N)$ consists of sets of the form

$$(6.2) \quad B = \{ \theta \in M(E^N) : |\theta(C_i) - \theta_0(C_i)| < \varepsilon_i, i = 1, \dots, n \},$$

where each C_i is a set of the form (6.1), $\theta_0 \in M(E^N)$ and each ε_i is positive.

So we need only show $Q(B)$ is positive for B as in (6.2). To do this, we begin by borrowing a trick from Ferguson [(1973), Proposition 3.3]. Call n the *dimension* of the set C in (6.1) and let b be the maximum of the dimensions of the C_i occurring in (6.2). Then each C_i is a disjoint union of at most 2^b sets of the form

$$(6.3) \quad D = \{x \in E^N: x_1 = j_1, \dots, x_b = j_b\}.$$

Let ε be the minimum of the ε_i in (6.2). Then B contains the set

$$(6.4) \quad F = \bigcap_D \{\theta \in M(E^N): |\theta(D) - \theta_0(D)| < \varepsilon/2^b\},$$

where the intersection is over all D of the form (6.3). So it suffices to show that $Q(F) > 0$.

Now each D occurring in (6.4) is of dimension b . So a strategy θ belongs to F if and only if its restriction $\theta^{(b)}$ to E^{b-1} belongs to

$$F_b = \bigcap_{D_b} \{\theta \in M(E^b): |\theta(D_b) - \theta_0(D_b)| < \varepsilon/2^b\},$$

where D_b is that subset of E^b such that

$$D = D_b \times E^N.$$

The set F_b is a nonempty, open subset of $M(E^b)$, and if Θ has distribution Q , then

$$Q(F) = P[\Theta \in F] = P[\Theta^{(b)} \in F_b].$$

The final probability is positive because $\Theta^{(b)}$ is $\mathbf{D}_b(u^{(b)})$ and has, under condition (d), a density which is positive on all of $M(E^b)$. [The space $M(E^b)$ can be identified with the set S_n where $n = k^b - 1$ and the density of $\Theta^{(b)}$ is taken with respect to Lebesgue measure.] \square

Let $\lambda \in M(E^N)$ and let u be an infinite Polya tree. Suppose data variables X_1, X_2, \dots are independent with distribution λ and consider the posterior distributions $Q_{u^{(n)}}$ calculated from these data variables. Following Freedman (1963) and Fabius (1964), call the pair (λ, Q_u) *consistent* if, with probability 1, $Q_{u^{(n)}}$ converges weakly to $\delta(\lambda)$, the measure concentrated on $\{\lambda\}$. For $\lambda \in M(I)$, the consistency of the pair (λ, \bar{Q}_u) is defined similarly under the assumption that λ gives mass zero to the collection of $k + 1$ -ary rationals so that $\bar{Q}_{u^{(n)}}$ is the correct posterior given data Y_1, \dots, Y_n which are independent with distribution λ .

THEOREM 6.2. *If λ belongs to the support of Q_u [\bar{Q}_u and assigns mass zero to the $(k + 1)$ -ary rationals], then the pair (λ, Q_u) [(λ, \bar{Q}_u)] is consistent.*

The proof of this theorem is similar to the proof of Theorem 2.2 in Fabius (1964), the main idea being a reduction to the finite, discrete case of Freedman (1963). Indeed, the measures \bar{Q}_u are "tail-free" in the sense of Fabius so that this result is almost immediate from his.

7. Estimation of a distribution function. Suppose Y_1, \dots, Y_n is a sample from a distribution $\Theta \in M(I)$ and we wish to estimate the distribution function F for Θ given by

$$F(y) = \Theta[0, y], \quad 0 \leq y \leq 1.$$

If Θ has prior distribution \bar{Q}_u , then, as Ferguson [(1973), Section 5 (a)] explains, a natural Bayes estimator is the distribution function $\hat{F}_n(y|Y_1, \dots, Y_n)$ which corresponds to the expected value of $F(y)$ under the posterior $\bar{Q}_{u^{(n)}}$ and can be written as

$$\hat{F}_n(y|Y_1, \dots, Y_n) = \int \theta[0, y] \bar{Q}_{u^{(n)}}(d\theta).$$

Since we know the form of the Polya tree $u^{(n)}$, the problem reduces to the no data case where F is the expected (or predictive) distribution function as in

$$\hat{F}(y) = \hat{F}(y; u) = \int \theta[0, y] \bar{Q}_u(d\theta).$$

For the calculation, introduce the notation $\Pi((y_1, \dots, y_n); u)$ for the probability that the Polya tree process X_{11}, \dots, X_{1n} traverses the finite path (y_1, \dots, y_n) ; that is,

$$(7.0) \quad \Pi((y_1, \dots, y_n); u) = \frac{u(\phi)_{y_1}}{|u(\phi)|} \dots \frac{u(y_1, \dots, y_{n-1})_{y_n}}{|u(y_1, \dots, y_{n-1})|}.$$

The distribution function $\hat{F}(y; u)$ is, of course, the distribution function for predictive distribution discussed in Section 5. That is, $\hat{F}(y; u)$ is the distribution function for the random variable

$$(7.1) \quad Y = .Y_1Y_2\dots \quad (\text{to base } k + 1),$$

where $(Y_1, Y_2, \dots) = (X_{11}, X_{12}, \dots)$ is the random sequence constructed in the infinite Polya urn scheme of the Introduction.

THEOREM 7.1. *The expected distribution function under the Polya tree prior \bar{Q}_u is*

$$(7.2) \quad \hat{F}(y; u) = \sum_{n=0}^{\infty} \left(\Pi((y_1, \dots, y_n); u) \sum_{i=0}^{y_{n+1}-1} \frac{u(y_1, \dots, y_n)_i}{|u(y_1, \dots, y_n)|} \right) + \Pi((y_1, y_2, \dots); u)$$

for every $y = .y_1y_2\dots$ (to base $k + 1$) in I . [Here $\Pi(\phi; u) = 1$ and $\Pi((y_1, y_2, \dots); u)$ is as in (5.1). Also, the inner summation is taken to be zero when $y_{n+1} = 0$ and we take the representation of y in which infinitely many y_i 's are zero if there is ambiguity.]

PROOF. Observe that

$$P[Y \leq y] = \sum_{n=0}^{\infty} P[(Y_1, \dots, Y_n) = (y_1, \dots, y_n), Y_{n+1} < y_{n+1}] + P[Y = y],$$

which by inspection is (7.2). \square

Formula (7.2) is annoyingly complex when compared with Ferguson's (1973) formula (5.2). However, it does simplify in interesting special cases.

Say that an infinite Polya tree u has *constant proportions* λ if there is a fixed probability vector $\lambda = (\lambda_0, \dots, \lambda_k)$ such that, for every $p \in E^*$ and $i \in E$,

$$\frac{u(p)_i}{|u(p)|} = \lambda_i.$$

Let $\lambda^* = (k + 1)^{-1}(1, 1, \dots, 1)$ be the probability vector all of whose coordinates are equal to $(k + 1)^{-1}$.

THEOREM 7.2. *Suppose u has constant proportions λ .*

- (a) *If $\lambda = \lambda^*$, then $\hat{F}(y; u) = y$ for all $y \in I$.*
- (b) *If $\lambda \neq \lambda^*$, then $\hat{F}(y; u)$ is singular with respect to Lebesgue measure.*

PROOF. If $\lambda = \lambda^*$, then, by (7.2),

$$\hat{F}(y; u) = \sum_{n=0}^{\infty} (k + 1)^{-n} \frac{y_{n+1}}{k + 1} = \sum_{n=1}^{\infty} y_n (k + 1)^{-n} = y.$$

Suppose now that $\lambda \neq \lambda^*$. So, for some $i \in E$, $\lambda_i \neq (k + 1)^{-1}$. Now the variables X_{11}, X_{12}, \dots of (7.1) are clearly independent with distribution λ . By the strong law of large numbers $\hat{F}(\cdot; u)$ assigns probability 1 to the set of all $y = .y_1 y_2 \dots$ such that

$$\#\{j \leq n : y_j = i\} / n \rightarrow \lambda_i \quad \text{as } n \rightarrow \infty.$$

But this set has Lebesgue measure zero. \square

Consider again the problem of calculating $\hat{F}_n(y | Y_1, \dots, Y_n) = \hat{F}(y; u^{(n)})$. In principle, the problem is solved since we know the form of $u^{(n)}$ and can apply (7.2). In practice, it may be difficult to evaluate the infinite sum of (7.2). Lavine (1992) explains how to calculate the density for Y when it exists and gives an example to illustrate how it depends on the choice of the Polya tree prior.

8. Estimation of a mean. As explained in Ferguson (1973), if the statistician is to estimate the mean of a distribution $\Theta \in M(I)$ with squared error loss, then the Bayes estimate is the expected mean of the posterior. So, if Θ

has prior distribution \bar{Q}_u , the Bayes estimate will be

$$(8.1) \quad \int m(\theta)\bar{Q}_{u^{(n)}}(d\theta),$$

where $m(\theta) = \int x\theta(dx)$ and $\bar{Q}_{u^{(n)}}$ is the posterior. Again the problem reduces, at least in principle, to the no data case for which a formula is given below.

Let u be a Polya tree and, for each $p \in E^*$, define $m(u(p))$ to be the mean of the variable corresponding to one draw from $u(p)$; that is,

$$m(u(p)) = \sum_{i=0}^k \frac{i u(p)_i}{|u(p)|}.$$

THEOREM 8.1. *The expected mean under the Polya tree prior \bar{Q}_u is*

$$(8.2) \quad \int m(\theta)\bar{Q}_u(d\theta) = \sum_{n=1}^{\infty} (k+1)^{-n} \sum_{p \in E^{n-1}} \Pi(p; u) m(u(p)),$$

where, for each $p = (i_1, \dots, i_n)$, $\Pi(p, u)$ denotes, as in (7.0), the probability that the Polya tree process traverses p and $\Pi(\phi; u) = 1$.

PROOF. For Y as in (7.1),

$$\begin{aligned} E[Y] &= E[.Y_1 Y_2 \dots] \\ &= E[.Y_1] + E[.0Y_2] + E[.00Y_3] + \dots \\ &= E[.Y_1] + E[E[.0Y_2 | Y_1]] + E[E[.00Y_3 | Y_1, Y_2]] + \dots \end{aligned}$$

and this final expression equals the right-hand side of (8.2). \square

If there is a constant m_0 such that $m(u(p)) = m_0$ for every $p \in E^*$, then (8.2) becomes

$$m_0 \sum_{n=1}^{\infty} (k+1)^{-n} \sum_{p \in E^{n-1}} \Pi(p; u) = m_0 \sum_{n=1}^{\infty} (k+1)^{-n} = m_0/k.$$

In the special case when every urn $u(p)$ has equal proportions of all $k+1$ colors,

$$m_0 = \sum_{i=0}^k i/(k+1) = k/2,$$

so that the expected mean is $1/2$.

Consider finally the problem of calculating the posterior mean in (8.1) based on observations Y_1, \dots, Y_n where

$$Y_i = .X_{i1} X_{i2} \dots \quad (\text{to base } k+1)$$

is a unique representation for each i . A direct calculation from (8.2) is sometimes feasible if one takes advantage of the formula

$$m(u^{(n)}(p)) = \frac{|u(p)|}{|u(p)| + n(p)} m(u(p)) + \frac{n(p)}{|u(p)| + n(p)} \bar{y}(p),$$

where, for each $p \in E^*$, $n(p)$ is the number of draws from $u(p)$ and $\bar{y}(p)$ is the mean of the numbers drawn.

Acknowledgments. Thanks to Tom Ferguson, Michael Lavine, a referee and an Associate Editor for their helpful comments.

REFERENCES

- ALDOUS, D. J. (1983). Exchangeability and related topics. *Ecole d'Été de Probabilités de Saint-Flour XIII. Lecture Notes in Math.* **1117** 1–197. Springer, New York.
- BLACKWELL, D. and KENDALL, D. (1964). The Martin boundary for Polya's urn scheme, and an application to stochastic population growth. *J. Appl. Probab.* **1** 284–296.
- BLACKWELL, D. and MACQUEEN, J. B. (1973). Ferguson distributions via Polya urn schemes. *Ann. Statist.* **1** 353–355.
- DICKEY, J. (1990). Personal communication.
- DUBINS, L. E. and FREEDMAN, D. A. (1967). Random distribution functions. *Proc. Fifth Berkeley Symp. Math. Statist. Probab.* **2** 183–214. Univ. California Press, Berkeley.
- DUBINS, L. E. and SAVAGE, L. J. (1965, 1976). *Inequalities for Stochastic Processes*. Dover, New York.
- FABIUS, J. (1964). Asymptotic behavior of Bayes estimates. *Ann. Math. Statist.* **35** 846–856.
- FERGUSON, T. S. (1973). A Bayesian analysis of some nonparametric problems. *Ann. Statist.* **1** 209–230.
- FERGUSON, T. S. (1974). Prior distributions on spaces of probability measures. *Ann. Statist.* **2** 615–629.
- FREEDMAN, D. A. (1963). On the asymptotic behavior of Bayes' estimates in the discrete case. *Ann. Math. Statist.* **34** 1386–1403.
- GRAF, S., MAULDIN, R. D. and WILLIAMS, S. C. (1986). Random homeomorphisms. *Adv. in Math.* **60** 239–359.
- HEWITT, E. and SAVAGE, L. J. (1955). Symmetric measures on Cartesian products. *Trans. Amer. Math. Soc.* **80** 470–501.
- KRAFT, C. H. (1964). A class of distribution function processes which have derivatives. *J. Appl. Probab.* **1** 385–388.
- LAVINE, M. (1992). Some aspects of Polya tree distributions for statistical modelling. *Ann. Statist.* **20** 1222–1235.
- MAULDIN, R. D. and WILLIAMS, S. C. (1990). Reinforced random walks and random distributions. *Proc. Amer. Math. Soc.* **110** 251–258.
- MEYER, P. A. (1966). *Probability and Potentials*. Blaisdell, Waltham, Mass.
- PARTHASARATHY, K. R. (1967). *Probability Measures on Metric Spaces*. Academic, New York.

R. DANIEL MAULDIN
DEPARTMENT OF MATHEMATICS
BOX 5116
UNIVERSITY OF NORTH TEXAS
DENTON, TEXAS 76203

WILLIAM D. SUDDERTH
DEPARTMENT OF THEORETICAL STATISTICS
270 VINCENT HALL
UNIVERSITY OF MINNESOTA
206 CHURCH STREET S.E.
MINNEAPOLIS, MINNESOTA 55455

S. C. WILLIAMS
DEPARTMENT OF MATHEMATICS
AND STATISTICS
UTAH STATE UNIVERSITY
LOGAN, UTAH 84322