

THE SIZE OF THE CLASS OF COUNTABLE SEQUENCES OF ORDINALS

WILLIAM CHAN, STEPHEN JACKSON, AND NAM TRANG

ABSTRACT. Assume $\text{ZF} + \text{AD} + \text{DC}_{\mathbb{R}}$. There is no injection of ${}^{<\omega_1}\omega_1$ (the set of countable length sequences of countable ordinals) into ${}^\omega\text{ON}$ (the class of ω length sequences of ordinals). There is no injection of $[\omega_1]^{\omega_1}$ (the powerset of ω_1) into ${}^{<\omega_1}\text{ON}$ (the class of countable length sequences of ordinals).

1. INTRODUCTION

Mathematical size between two sets is compared through injections and bijections. If A and B are two sets, then $|A| \leq |B|$ indicates that there is an injection from A to B . One writes $|A| < |B|$ if and only if $|A| \leq |B|$ and not $|B| \leq |A|$. One writes $|A| = |B|$ if there is a bijection between A and B . In ZF , the Cantor-Schröder-Bernstein theorem asserts that $|A| \leq |B|$ and $|B| \leq |A|$ imply that $|A| = |B|$. The axiom of choice, AC , implies every set can be wellordered and is in bijection with an ordinal and the least such one is called its cardinality. Thus under AC , the class of cardinals is wellordered under the injection relation. A frequent phenomenon is that the relations between the size of two sets with explicit definitions are often independent of $\text{ZF} + \text{AC}$. The continuum hypothesis is a notable example.

The axiom of determinacy, AD , asserts that all integer games between two players of a certain form have a winning strategy for one of the two players. Cardinalities of sets are no longer wellorderable under injections. However, under AD and its extensions, mathematical size of sets become more natural in that size corresponds more closely to the identity of the object or its fundamental combinatorial properties. The computation of size under determinacy involves techniques that are closely connected to definability. For instance, under AD , $|\mathbb{R}|$ and ω_1 are incomparable cardinalities. Even under AC , it seems that one cannot explicitly specify a wellordering of the reals without imposing conditions on the structure of the universe, such as the reals all belong to the constructible universe L or some other canonical inner model. Moreover, various large cardinal principles imply that wellordering of the reals must necessarily be quite complicated. Under AD , the incomparability of $|\mathbb{R}|$ and ω_1 follows from the measurability of ω_1 . Measurability of ω_1 can be proved using the Martin measure on the Turing degrees. Alternatively, Solovay showed the club measure on ω_1 is a normal measure using the Σ_1^1 boundedness principle.

Let ON denote the class of ordinals. Let $\epsilon \in \text{ON}$ and $X \subseteq \text{ON}$ be a set or class. Let ${}^\epsilon X$ be the set of functions $f : \epsilon \rightarrow X$. Let ${}^{<\epsilon}X = \bigcup_{\delta < \epsilon} {}^\delta X$. Let $[X]^\epsilon$ be the set of functions $f : \epsilon \rightarrow X$ which are increasing. Let $[X]^{<\epsilon} = \bigcup_{\delta < \epsilon} [X]^\delta$.

Suppose κ and δ are two ordinals greater than 1. The main question in this paper is whether ${}^{<\omega_1}\kappa$ can inject into ${}^\omega\delta$. These two sets seem to have a fundamental difference. ${}^{<\omega_1}\kappa$ consists of sequences of arbitrary countable length and ${}^\omega\delta$ consists entirely of sequences of one fixed length ω . Assuming the axiom of choice, these sets may not be distinguishable through size since, for example, $||[\omega_1]^\omega| = |[\omega_1]^{<\omega_1}|$ under ZFC .

Observe that $|{}^{<\omega_1}2| = |[\omega_1]^{<\omega_1}| = |{}^{<\omega_1}\omega_1|$. Thus a negative answer to the above question follows from a negative answer to the following question.

Question 1.1. Is there an injection from $[\omega_1]^{<\omega_1}$ into ${}^\omega\text{ON}$?

Another related question is that if κ and δ are two ordinals greater than 1, then does $|{}^{\omega_1}\kappa| \leq |{}^{<\omega_1}\delta|$ hold? Again these two sets have a fundamental difference: ${}^{\omega_1}\kappa$ consists of sequences of length ω_1 and ${}^{<\omega_1}\delta$ consists of countable length sequences. Observe that $|{}^{\omega_1}2| = |\mathcal{P}(\omega_1)| = |[\omega_1]^{\omega_1}|$. Thus a negative answer to the above question follows from a negative answer to the following question.

Question 1.2. Is there an injection from $[\omega_1]^{\omega_1}$ into ${}^{<\omega_1}\text{ON}$?

January 22, 2021. The first author was supported by NSF grant DMS-1703708. The second author was supported by NSF grant DMS-1800323. The third author was supported by NSF grant DMS-1855757 and DMS-1945592.

This paper will work in extensions of AD. A very useful extension of AD is a theory called AD^+ isolated by Woodin ([12] Section 9.1). AD^+ consists of the following statements.

- $\text{DC}_{\mathbb{R}}$.
- Every set of reals has an ∞ -Borel code. (An ∞ -Borel code is a pair (S, φ) where S is a set of ordinals and φ is a formula of set theory. Let $\mathfrak{B}_{(S, \varphi)} = \{r \in \mathbb{R} : L[S, r] \models \varphi(S, r)\}$. (S, φ) is an ∞ -Borel code for a set $A \subseteq \mathbb{R}$ if and only if $A = \mathfrak{B}_{(S, \varphi)}$.)
- **Ordinal Determinacy**, which is the statements that for every $\lambda < \Theta$, $X \subseteq \mathbb{R}$, and continuous function $\pi : {}^\omega \lambda \rightarrow \mathbb{R}$, the two player game on λ with payoff set $\pi^{-1}(X)$ is determined.

Results of Kechris and Woodin showed that if AD holds, then $L(\mathbb{R}) \models \text{AD}^+$. The relation between AD and AD^+ as well as the relations between the three statements in AD^+ are not known.

Woodin [11] was aware of a negative answer to a particular instance of Question 1.1 under $\text{ZF} + \text{DC} + \text{AD}_{\mathbb{R}}$. This involves investigating the cardinality of a set called $S_1 = \{\sigma \in [\omega_1]^{<\omega_1} : \omega_1^{L[\sigma]} = \sup(\sigma)\}$. In $\text{ZF} + \text{AD}_{\mathbb{R}} + \text{DC}$, $|S_1| \leq |[\omega_1]^{<\omega_1}|$ however S_1 does not inject into $[\omega_1]^\omega$. Since $S_1 \subseteq [\omega_1]^{<\omega_1}$, a negative answer to Question 1.1 follows from AD^+ by the following result.

Fact 1.3. ([3]) *Assume $\text{ZF} + \text{AD} + \text{DC}_{\mathbb{R}}$ and all sets of reals have an ∞ -Borel code. Then there is no injection of S_1 into ${}^\omega \text{ON}$. As a consequence, there is no injection of $[\omega_1]^{<\omega_1}$ into ${}^\omega \text{ON}$.*

This result uses ∞ -Borel codes to absorb fragments of functions into suitable ZFC models. To the authors' knowledge, the most interesting properties about S_1 (and even to distinguish $|S_1|$ from $|\mathbb{R}|$) require arguments using ∞ -Borel codes. Unlike S_1 , $[\omega_1]^{<\omega_1}$ is a more combinatorial object and [5] distinguished $[\omega_1]^\omega$ and $[\omega_1]^{<\omega_1}$ in AD alone using the almost everywhere continuity property (on sequences of a fixed countable length).

First, this paper will show that under just $\text{ZF} + \text{AD}$, one can prove the following.

Theorem 2.9. Assuming $\text{ZF} + \text{AD}$, $\neg(|[\omega_1]^{<\omega_1}| \leq |{}^\omega(\omega_\omega)|)$.

Then one will obtain the conclusion of Fact 1.3 under just $\text{ZF} + \text{AD} + \text{DC}_{\mathbb{R}}$.

Theorem 4.4. Assume $\text{ZF} + \text{AD} + \text{DC}_{\mathbb{R}}$. There is no injection of $[\omega_1]^{<\omega_1} \rightarrow {}^\omega \text{ON}$.

These two theorems are then used to prove the following theorem.

Theorem 4.7. Assume $\text{ZF} + \text{AD} + \text{DC}_{\mathbb{R}}$. There is no injection of $[\omega_1]^{\omega_1}$ into ${}^{<\omega_1} \text{ON}$.

Assuming just $\text{ZF} + \text{AD}$, $\neg(|[\omega_1]^{\omega_1}| \leq |{}^{<\omega_1}(\omega_\omega)|)$.

Fact 1.3 is proved using techniques that clearly have an AD^+ flavor. In contrast, the results of this paper are proved using an eclectic combination of classical determinacy arguments and more recent results of classical flavor. Almost everywhere continuity results for functions $\Phi : [\omega_1]^\epsilon \rightarrow \omega_1$ from [5] use the Kunen tree which is an important tool for analyzing the ultrapower of ω_1 by the partition measures. Various almost everywhere club uniformization results will be employed. One consequence of these club uniformization results is the almost everywhere continuity property for functions of the form $\Phi : [\omega_1]^{\omega_1} \rightarrow \omega_1$ from [4]. Generic coding arguments, category notions, and the Banach-Mazur games will be used to make uniform selection of cofinal sets and uniform selection of Σ_2^1 bounding prewellorderings. Martin's good coding system and the Martin style games which are used to prove the partition relations will indirectly appear in the almost everywhere good code uniformization.

Ideas involving S_1 require forcing techniques over ZFC-models that do not seem to generalize to cardinals higher than ω_1 . [5] used classical determinacy arguments to prove $\neg(|[\omega_1]^{<\omega_1}| \leq |[\omega_1]^\omega|)$ under just AD, but the techniques used could be generalized to prove $\neg(|[\omega_2]^{<\omega_2}| \leq |[\omega_2]^{\omega_1}|)$ under AD which has no known AD^+ style proof to the authors' knowledge. The methods used here seem to be more suitable to generalizations of the main questions to higher strong partition cardinals such as δ_3^1 .

2. BELOW ω_ω

Let $\epsilon \in \text{ON}$ and $f : \epsilon \rightarrow \text{ON}$. For $\alpha \leq \epsilon$, let $\text{bound}(f, \alpha) = \sup\{f(\beta) : \beta < \alpha\}$. The function f is discontinuous everywhere if and only if for all $\alpha < \epsilon$, $\text{bound}(f, \alpha) < f(\alpha)$. The function f has uniform cofinality ω if and only if there is a function $F : \epsilon \times \omega \rightarrow \text{ON}$ so that for all $\alpha < \epsilon$ and $n \in \omega$, $F(\alpha, n) < F(\alpha, n+1)$ and $F(\alpha) = \sup\{F(\alpha, k) : k \in \omega\}$. The function f has the correct type if and only if it is discontinuous everywhere and has uniform cofinality ω . If X is a set or class of ordinals, then $[X]_*^\epsilon$ and $[X]_*^{<\epsilon}$ are the subsets of $[X]^\epsilon$ and $[X]^{<\epsilon}$ (respectively) consisting of those functions of the correct type. One can show that if $\epsilon \leq \kappa$ and κ is a cardinal, then $||[\kappa]^\epsilon|| = ||[\kappa]_*^\epsilon||$. The notion of having correct type is used to formulate a correct type partition property which provides club homogeneous sets.

Definition 2.1. Let $\epsilon \leq \kappa$ be ordinals. Let $\kappa \rightarrow_* (\kappa)_2^\epsilon$ indicate that for all $P : [\kappa]_*^\epsilon \rightarrow 2$, there is a club $C \subseteq \kappa$ and an $i \in 2$ so that for all $f \in [C]_*^\epsilon$, $P(f) = i$. If $\kappa \rightarrow_* (\kappa)_2^\epsilon$ for all $\epsilon < \kappa$, then κ is said to be a weak partition cardinal. If $\kappa \rightarrow_* (\kappa)_2^\kappa$, then κ is said to be a strong partition cardinal.

Fact 2.2. (Martin; [7] Theorem 12.2, [2] Fact 4.9, [2] Corollary 4.27) Assume ZF + AD. For all $\epsilon \leq \omega_1$, $\omega_1 \rightarrow_* (\omega_1)_2^\epsilon$. (Martin and Paris; [2] Theorem 5.19) For all $\epsilon < \omega_2$, $\omega_2 \rightarrow_* (\omega_2)_2^\epsilon$.

For $1 \leq \epsilon \leq \omega_1$, define the filter W_1^ϵ on $[\omega_1]_*^\epsilon$ by $A \in W_1^\epsilon$ if and only if there is a club $C \subseteq \omega_1$ so that $[C]_*^\epsilon \subseteq A$. The partition relations imply that W_1^ϵ is a countably complete measure. In particular, ω_1 is a measurable cardinal. Using the Kunen tree analysis, it can be shown in ZF + AD (without $\text{DC}_\mathbb{R}$) for each $1 \leq n < \omega_1$, $\prod_{[\omega_1]^n} \omega_1/W_1^n$ is a wellordering under the usual ultrapower ordering and in fact $\omega_{n+1} = \prod_{[\omega_1]^n} \omega_1/W_1^n$. The Martin and Paris weak partition property for ω_2 from Fact 2.2 follows from the ultrapower representation of ω_2 . These partition properties on ω_2 imply that ω_2 is measurable (for instance using the ω -club filter or the ω_1 -club filter on ω_2) and hence regular. The ultrapower representation also shows that for all $n \geq 3$, $\text{cof}(\omega_n) = \omega_2$.

One can explicitly give an ω_2 cofinal sequence through ω_n when $n \geq 2$. Let \mathfrak{V} be the subset of the ultrapower $\prod_{\omega_1} \omega_1/W_1^1$ which has a representative $f : \omega_1 \rightarrow \omega_1$ which is an increasing function of the correct type. Since $|\prod_{\omega_1} \omega_1/W_1^1| = \omega_2$, one also has that $|\mathfrak{V}| = \omega_2$. For $n \geq 2$ and $f : \omega_1 \rightarrow \omega_1$, define $K_n(f) : [\omega_1]^{n-1} \rightarrow \omega_1$ by $K_n(f)(\alpha_1, \dots, \alpha_{n-1}) = f(\alpha_{n-1})$. Define $\rho_n : \mathfrak{V} \rightarrow \prod_{[\omega_1]^{n-1}} \omega_1/W_1^{n-1} = \omega_n$ by $\rho_n([f]_{W_1^1}) = [K_n(f)]_{W_1^{n-1}}$. (Observe that ρ_n is well defined and independent of the choice of representative f .) One can check that $\rho_n : \mathfrak{V} \rightarrow \omega_n$ is cofinal.

Fact 2.3. (Kunen; [2] Theorem 5.10; [6] Lemma 4.1) Assume ZF + AD. Let $f : \omega_1 \rightarrow \omega_1$ be a function. There is a function $\mathcal{K} : \omega_1 \times \omega_1 \rightarrow \omega_1$ so that for each $\omega < \alpha < \omega_1$, $f(\alpha) < \sup\{\mathcal{K}(\alpha, \beta) : \beta < \alpha\} = \{\mathcal{K}(\alpha, \beta) : \beta < \alpha\}$.

Fact 2.4. Assume ZF + AD. Let $\delta < \epsilon \leq \omega_1$ and $\Phi : [\omega_1]_*^\epsilon \rightarrow \omega_1$ have the property that there is a club $C \subseteq \omega_1$ so that for all $f \in [C]_*^\epsilon$, $\Phi(f) < f(\delta)$. Then there is a club $F \subseteq \omega_1$ so that for all $f, g \in [F]_*^\epsilon$, if $f \upharpoonright \delta = g \upharpoonright \delta$, then $\Phi(f) = \Phi(g)$.

Proof. Let δ, ϵ, Φ , and C be as in the statement. Let $\epsilon' = \delta + 1 + (\epsilon - \delta)$. Let $h : \epsilon' \rightarrow \omega_1$. Define $\text{main}(h) : \epsilon \rightarrow \omega_1$ by

$$\text{main}(h)(\alpha) = \begin{cases} h(\alpha) & \alpha < \delta \\ h(\delta + 1 + (\alpha - \delta)) & \alpha \geq \delta \end{cases}$$

Let $\text{extra}(h) \in \omega_1$ by $\text{extra}(h) = h(\delta)$. Define a partition $\Phi : [C]_*^{\epsilon'} \rightarrow 2$ by $P(h) = 0$ if and only if $\Phi(\text{main}(h)) < \text{extra}(h)$. By $\omega_1 \rightarrow_* (\omega_1)_2^{\epsilon'}$, let $D_0 \subseteq C$ be a club homogeneous for P . Let $D_1 \subseteq D_0$ be the set of limit points of D_0 . Pick $f \in [D_1]_*^{\epsilon'}$. Since $f \in [C]_*^{\epsilon'}$, $\Phi(f) < f(\delta) \in D_1$. Let $\gamma \in D_0$ be so that $\text{bound}(f, \delta) < \gamma < f(\delta)$. Let $h : \epsilon' \rightarrow D_0$ be such that $\text{main}(h) = f$ and $\text{extra}(h) = \gamma$. Since $P(h) = 0$, D_0 is homogeneous for P taking value 0.

Let $\text{next}_{D_0} : \omega_1 \rightarrow D_0$ be defined by $\text{next}_{D_0}(\alpha)$ is the least element of D_0 larger than α . For any $f \in [D_1]_*^{\epsilon'}$, $\Phi(f) < \text{next}_{D_0}(\text{bound}(f, \delta))$. This follows from $P(h) = 0$ where $h : \epsilon' \rightarrow D_0$ is defined so that $\text{main}(h) = f$ and $\text{extra}(h) = \text{next}_{D_0}(\text{bound}(f, \delta))$. For each $\sigma \in [D_1]^\delta$, let $V_\sigma : [D_1]^{\epsilon-\delta} \rightarrow \text{next}_{D_0}(\text{sup}(\sigma))$ by $V_\sigma(k) = \Phi(\sigma \hat{\ } k)$. By the countable completeness of $W_1^{\epsilon-\delta}$, there is a club E and a $\gamma_\sigma < \text{next}_{D_0}(\text{sup}(\sigma))$ so that for all $k \in [E]_*^{\epsilon-\delta}$, $V_\sigma(k) = \gamma_\sigma$.

Define $Q : [D_1]_*^\epsilon \rightarrow 2$ by $Q(f) = 0$ if and only if $\Phi(f) = \gamma_{f \upharpoonright \delta}$. By $\omega_1 \rightarrow_* (\omega_1)_2^\epsilon$, there is a club $F \subseteq D_1$ which is homogeneous for Q . Pick any $\sigma \in [F]_*^\delta$. There is a club $E \subseteq F$ so that for all $k \in [E]_*^{\epsilon-\delta}$, $V_\sigma(k) = \gamma_\sigma$. Pick a $k \in [E]_*^{\epsilon-\delta}$ and let $f = \sigma \hat{\ } k$. Then $\Phi(f) = \Phi(\sigma \hat{\ } k) = V_\sigma(k) = \gamma_\sigma = \gamma_{f \upharpoonright \delta}$. So $Q(f) = 0$ and hence F must be homogeneous for Q taking value 0. Thus F is the desired club. \square

The following is the almost everywhere continuity property for functions $\Phi : [\omega_1]^\epsilon \rightarrow \omega_1$ when $\epsilon < \omega_1$.

Fact 2.5. ([5] *Theorem 2.15*) *Assume ZF + AD. Let $\epsilon < \omega_1$ and $\Phi : [\omega_1]_*^\epsilon \rightarrow \omega_1$. There is an $\delta < \epsilon$ and a club $C \subseteq \omega_1$ so that for all $f, g \in [C]_*^\epsilon$, if $\sup(f) = \sup(g)$ and $f \upharpoonright \delta = g \upharpoonright \delta$, then $\Phi(f) = \Phi(g)$.*

Proof. In [5], this fact is derived from a finer and complete analysis of continuity given by [5] Theorem 2.14. The following is a simpler proof of just the coarser continuity stated above.

Define a partition $P : [\omega_1]_*^{\epsilon+1} \rightarrow 2$ by $P(h)$ if and only if $\Phi(h \upharpoonright \epsilon) < h(\epsilon)$. By $\omega_1 \rightarrow_* (\omega_1)_2^{\epsilon+1}$, there is a club $C_0 \subseteq \omega_1$ which is homogeneous for P . Pick an $f \in [C_0]_*^\epsilon$ and let $\gamma \in C_0$ be such that $\Phi(f) < \gamma$. Define $h = f \hat{\ } \gamma$. Since $h \in [C_0]_*^\epsilon$ and $P(h) = 0$, C_0 is homogeneous for P taking value 0. For all $f \in [C_0]_*^\epsilon$, $\Phi(f) < \text{next}_{C_0}(\sup(f))$ by using the fact that $P(h) = 0$ where $h \in [C_0]_*^{\epsilon+1}$ is defined so that $h \upharpoonright \epsilon = f$ and $h(\epsilon) = \text{next}_{C_0}(\sup(f))$. By Fact 2.3, there is a function $\mathcal{K} : \omega_1 \times \omega_1 \rightarrow \omega_1$ with the property that for all $\omega \leq \alpha < \omega_1$, $\text{next}_{C_0}(\alpha) < \sup\{\mathcal{K}(\alpha, \beta) : \beta < \alpha\} = \{\mathcal{K}(\alpha, \beta) : \beta < \alpha\}$. Since for all $f \in [C_0]_*^\epsilon$, $\Phi(f) < \text{next}_{C_0}(\sup(f))$, define $\Psi : [C_0]_*^\epsilon \rightarrow \omega_1$ by $\Psi(f)$ is the least $\beta < \sup(f)$ so that $\Phi(f) = \mathcal{K}(\sup(f), \beta)$. For each $f \in [C_0]_*^\epsilon$, let δ_f be the least δ so that $\Psi(f) < f(\delta)$. Since W_1^ϵ is countably additive, there is a $\delta < \epsilon$ and a club $C_1 \subseteq C_0$ so that for all $f \in [C_1]_*^\epsilon$, $\delta_f = \delta$. Fact 2.4 implies there is a club $C \subseteq C_1$ so that for all $f, g \in [C]_*^\epsilon$, if $f \upharpoonright \delta = g \upharpoonright \delta$, then $\Psi(f) = \Psi(g)$. For any $f, g \in [C]_*^\epsilon$, if $f \upharpoonright \delta = g \upharpoonright \delta$ and $\sup(f) = \sup(g)$, then $\Phi(f) = \mathcal{K}(\sup(f), \Psi(f)) = \mathcal{K}(\sup(g), \Psi(g)) = \Phi(g)$. C is the desired club. \square

[5] uses Fact 2.5 to give an argument in ZF + AD that $[\omega_1]^{<\omega_1}$ does not inject into ${}^\omega\omega_1$. Since this result is the backbone of all other results in the paper, the proof will be given for completeness.

Fact 2.6. ([5] *Theorem 2.16*) *Assuming ZF + AD, $\neg(|[\omega_1]^{<\omega_1}| \leq |{}^\omega\omega_1|)$. In particular, $|[\omega_1]^\omega| < |[\omega_1]^{<\omega_1}|$.*

Proof. Suppose there is an injection $\Phi : [\omega_1]^{<\omega_1} \rightarrow {}^\omega\omega_1$. For each $\epsilon < \omega_1$ and $n \in \omega$, define $\Phi_n^\epsilon : [\omega_1]^\epsilon \rightarrow \omega_1$ by $\Phi_n^\epsilon(f) = \Phi(f)(n)$. By Fact 2.5, there is a $\delta < \epsilon$ and a club $C \subseteq \omega_1$ so that for all $f, g \in [C]_*^\epsilon$, if $f \upharpoonright \delta = g \upharpoonright \delta$ and $\sup(f) = \sup(g)$, then $\Phi_n^\epsilon(f) = \Phi_n^\epsilon(g)$. Let δ_n^ϵ be the least such δ . For each $n \in \omega$, let $\Lambda_n : \omega_1 \rightarrow \omega_1$ be defined by $\Lambda_n(\epsilon) = \delta_n^\epsilon$. Since Λ_n is regressive and the club measure W_1^1 is normal, there is a $\delta_n < \omega_1$ so that there exists a club $C \subseteq \omega_1$ with the property that $\Lambda_n(\epsilon) = \delta_n$ for all $\epsilon \in C$. By the Moschovakis coding lemma, there is a surjection $\pi : \mathbb{R} \rightarrow \mathcal{P}(\omega_1)$. Define $R \subseteq \omega \times \mathbb{R}$ by $R(n, x)$ if and only if $\pi(x)$ is a club with the property that for all $\epsilon \in \pi(x)$, $\Lambda_n(\epsilon) = \delta_n$. By $\text{AC}_\omega^\mathbb{R}$, there is a sequence $\langle x_n : n \in \omega \rangle$ so that for all $n \in \omega$, $R(n, x_n)$. Let $C_n = \pi(x_n)$ and $C = \bigcap_{n \in \omega} C_n$. Let $\delta = \sup\{\delta_n : n \in \omega\}$ and note that $\delta < \omega_1$ since ω_1 is regular. Fix an ordinal $\epsilon > \delta$ with $\epsilon \in C$. Using the Moschovakis coding lemma and $\text{AC}_\omega^\mathbb{R}$ again, there is a sequence $\langle D_n : n \in \omega \rangle$ of club subsets of ω_1 with the property that for all $n \in \omega$, for all $f, g \in [D_n]_*^\epsilon$, if $f \upharpoonright \delta_n^\epsilon = g \upharpoonright \delta_n^\epsilon$ and $\sup(f) = \sup(g)$, then $\Phi_n^\epsilon(f) = \Phi_n^\epsilon(g)$. Let $D = \bigcap_{n \in \omega} D_n$. Pick $f, g \in [D]_*^\epsilon$ so that $f \upharpoonright \delta = g \upharpoonright \delta$, $\sup(f) = \sup(g)$, and $f \neq g$. Since $\delta \geq \delta_n = \delta_n^\epsilon$ (since $\epsilon \in C$) for all $n \in \omega$, one has that $\Phi(f) = \Phi(g)$. This contradicts Φ being an injection. \square

Let club_{ω_1} denote the collection of club subsets of ω_1 . The following is the everywhere ω_1 club uniformization.

Fact 2.7. ([2] *Fact 4.8*) *Assume ZF + AD. Suppose $R \subseteq \omega_1 \times \text{club}_{\omega_1}$ is a relation which is \subseteq -downward closed in the club coordinate. (This means for all $\alpha \in \omega_1$, if $C \subseteq D$ are club subsets of ω_1 and $R(\alpha, D)$, then $R(\alpha, C)$.) Then there is a function $\Phi : \text{dom}(R) \rightarrow \text{club}_{\omega_1}$ so that for all $\alpha \in \text{dom}(R)$, $R(\alpha, \Phi(\alpha))$.*

The following is useful notation. If $f : \omega_1 \rightarrow \omega_1$ and $\epsilon < \omega_1$, then let $\text{drop}(f, \epsilon) : \omega_1 \rightarrow \omega_1$ be defined by $\text{drop}(f, \epsilon)(\alpha) = f(\epsilon + \alpha)$. The following argument appears in [5] for just ω_2 . The following adapts the arguments for all ω_n such that $2 \leq n < \omega$.

Theorem 2.8. *Assume ZF + AD, $\neg(|[\omega_1]^{<\omega_1}| \leq {}^\omega\omega_n)$ for all $n \in \omega$.*

Proof. Suppose $\Phi : [\omega_1]^{<\omega_1} \rightarrow {}^\omega\omega_n$ is an injection. Fact 2.6 implies this is impossible if $n \leq 1$. So suppose $n \geq 2$ and inductively this result has been shown for all $k < n$.

For each $\epsilon < \omega_1$, define a partition $P_\epsilon : [\omega_1]_*^{\omega_1} \rightarrow 2$ by $P_\epsilon(f) = 0$ if and only if $\sup(\Phi(f \upharpoonright \epsilon)) < \rho_n([\mathbf{drop}(f, \epsilon)]_{W_1^1})$, where $\rho_n : \mathfrak{V} \rightarrow \omega_n$ is cofinal. (Note that $\sup(\Phi(f \upharpoonright \epsilon)) < \omega_n$ since $\Phi(f \upharpoonright \epsilon) \in {}^\omega \omega_n$ and $\text{cof}(\omega_n) = \omega_2$.) By $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$, there is a club $C \subseteq \omega_1$ which is homogeneous for P_ϵ . Pick any $\sigma \in [C]_*^\epsilon$. Since ρ_n is cofinal, there is an $h \in [C]_*^{\omega_1}$ with $\sup(\sigma) < h(0)$ and $\sup(\Phi(\sigma)) < \rho_n([h]_{W_1^1})$. Let $f = \sigma \hat{\ } h$. Note that $f \in [C]_*^{\omega_1}$ and $P_\epsilon(f) = 0$ since $\sup(\Phi(f \upharpoonright \epsilon)) = \sup(\Phi(\sigma)) < \rho_n([h]_{W_1^1}) = \rho_n([\mathbf{drop}(f, \epsilon)]_{W_1^1})$. Thus C is homogeneous for P_ϵ taking value 0. Fix a $g \in [C]_*^{\omega_1}$. Let $\beta = \rho_n([g]_{W_1^1})$. For any $\sigma \in [C]_*^\epsilon$, let γ_σ be the least $\gamma < \omega_1$ so that $\sup(\sigma) < g(\gamma)$. Let $f_\sigma = \sigma \hat{\ } \mathbf{drop}(g, \gamma_\sigma)$ and note that $f_\sigma \in [C]_*^{\omega_1}$. $P(f_\sigma) = 0$ implies that $\sup(\Phi(\sigma)) < \rho_n([\mathbf{drop}(f_\sigma, \epsilon)]_{W_1^1}) = \rho_n([g]_{W_1^1}) = \beta$ since $[\mathbf{drop}(f_\sigma, \epsilon)]_{W_1^1} = [g]_{W_1^1}$. Let β_ϵ be the least ordinal β for which there exists a club $C \subseteq \omega_1$ so that for all $\sigma \in [C]_*^\epsilon$, $\sup(\Phi(\sigma)) < \beta$.

Define a relation $R \subseteq \omega_1 \times \text{club}_{\omega_1}$ by $R(\epsilon, C)$ if and only if for all $\sigma \in [C]_*^\epsilon$, $\sup(\Phi(\sigma)) < \beta_\epsilon$. Note that R is \subseteq -downward in the club_{ω_1} coordinate and $\text{dom}(R) = \omega_1$. By Fact 2.7, there is a sequence $\langle C_\epsilon : \epsilon < \omega_1 \rangle$ so that for all $\epsilon < \omega_1$, $R(\epsilon, C_\epsilon)$. Let $\delta = \sup\{\beta_\epsilon : \epsilon < \omega_1\}$ and note that $\delta < \omega_n$ since $\text{cof}(\omega_n) = \omega_2$. Let $B : \delta \rightarrow \omega_{n-1}$ be an injection. Define an injection $\Psi : \bigcup_{\epsilon < \omega_1} [C_\epsilon]_*^\epsilon \rightarrow {}^\omega \omega_{n-1}$ by $\Psi(\sigma)(n) = B(\Phi(\sigma)(n))$. This is well defined since for all $\sigma \in [C_\epsilon]_*^\epsilon$, $\Phi(\sigma)(n) < \beta_\epsilon < \delta$. Since $|\bigcup_{\epsilon < \omega_1} [C_\epsilon]_*^\epsilon| = |[\omega_1]^{<\omega_1}|$, Ψ induces an injection of $[\omega_1]^{<\omega_1}$ into ${}^\omega \omega_{n-1}$, which is impossible by the induction hypothesis. \square

Note that the next result implies Theorem 2.8; however, the proof is more tedious.

Theorem 2.9. *Assume ZF + AD, $\neg(|[\omega_1]^{<\omega_1}| \leq |{}^\omega(\omega_\omega)|)$.*

Proof. Suppose there is an injection $\Phi : [\omega_1]^{<\omega_1} \rightarrow {}^\omega \omega_\omega$. By the countable additivity of W_1^1 , for each $n \in \omega$ and $\epsilon < \omega_1$, there is a club C and an integer b so that $\Phi(\sigma)(n) < \omega_b$ for all $\sigma \in [C]_*^\epsilon$. Let b_n^ϵ be the least such integer. Again by the countable additivity of W_1^1 , for each $n \in \omega$, there is a club D and an integer b_n so that for all $\epsilon \in D$, $b_n^\epsilon = b_n$. By the Moschovakis coding lemma and $\text{AC}_\omega^\mathbb{R}$, there is a sequence $\langle D_n : n \in \omega \rangle$ of club subsets of ω_1 so that for all $\epsilon \in D_n$, $b_n^\epsilon = b_n$. Let $D^* = \bigcap_{n \in \omega} D_n$.

Claim 1: For each $n \in \omega$, there is a sequence $\langle E_\epsilon : \epsilon \in D^* \rangle$ of club subsets of ω_1 and an injection $I : A \rightarrow \omega_1$, where $A = \{\Phi(\sigma)(n) : \sigma \in \bigcup_{\epsilon \in D^*} [E_\epsilon]_*^\epsilon\}$.

To see Claim 1: Fix $n \in \omega$. Recall for each $i \geq 2$, there are cofinal maps $\rho_i : \mathfrak{V} \rightarrow \omega_i$ of \mathfrak{V} (which has cardinality ω_2) into ω_i . Define $R \subseteq D^* \times \text{club}_{\omega_1}$ by $R(\epsilon, C)$ if and only if for all $\sigma \in [C]_*^\epsilon$, $\Phi(\sigma)(n) < \omega_{b_n}$. Since R is \subseteq -downward closed in the club_{ω_1} -coordinate and $\text{dom}(R) = D^*$, Fact 2.7 implies there is a sequence $\langle C'_\epsilon : \epsilon \in D^* \rangle$ so that for all $\epsilon \in D^*$, $R(\epsilon, C'_\epsilon)$. Let $C_\epsilon^{b_n} = C'_\epsilon$, $a_{b_n} = \omega_{b_n}$, and $\Psi_{b_n} : \bigcup_{\epsilon \in D^*} [C_\epsilon^{b_n}]_*^\epsilon \rightarrow \omega_{a_{b_n}}$ be defined by $\Psi_{b_n}(\sigma) = \Phi(\sigma)(n)$.

Suppose for $0 < k \leq b_n$, the following objects have been defined.

- For all $k \leq j \leq b_n$, $a_j \leq b_n$ and for all $k < j \leq b_n$, if $a_j > 1$, then $a_{j-1} < a_j$.
- For all $k \leq j \leq b_n$, $\langle C_\epsilon^j : \epsilon \in D^* \rangle$ is a sequence of clubs and for all $k \leq j_0 < j_1 \leq b_n$, $C_\epsilon^{j_0} \subseteq C_\epsilon^{j_1}$.
- For all $k \leq j \leq b_n$, $\Psi_j : \bigcup_{\epsilon \in D^*} [C_\epsilon^j]_*^\epsilon \rightarrow \omega_{a_j}$.
- For $k < j \leq b_n$, $I_j : T_j \rightarrow \omega_{a_{j-1}}$ is an injection where $T_j = \{\Psi_j(\sigma) : \sigma \in \bigcup_{\epsilon \in D^*} [C_\epsilon^j]_*^\epsilon\}$.

(Case I: $a_k > 1$.) Fix $\epsilon \in D^*$. Define $P_\epsilon^k : [C_\epsilon^k]_*^{\omega_1} \rightarrow 2$ by $P_\epsilon^k(f) = 0$ if and only if $\Psi_k(f \upharpoonright \epsilon) < \rho_{a_k}([\mathbf{drop}(f, \epsilon)]_{W_1^1})$. By $\omega_1 \rightarrow_* (\omega_1)_2^{\omega_1}$, there is a club $K \subseteq C_\epsilon^k$ which is homogeneous for P_ϵ^k . Fix a $\sigma \in [K]_*^\epsilon$. Pick any $\ell \in [K]_*^{\omega_1}$ so that $\ell(0) > \sup(\sigma)$ and $\Psi_k(\sigma) < \rho_{a_k}([\ell]_{W_1^1})$ which is possible since $\rho_{a_k} : \mathfrak{V} \rightarrow \omega_{a_k}$ is cofinal. Let $f = \sigma \hat{\ } \ell$. Note that $P_\epsilon^k(f) = 0$ since $\Psi_k(f \upharpoonright \epsilon) = \Psi_k(\sigma) < \rho_{a_k}([\ell]_{W_1^1}) = \rho_{a_k}([\mathbf{drop}(f, \epsilon)]_{W_1^1})$. Since $f \in [K]_*^{\omega_1}$, K is homogeneous for P_ϵ^k taking value 0. Now fix an $\ell \in [K]_*^{\omega_1}$ and let $\beta = \rho_{a_k}([\ell]_{W_1^1})$. For any $\sigma \in [K]_*^\epsilon$, let γ_σ be the least γ so that $\sup(\sigma) < \ell(\gamma)$. Let $f_\sigma = \sigma \hat{\ } \mathbf{drop}(\ell, \gamma_\sigma)$. $P_\epsilon^k(f_\sigma) = 0$ implies that $\Psi_k(\sigma) = \Psi_k(f_\sigma \upharpoonright \epsilon) < \rho_{a_k}([\mathbf{drop}(f_\sigma, \epsilon)]_{W_1^1}) = \rho_{a_k}([\ell]_{W_1^1}) = \beta$ since $[\mathbf{drop}(f_\sigma, \epsilon)]_{W_1^1} = [\ell]_{W_1^1}$. It has been shown that there is a β so that there exists a club $K \subseteq C_\epsilon^k$ with the property that for all $\sigma \in [K]_*^\epsilon$, $\Psi_k(\sigma) < \beta$. Let β_ϵ be the least such $\beta \geq \omega$ with this property. Let $\delta = \sup\{\beta_\epsilon : \epsilon \in D^*\}$. Since $a_k > 1$, $\text{cof}(\omega_{a_k}) = \omega_2$ so $\delta < \omega_{a_k}$.

Now define $S \subseteq D^* \times \text{club}_{\omega_1}$ by $S(\epsilon, K)$ if and only if $K \subseteq C_\epsilon^k$ and for all $\sigma \in [K]_*^\epsilon$, $\Psi_k(\sigma) < \delta$. Note that $\text{dom}(S) = D^*$ by the previous discussion and S is \subseteq -downward closed in the club_{ω_1} -coordinate. By Fact 2.7, there is a sequence of clubs $\langle C_\epsilon^{k-1} : \epsilon \in D^* \rangle$ with the property that for all $\epsilon \in D^*$, $S(\epsilon, C_\epsilon^{k-1})$. For all $\sigma \in \bigcup_{\epsilon \in D^*} [C_\epsilon^{k-1}]_*^\epsilon$, $\Psi_k(\sigma) < \beta_{|\sigma|} < \delta < \omega_{a_k}$. Since $\omega \leq \delta < \omega_{a_k}$, there is an $a_{k-1} < a_k$ so that $\omega_{a_{k-1}} \leq \delta < \omega_{a_{k-1}+1}$. Let $I_k : \delta \rightarrow \omega_{a_{k-1}}$ be an injection. Let $T_k = \{\Psi_k(\sigma) : \sigma \in \bigcup_{\epsilon \in D^*} [C_\epsilon^{k-1}]_*^\epsilon\}$ and note that the restriction $I_k : T_k \rightarrow \omega_{a_{k-1}}$ is an injection. Let $\Psi_{k-1} : \bigcup_{\epsilon \in D^*} [C_\epsilon^{k-1}]_*^\epsilon \rightarrow \omega_{a_{k-1}}$ be defined by $I_k \circ \Psi_k$.

(Case II: $a_k \leq 1$) Let $a_{k-1} = a_k$, $C_\epsilon^{k-1} = C_\epsilon^k$ for each $\epsilon \in D^*$, $T_k = \{\Psi_k(\sigma) : \sigma \in \bigcup_{\epsilon \in D^*} [C_\epsilon^{k-1}]_\epsilon^*\}$, $I_k : T_k \rightarrow \omega_{a_{k-1}}$ be the inclusion map, and $\Psi_{k-1} = \Psi_k$.

By recursion, one has constructed $\langle a_k : 0 \leq k \leq b_n \rangle$, $\langle \langle C_\epsilon^k : \epsilon \in D^* \rangle : 0 \leq k \leq b_n \rangle$, $\langle T_k : 0 < k \leq b_n \rangle$, and $\langle I_k : 0 < k \leq b_n \rangle$. Since $a_{k-1} < a_k$ for all $0 < k \leq b_n$ such that $a_k > 1$ and $a_{b_n} = b_n$, one must have that $a_0 \leq 1$. For each $\epsilon \in D^*$, let $E_\epsilon = C_\epsilon^0$ and $I : A \rightarrow \omega_1$ be defined by $I = I_1 \circ \dots \circ I_{b_n}$, where recall that $A = \{\Phi(\sigma)(n) : \sigma \in \bigcup_{\epsilon \in D^*} [E_\epsilon]_\epsilon^*\}$. This completes the proof of Claim 1.

Using Claim 1, the Moschovakis coding lemma, and $\text{AC}_\omega^\mathbb{R}$, there exist sequences $\langle \langle E_\epsilon^n : \epsilon \in D^* \rangle : n \in \omega \rangle$ and $\langle I_n : n \in \omega \rangle$ so that for all $n \in \omega$, $I_n : A_n \rightarrow \omega_1$ is an injection where $A_n = \{\Phi(\sigma)(n) : \sigma \in \bigcup_{\epsilon \in D^*} [E_\epsilon^n]_\epsilon^*\}$. For each $\epsilon \in D^*$, let $E_\epsilon = \bigcap_{n \in \omega} E_\epsilon^n$. Let $\Sigma : \bigcup_{\epsilon \in D^*} [E_\epsilon]_\epsilon^* \rightarrow {}^\omega \omega_1$ be defined by $\Sigma(\sigma)(n) = I_n(\Phi(\sigma)(n))$. Now suppose $\sigma_0, \sigma_1 \in \bigcup_{\epsilon \in D^*} [E_\epsilon]_\epsilon^*$ and $\sigma_0 \neq \sigma_1$. Since Φ is an injection, $\Phi(\sigma_0) \neq \Phi(\sigma_1)$. Thus there is some $n \in \omega$ so that $\Phi(\sigma_0)(n) \neq \Phi(\sigma_1)(n)$. Since $\Phi(\sigma_0)(n), \Phi(\sigma_1)(n) \in A_n$ and $I_n : A_n \rightarrow \omega_1$ is an injection, one has that $\Sigma(\sigma_0) \neq \Sigma(\sigma_1)$ because $\Sigma(\sigma_0)(n) = I_n(\Phi(\sigma_0)(n)) \neq I_n(\Phi(\sigma_1)(n)) = \Sigma(\sigma_1)(n)$. It has been shown that Σ is an injection. Since $|\bigcup_{\epsilon \in D^*} [E_\epsilon]_\epsilon^*| = |\omega_1|^{<\omega_1}$, Σ induces an injection from $[\omega_1]^{<\omega_1}$ to ${}^\omega \omega_1$. This is impossible by Fact 2.6. \square

3. UNIFORM CHOICE OF UNBOUNDED SUBSETS AND BOUNDING PREWELLORDERINGS

Definition 3.1. Let $2 \leq \alpha < \omega_1$. For $s \in {}^{<\omega} \alpha$, let $N_s^\alpha = \{f \in {}^\omega \alpha : s \subseteq f\}$. Give ${}^\omega \alpha$ the topology generated by $\{N_s^\alpha : s \in {}^{<\omega} \alpha\}$ as a basis. Using this topology, one can define the usual category notions. Note that since α is countable, ${}^\omega \alpha$ is homeomorphic to the usual topology on ${}^\omega \omega$. Let surj_α be the set of functions $f : \omega \rightarrow \alpha$ which are surjections, i.e. $f[\omega] = \alpha$. Observe that surj_α is a comeager subset of ${}^\omega \alpha$.

Under AD, the meager ideal on ${}^\omega \omega$ has full wellordered additivity. That is, if $\lambda \in \text{ON}$ and $\langle A_\alpha : \alpha < \lambda \rangle$ is a sequence of meager subsets of ${}^\omega \omega$, then $\bigcup_{\alpha < \lambda} A_\alpha$ is a meager subset of ${}^\omega \omega$. Since, ${}^\omega \omega$ and ${}^\omega \alpha$ are homeomorphic for each $\alpha < \omega_1$, the meager ideal on ${}^\omega \alpha$ also has full wellordered additivity.

The following is a simple form of the Kechris-Woodin generic coding function [9] for ω_1 .

Fact 3.2. *There is a function $\mathfrak{G} : {}^\omega \omega_1 \rightarrow \text{WO}$ so that for $\alpha < \omega_1$, if $f \in \text{surj}_\alpha$, then $\text{ot}(\mathfrak{G}(f)) = \alpha$.*

Proof. Let $f \in {}^\omega \alpha$. Let $A_f = \{n \in \omega : (\forall m)(m < n \Rightarrow f(m) \neq f(n))\}$. Define $\mathfrak{G}(f)$ to be the element of WO with domain A_f so that for all $m, n \in A_f$, $m <_{\mathfrak{G}(f)} n$ if and only if $f(m) < f(n)$. Note that if $f \in \text{surj}_\alpha$, then $(A_f, <_{\mathfrak{G}(f)})$ is order isomorphic to α . \square

Fact 3.3. *Assume ZF + AD. Let $\langle \nu_\alpha : \alpha < \omega_1 \rangle$ be a sequence of ordinals so that for all $\alpha < \omega_1$, $\text{cof}(\nu_\alpha) \leq \omega_1$ and $\sup\{\nu_\alpha : \alpha < \omega_1\} < \Theta$. Then there is a sequence $\langle K_\alpha : \alpha < \omega_1 \rangle$ so that for all $\alpha < \omega_1$, $K_\alpha \subseteq \nu_\alpha$, $|K_\alpha| \leq \omega_1$, and $\sup K_\alpha = \nu_\alpha$.*

Proof. Let $\delta = \sup\{\nu_\alpha : \alpha < \omega_1\} < \Theta$. By the Moschovakis coding lemma, there is a surjection $\pi : \mathbb{R} \rightarrow \mathcal{P}(\delta)$. Define a relation $S \subseteq \text{WO} \times \mathbb{R}$ by $S(w, r)$ if and only if $\pi(r)$ codes a function $\rho : \omega_1 \rightarrow \nu_{\text{ot}(w)}$ such that $\sup \rho[\omega_1] = \nu_{\text{ot}(w)}$. By the Moschovakis coding lemma, there is relation R with the following properties.

- $R \subseteq S$ and R is Σ_2^1 .
- For all $\alpha < \omega_1$, $R \cap (\text{WO}_\alpha \times \mathbb{R}) \neq \emptyset$.

Let $T \subseteq \text{WO} \times \mathbb{R}$ be defined by

$$T(w, r) \Leftrightarrow (\exists v)(v \in \text{WO} \wedge \text{ot}(v) = \text{ot}(w) \wedge R(v, r)).$$

T is Σ_2^1 , $\text{dom}(T) = \text{WO}$, and for all $w \in \text{WO}$, $T(w, r)$ if and only if $\pi(r)$ codes a function $\rho : \omega_1 \rightarrow \nu_{\text{ot}(w)}$ such that $\sup \rho[\omega_1] = \nu_{\text{ot}(w)}$. Since T is Σ_2^1 , AD implies that T has a uniformization function $\Psi' : \text{WO} \rightarrow \mathbb{R}$, i.e. for all $w \in \text{WO}$, $T(w, \Psi'(w))$. For each $w \in \text{WO}$, let $\Psi(w) = \pi(\Psi'(w))$, i.e. $\Psi(w)$ is the unbounded function from ω_1 into $\nu_{\text{ot}(w)}$ coded by $\Psi'(w)$.

Define a partial function H as follows. For $\alpha < \omega_1$, $\beta < \omega_1$, and $s \in {}^{<\omega} \alpha$, $(\alpha, \beta, s) \in \text{dom}(H)$ if and only if there is an $\eta < \nu_\alpha$ such that $\{f \in N_s^\alpha \cap \text{surj}_\alpha : \Psi(\mathfrak{G}(f))(\beta) = \eta\}$ is comeager in N_s^α . If $(\alpha, \beta, s) \in \text{dom}(H)$, then let $H(\alpha, \beta, s)$ be the unique η with the above property.

Define $K_\alpha = \{H(\alpha, \beta, s) : (\alpha, \beta, s) \in \text{dom}(H)\}$. Note $|K_\alpha| \leq \omega_1$. It remains to show that $\sup K_\alpha = \nu_\alpha$. Fix $\gamma < \nu_\alpha$. For each $f \in \text{surj}_\alpha$, let β_f be the least $\beta < \omega_1$ so that $\Psi(\mathfrak{G}(f))(\beta) > \gamma$. For each $\beta < \omega_1$, let $B_\beta = \{f \in \text{surj}_\alpha : \beta_f = \beta\}$. Since surj_α is comeager, $\text{surj}_\alpha = \bigcup_{\beta < \omega_1} B_\beta$, and wellordered unions of meager subsets of ${}^\omega \alpha$ are meager, there is a $\beta^* < \omega_1$ so that B_{β^*} is nonmeager. For each $\zeta > \gamma$, let

$C_\zeta = \{f \in B_{\beta^*} : \Psi(\mathfrak{G}(f))(\beta^*) = \zeta\}$. Since B_{β^*} is nonmeager, $B_{\beta^*} = \bigcup_{\zeta > \gamma} C_\zeta$, and wellordered unions of meager subsets of ${}^\omega\alpha$ are meager, there is a $\zeta^* > \gamma$ so that C_{ζ^*} is nonmeager. By the Baire property, there is an $s \in {}^\omega\alpha$ so that C_{ζ^*} is comeager in N_s^α . Then for comeagerly many $f \in N_s^\alpha$, $\Psi(\mathfrak{G}(f))(\beta^*) = \zeta^* > \gamma$. Hence $(\alpha, \beta^*, s) \in \text{dom}(H)$ and $H(\alpha, \beta^*, s) = \zeta^* > \gamma$. Thus $\zeta^* \in K_\alpha$. Since $\gamma < \nu_\alpha$ was arbitrary, $\sup K_\alpha = \nu_\alpha$. \square

Let Γ be a (boldface) pointclass, $\check{\Gamma}$ be the dual pointclass of Γ , and $\Delta = \Gamma \cap \check{\Gamma}$. Let $\delta(\Gamma)$ be the supremum of the prewellorderings on \mathbb{R} which belong to Δ . Let $v(\Gamma)$ be the supremum of the $\check{\Gamma}$ wellfounded relations on \mathbb{R} . If $A \in \mathcal{P}(\mathbb{R})$, then let $\text{rk}_W(A)$ denote the Wadge rank of A (which exists assuming $\text{DC}_\mathbb{R}$). If Γ is a pointclass, then let $o(\Gamma) = \sup\{\text{rk}_W(A) : A \in \Gamma\}$.

Fact 3.4. ([6] Lemma 2.13 and 2.16) *Suppose Γ is a nonselfdual pointclass, closed under $\forall^\mathbb{R}$, \vee , \wedge , and has the prewellordering property. Then $\delta(\Gamma) = v(\Gamma)$ and is a regular cardinal.*

Fact 3.5. ([8] Lemma 2.3) *Assume $\text{ZF} + \text{AD} + \text{DC}_\mathbb{R}$. Suppose Δ is a (boldface) pointclass closed under \neg , \wedge , and $\forall^\mathbb{R}$. Then $o(\Delta) = \delta(\Delta)$.*

Fact 3.6. *Assume $\text{ZF} + \text{AD} + \text{DC}_\mathbb{R}$. Suppose Γ is a nonselfdual pointclass closed under $\exists^\mathbb{R}$, $\forall^\mathbb{R}$, \wedge , \vee , and has the prewellordering property. $\delta = \delta(\Gamma) = \delta(\Delta) = o(\Delta)$ is a regular cardinal. Let C be the set of $\eta < \delta$ so that $\Upsilon_\eta = \{A \subseteq \mathbb{R} : \text{rk}_W(A) < \eta\}$ is a pointclass closed under $\exists^\mathbb{R}$. C is a club subset of δ .*

Proof. The first statement follows from Fact 3.4 and Fact 3.5. It remains to show that the set C defined above is a club subset of δ . Let $\gamma < \delta$. Since $\delta = o(\Delta)$, find some $A \in \Delta$ so that $\gamma < \text{rk}_W(A)$. The pointclass $\Sigma_1^1(A)$ is the smallest nonselfdual pointclass containing A and closed under $\exists^\mathbb{R}$, \wedge , and \vee . Let $U \in \Sigma_1^1(A)$ be a universal set. Let $\xi = \text{rk}_W(U) + 1$. Note that $\gamma < \xi$ and since $\Upsilon_\xi = \Sigma_1^1(A)$, one has that $\xi \in C$. This shows that C is unbounded. Suppose ξ is a limit of points in C . Suppose $B \subseteq \mathbb{R} \times \mathbb{R}$ and $\text{rk}_W(B) < \xi$. There is some $\xi' < \xi$ with $\xi' \in C$ so that $\text{rk}_W(B) < \xi' < \xi$. Since $B \in \Upsilon_{\xi'}$ and $\Upsilon_{\xi'}$ is closed under $\exists^\mathbb{R}$, $\exists^\mathbb{R} B \in \Upsilon_{\xi'}$. Hence $\text{rk}_W(\exists^\mathbb{R} B) \leq \xi' < \xi$. Thus Υ_ξ is closed under $\exists^\mathbb{R}$. It has been shown that C is a club. \square

For any $A \subseteq \mathbb{R}$, an example of such a pointclass closed under $\exists^\mathbb{R}$, $\forall^\mathbb{R}$, \wedge , \vee , having the prewellordering property, and containing A is $\Sigma_1^{L(A, \mathbb{R})}$. Let δ_A denote the first Σ_1 -stable ordinal of $L(A, \mathbb{R})$, i.e. the least δ so that $L_\delta(A, \mathbb{R}) \prec_1 L(A, \mathbb{R})$. It can be shown that $\delta_A = o(\Delta_1^{L(A, \mathbb{R})}) = \delta(\Sigma_1^{L(A, \mathbb{R})})$.

Fact 3.7. (Steel; [10] Theorem 2.1) *Assume $\text{ZF} + \text{AD} + \text{DC}_\mathbb{R}$. Let Γ be a nonselfdual pointclass and $\Delta = \Gamma \cap \check{\Gamma}$. Suppose $\exists^\mathbb{R} \Delta \subseteq \Delta$. Suppose $\kappa < \text{cof}(o(\Delta))$. If A is κ -Suslin and $B \in \Gamma$, then $A \cap B \in \Gamma$.*

The next result is a finer version of Steel's result concerning Suslin bounded prewellorderings with particular emphasis on a bound for the Wadge rank of the desired prewellordering.

Fact 3.8. (Steel) *Assume $\text{ZF} + \text{AD} + \text{DC}_\mathbb{R}$. Let δ be such that $\text{cof}(\delta) \geq \omega_2$. Suppose κ is such that $\kappa > \delta$ and there is a pointclass Γ^* which is closed under $\forall^\mathbb{R}$, $\exists^\mathbb{R}$, \wedge , \vee , has the prewellordering property, and $o(\Gamma^*) = o(\Delta^*) = \kappa$, where $\Delta^* = \Gamma^* \cap \check{\Gamma}^*$. Then there is a prewellordering (P, \preceq) with the following properties.*

- *The length of (P, \preceq) is δ . Let $\varphi : P \rightarrow \delta$ be the associated norm of \preceq .*
- *$\varphi : P \rightarrow \delta$ satisfies Σ_2^1 bounding, which means that for all $\Sigma_2^1 S \subseteq P$, there is a $\zeta < \delta$ so that $\varphi[S] \subseteq \zeta$.*
- *$\text{rk}_W(P) < \kappa$ and $\text{rk}_W(\preceq) < \kappa$.*

Proof. This argument follows the template from [6] Theorem 2.28 with additional complexity calculations. Let $\nu = \text{cof}(\delta)$. Let η be the ν^{th} -element of the club $C \subseteq \kappa$ from Fact 3.6. Then $\Upsilon_\eta = \{A \subseteq \mathbb{R} : \text{rk}_W(A) < \eta\}$ is a pointclass closed under $\exists^\mathbb{R}$. Let $B \subseteq \mathbb{R}$ be such that $\text{rk}_W(B) = \eta$. Let $\Gamma = \Upsilon_{\eta+1} = \{A \subseteq \mathbb{R} : A \leq_W B\}$. By a basic property of the Wadge degrees, Γ is nonselfdual because $\text{cof}(\text{rk}_W(B)) > \omega$. Also observe $\Upsilon_\eta = \Gamma \cap \check{\Gamma}$.

Fix some recursive coding of continuous functions $F : \mathbb{R} \rightarrow \mathbb{R}$ by \mathbb{R} . If $x \in \mathbb{R}$, then let $\Sigma_x : \mathbb{R} \rightarrow \mathbb{R}$ denote the continuous function coded by x . Let $\pi : \omega \times \omega \rightarrow \omega$ be a recursive bijection. If $x \in \mathbb{R} = {}^\omega\omega$ and $n \in \omega$, then let $x^{[n]}(k) = x(\pi(n, k))$.

Let $E \subseteq \mathbb{R}$ be defined by $x \in E$ if and only if $\Sigma_{x[0]}^{-1}[B] = \mathbb{R} \setminus \Sigma_{x[1]}^{-1}[B]$. Note that $E \in \Delta^*$ since Δ^* is closed under \vee , \wedge , \neg , $\forall^\mathbb{R}$, and $\exists^\mathbb{R}$. Therefore $o(E) < \kappa$. Note that if $x \in E$, then $\Sigma_{x[0]}^{-1}[B] \in \Upsilon_\eta$ and for every $A \in \Upsilon_\eta$,

there is some $x \in E$ so that $A = \Sigma_{x^{[0]}}^{-1}[B]$. Define a prewellordering $\varphi_0 : E \rightarrow \eta$ by $\varphi_0(x) = \text{rk}_W(\Sigma_{x^{[0]}}^{-1}[B])$. For all $x, y \in E$, define $x \preceq_{\varphi_0} y$ if and only if $\varphi_0(x) \leq \varphi_0(y)$. Note that $x \preceq_{\varphi_0} y$ if and only if

$$(\exists^{\mathbb{R}} z)(\Sigma_{x^{[0]}}^{-1}[B] = \Sigma_z^{-1}[\Sigma_{y^{[0]}}^{-1}[B]]).$$

Thus the prewellordering \preceq_{φ_0} associated to φ_0 belongs to Δ^* and hence has Wadge rank below κ .

Next, one seeks to show that $\varphi_0 : E \rightarrow \eta$ is Σ_2^1 bounded. Let $S \subseteq E$ be a Σ_2^1 set (so it is also an ω_1 -Suslin set). Suppose $\sup \varphi_0[S] = \eta$. Define $F_0 \subseteq \mathbb{R} \times \mathbb{R}$ by

$$F_0(x, y) \Leftrightarrow (x \in S \wedge \Sigma_{x^{[0]}}(y) \in B) \Leftrightarrow (x \in S \wedge \Sigma_{x^{[1]}}(y) \notin B).$$

Since $\Upsilon_\eta = \Gamma \cap \check{\Gamma}$ is closed under $\exists^{\mathbb{R}}$ and $\omega_1 < \text{cof}(\eta) = \text{cof}(o(\Upsilon_\eta))$, one has that F_0 belongs to Υ_η by applying Fact 3.7 to the ω_1 -Suslin set S , the set $B \in \Gamma$, and the set $(\mathbb{R} \setminus B) \in \check{\Gamma}$. Now suppose $A \in \Upsilon_\eta$ and thus $\text{rk}_W(A) < \eta$. Since $\sup \varphi_0[S] = \eta$, there is some $x \in S$ so that $\varphi_0(x) > \text{rk}_W(A)$. Thus $A \leq_W \Sigma_{x^{[0]}}^{-1}[B] = (F_0)_x \leq_W F_0$. This shows that every set in Υ_η is Wadge reducible to F_0 . Thus $\text{rk}_W(F_0) \geq \eta$ however $\text{rk}_W(F_0) < \eta$ since $F_0 \in \Upsilon_\eta$. Thus $\sup \varphi_0[S] < \eta$. It has been shown that there is Σ_2^1 bounded prewellordering $\varphi_0 : E \rightarrow \eta$ where $\eta < \kappa$, $\text{cof}(\eta) = \nu$, and the Wadge rank of the associated prewellordering \preceq_{φ_0} is less than κ .

Since $\text{cof}(\eta) = \nu$, let $\rho_0 : \nu \rightarrow \eta$ be an increasing cofinal map. Define $\varphi_1 : E \rightarrow \nu$ by $\varphi_1(x)$ is the least $\alpha < \nu$ so that $\rho_0(\alpha) \geq \varphi_0(x)$. Suppose $S \subseteq E$ is Σ_2^1 . Since φ_0 is Σ_2^1 bounded, there is some $\zeta < \eta$ so that $\varphi_0[S] \subseteq \zeta$. Since ρ_0 is cofinal through η , there is some $\xi < \nu$ so that $\rho_0(\xi) > \zeta$. Thus $\varphi_1[S] \subseteq \xi$. Hence φ_1 is also Σ_2^1 bounded.

Since $\nu < \kappa = o(\Delta^*) = \delta(\Delta^*)$, there is a norm $\psi_0 : \mathbb{R} \rightarrow \nu$ whose associated prewellordering \preceq_{ψ_0} belongs to Δ^* . Define a relation $S \subseteq \mathbb{R} \times E$ by $S(x, y)$ if and only if $\rho_0(\psi_0(x)) = \varphi_0(y)$. By the Moschovakis coding lemma, there is an $R \subseteq S$ so that for all $\alpha < \nu$, $R \cap (\psi_0^{-1}[\{\alpha\}] \times E) \neq \emptyset$ and $R \in \Delta^*$ (in fact $R \in \Sigma_1^1(\preceq_{\psi_0}) \subseteq \Delta^*$). Note that $x \preceq_{\varphi_1} y$ if and only if

$$(\forall^{\mathbb{R}} a)(\forall^{\mathbb{R}} b)[(R(a, b) \wedge y \preceq_{\varphi_0} b) \Rightarrow x \preceq_{\varphi_0} b].$$

Thus $\preceq_{\varphi_1} \in \Delta^*$ and $\text{rk}_W(\preceq_{\varphi_1}) < \kappa$. It has been shown that there is a prewellordering $\varphi_1 : E \rightarrow \nu$ which is Σ_2^1 bounded and $\text{rk}_W(\preceq_{\varphi_1}) < \kappa$.

Since $\text{cof}(\delta) = \nu$, let $\rho_1 : \nu \rightarrow \delta$ be an increasing cofinal map. Since $o(\Delta^*) = \delta(\Delta^*) = \kappa$ and $\delta < \kappa$, let $\psi_1 : \mathbb{R} \rightarrow \delta$ be a surjective map so that $\text{rk}_W(\preceq_{\psi_1}) < \kappa$. Define $S_1 \subseteq E \times \mathbb{R}$ by $S_1(x, y)$ if and only if $\rho_1(\varphi_1(x)) = \psi_1(y)$. By the Moschovakis coding lemma, there is a relation $R_1 \subseteq S_1$ such that $R_1 \in \Sigma_1^1(\preceq_{\varphi_1}) \subseteq \Delta^*$ and for all $\alpha < \nu$, $R_1 \cap (\varphi_1^{-1}[\{\alpha\}] \times \mathbb{R}) \neq \emptyset$.

For each $\beta < \delta$, define $P_\beta \subseteq \mathbb{R}$ by $x \in P_\beta$ if and only if

$$\psi_1(x^{[0]}) = \beta \wedge x^{[1]} \in E \wedge \beta < \rho_1(\varphi_1(x^{[1]})).$$

Fix some $w \in \mathbb{R}$ so that $\psi_1(w) = \beta$. Then $x \in P_\beta$ if and only if

$$\psi_1(x^{[0]}) = \psi_1(w) \wedge (\exists^{\mathbb{R}} z)(\exists^{\mathbb{R}} v)(x^{[1]} \in E \wedge z \in E \wedge \varphi_1(x^{[1]}) = \varphi_1(z) \wedge R_1(z, v) \wedge \psi_1(w) < \psi_1(v)).$$

Using $\preceq_{\psi_1} \in \Delta^*$, $R_1 \in \Delta^*$, and the closure properties of Δ^* , one has that $P_\beta \in \Delta^*$ for any $\beta < \delta$ and thus $\text{rk}_W(P_\beta) < \kappa$. Let $P = \bigcup_{\beta < \delta} P_\beta$. Since κ is regular, $\sup\{\text{rk}_W(P_\beta) : \beta < \delta\} < \kappa$. Pick a set $D \in \Delta^*$ such that for all $\beta < \delta$, $\text{rk}_W(P_\beta) < \text{rk}_W(D)$. Define $S_2 \subseteq \mathbb{R} \times \mathbb{R}$ by $S_2(x, y)$ if and only if $P_{\psi_1(x)} = \Sigma_y^{-1}[D]$. By the Moschovakis coding lemma, there is a $R_2 \subseteq S_2$ with $R_2 \in \Delta^*$ so that for all $\alpha < \delta$, $R_2 \cap (\psi_1^{-1}[\{\alpha\}] \times \mathbb{R}) \neq \emptyset$. Then $x \in P$ if and only if

$$(\exists^{\mathbb{R}} w)(\exists^{\mathbb{R}} y)(R_2(w, y) \wedge \Sigma_y(x) \in D).$$

Thus $P \in \Delta^*$ and hence $\text{rk}_W(P) < \kappa$.

Define a norm $\varphi : P \rightarrow \delta$ by $\varphi(x) = \psi_1(x^{[0]})$. Since $\text{rk}_W(\preceq_{\psi_1}) < \kappa$ and $\text{rk}_W(P) < \kappa$, one has that $\text{rk}_W(\preceq_\varphi) < \kappa$. Let $T \subseteq P$ be Σ_2^1 . Let $T' = \{x : (\exists y)(y \in T \wedge x = y^{[1]})\}$. Since $T \subseteq P$, one has that T' is a Σ_2^1 subset of E . Since φ_1 is Σ_2^1 bounded, there is a $\zeta < \nu$ so that $\varphi_1[T'] \subseteq \zeta$. By definition of $P = \bigcup_{\beta < \delta} P_\beta$, one has that $\varphi[T] \subseteq \rho_1(\zeta) < \delta$. It has been shown that $\varphi : P \rightarrow \delta$ is a Σ_2^1 bounded prewellordering of length δ so that $\text{rk}_W(\preceq_\varphi) < \kappa$. This completes the proof. \square

Fix a coding of strategies by reals. If $x \in \mathbb{R}$, let $\rho_x : {}^{<\omega}\omega \rightarrow \omega$ denote a strategy on ω coded by x . If $w \in \text{WO}_{\geq \omega}$, then $\rho_x^w : {}^{<\omega}\text{ot}(w) \rightarrow \text{ot}(w)$ denote the strategy on $\text{ot}(w)$ which results from transferring ρ_x via the bijection $B_w : \omega \rightarrow \text{ot}(w)$ naturally induced from w . In this way, one says that (w, x) with $w \in \text{WO}$ and $x \in \mathbb{R}$ code the strategy ρ_x^w .

The following (in the ω case) is a well-known result concerning the unfolded Banach-Mazur game. If $2 \leq \alpha < \omega_1$, then one has that ${}^\omega\alpha$ is homeomorphic to ${}^\omega\omega$ so the result transfers to the countable ordinal α .

Fact 3.9. *Let $\alpha < \omega_1$. Let $A \subseteq {}^\omega\alpha$ and $B \subseteq {}^\omega\alpha \times {}^\omega\omega$. Consider the following game $G_{A,B}^*$ defined as follows.*

$G_{A,B}^*$	1	s_0	s_2	s_4	s_6	\dots	f
		s_1	s_3	s_5	s_7	\dots	
	2	z_0	z_1	z_3	z_4	\dots	z

For all $i \in \omega$, $s_i \in {}^{<\omega}\alpha$ and $z_i \in \omega$. Player 1 plays s_{2i} for all $i \in \omega$. Player 2 plays s_{2i+1} and z_i for all $i \in \omega$. Let $f = s_0 \hat{\ } s_1 \hat{\ } s_2 \dots$ and $z \in {}^\omega\omega$ be defined by $z(i) = z_i$. Player 2 wins $G_{A,B}^*$ if and only if $f \in A$ and $B(f, z)$.

If A is comeager in ${}^\omega\alpha$ and $A \subseteq \text{dom}(B)$, then Player 2 has a winning strategy. For every $w \in \text{WO}$ so that $\text{ot}(w) = \alpha$ and coding finite sequence of ordinals in α by elements of α , one can find some $x \in \mathbb{R}$ so that ρ_x^w is a Player 2 winning strategy for $G_{A,B}^*$.

Fact 3.10. *Assume $\text{ZF} + \text{AD} + \text{DC}_{\mathbb{R}}$. Let $\langle \delta_\alpha : \alpha < \omega_1 \rangle$ be such that $\sup\{\delta_\alpha : \alpha < \omega_1\} < \Theta$ and for all $\alpha < \omega_1$, $\text{cof}(\delta_\alpha) \geq \omega_2$. Then there is a sequence $\langle (P_\alpha, \preceq_\alpha) : \alpha < \omega_1 \rangle$ so that for all $\alpha < \omega_1$, \preceq_α is a prewellordering on P_α of length δ_α so that the associated surjective norm $\varphi_\alpha : P_\alpha \rightarrow \delta_\alpha$ is Σ_2^1 bounded.*

Proof. Let $\delta = \sup\{\delta_\alpha : \alpha < \omega_1\}$ which is less than Θ by assumption. Let \preceq_δ denote a prewellordering on \mathbb{R} of length δ . Let Γ be a nonselfdual pointclass closed under $\forall^{\mathbb{R}}, \exists^{\mathbb{R}}, \wedge, \vee$, having the prewellordering property, and containing \preceq_δ (for example, $\Sigma_1^{L(\preceq_\delta, \mathbb{R})}$). Then $o(\Gamma) > \delta$. Let A^* be a universal set in Γ and note that $\text{rk}_W(A^*) = o(\Gamma)$. Define a relation $S \subseteq \text{WO} \times \mathbb{R}$ by $S(w, x)$ if and only if $\Sigma_x^{-1}[A^*]$ is a Σ_2^1 bounded prewellordering of length $\delta_{\text{ot}(w)}$, where recall that $\Sigma_x : \mathbb{R} \rightarrow \mathbb{R}$ is the continuous function coded by the real x . Note that $\text{dom}(S) = \text{WO}$ since by Fact 3.8, for each $\alpha < \omega_1$, there is a Σ_2^1 bounded prewellordering (P, \preceq) of length δ_α so that $(P, \preceq) \leq_W A^*$. By the Moschovakis coding lemma, there is a Σ_2^1 relation $R' \subseteq S$ so that for all $\alpha < \omega_1$, $R' \cap (\text{WO}_\alpha \times \mathbb{R}) \neq \emptyset$. Define $R \subseteq \text{WO} \times \mathbb{R}$ by $R(w, x)$ if and only if $(\exists v)(v \in \text{WO} \wedge \text{ot}(v) = \text{ot}(w) \wedge R'(v, x))$. Note that R is Σ_2^1 and $\text{dom}(R) = \text{WO}$. Since AD can uniformize projective relations, let $\Phi : \text{WO} \rightarrow \mathbb{R}$ be a uniformization for R , which means for all $w \in \text{WO}$, $R(w, \Phi(w))$. For each $w \in \text{WO}$, let (Q_w, \preceq_w) denote the Σ_2^1 bounded prewellordering of length $\delta_{\text{ot}(w)}$ coded by $\Sigma_{\Phi(w)}^{-1}[A^*]$. Let $\varphi_w : Q_w \rightarrow \delta_{\text{ot}(w)}$ be the associated surjective norm of (Q_w, \preceq_w) .

For each $\alpha < \omega_1$, let P_α consists of the collection of (w, x) such that $w \in \text{WO}_\alpha$ and $x \in \mathbb{R}$ with the following properties.

- (1) ρ_x^w is a Player 2 strategy in games of the following form.

1	s_0	s_2	s_4	s_6	\dots	f
	s_1	s_3	s_5	s_7	\dots	
2	z_0	z_1	z_3	z_4	\dots	z

For each $i \in \omega$, $s_i \in {}^{<\omega}\alpha$. For each i , s_{2i} is played by Player 1 and s_{2i+1} is played by Player 2. For all $i \in \omega$, $z_i \in \omega$ and is played by Player 2. Let $f \in {}^\omega\alpha$ be defined by $f = s_0 \hat{\ } s_1 \hat{\ } s_2 \dots$ and $z \in \mathbb{R}$ be defined by $z(i) = z_i$ for all $i \in \omega$. (These are the same condition as the unfolded Banach-Mazur game on α from Fact 3.9 except that no payoff set has been specified.)

For each $p \in {}^\omega({}^{<\omega}\alpha)$, let $s_{2n}^p = p(n)$. Let $\langle s_{2n+1}^p : n \in \omega \rangle$ and $\langle z_i^p : i \in \omega \rangle$ be the response of Player 2 using ρ_x^w . Let $f(w, x, p) = s_0^p \hat{\ } s_1^p \hat{\ } s_2^p \dots$ and let $z(w, x, p) \in \mathbb{R}$ be defined by $z(w, x, p)(i) = z_i^p$.

- (2) There exists a $\beta_{w,x} < \delta_\alpha$ so that for all $p \in {}^\omega({}^{<\omega}\alpha)$,

$$z(w, x, p) \in Q_{\mathfrak{G}(f(w,x,p))} \quad \text{and} \quad \varphi_{\mathfrak{G}(f(w,x,p))}(z(w, x, p)) = \beta_{w,x}.$$

Define $\varphi_\alpha : P_\alpha \rightarrow \delta_\alpha$ by $\varphi_\alpha((w, x)) = \beta_{w, x}$. Let \preceq_α be the prewellordering on P_α induced from φ_α .

Claim 1: φ_α is surjective.

To see Claim 1: Fix a $\beta < \delta_\alpha$. Let $A = \text{surj}_\alpha$ which is a comeager subset of ${}^\omega\alpha$. Let $B \subseteq \text{surj}_\alpha \times \mathbb{R}$ be defined by $(f, z) \in B$ if and only if $z \in Q_{\mathfrak{G}(f)}$ and $\varphi_{\mathfrak{G}(f)}(z) = \beta$. Since $f \in \text{surj}_\alpha$ implies $\mathfrak{G}(f) = \alpha$, one has that $\text{dom}(B) = \text{surj}_\alpha = A$. Fact 3.9 implies that there is a (w, x) with $w \in \text{WO}_\alpha$ and $x \in \mathbb{R}$ so that ρ_x^w is a Player 2 winning strategy in $G_{A, B}^*$. From the definitions, one has that $(w, x) \in P_\alpha$ and $\varphi_\alpha((w, x)) = \beta$.

Claim 2: φ_α is Σ_2^1 bounded.

To see Claim 2: Recall the following basic fact about Banach-Mazur type games. For any $(w, x) \in P_\alpha$, there is a comeager set $K_{w, x}$ of $f \in {}^\omega\alpha$ so that there exists a $p : \omega \rightarrow <^\omega\alpha$ such that $f(w, x, p) = f$. For each $f \in K_{w, x}$, there is a canonical $p^{f, w, x} : \omega \rightarrow <^\omega\alpha$ so that $f(w, x, p^{f, w, x}) = f$. For a fixed f , there is a Borel function with the property that given a (w, x) such that $f \in K_{w, x}$, the function will output $p^{f, w, x}$.

Let $S \subseteq P_\alpha$ be Σ_2^1 . Fix an $f \in \text{surj}_\alpha$. Let $S_f = \{(w, x) \in S : (\exists p)(f(w, x, p) = f)\}$ which is a Σ_2^1 set. For any $(w, x) \in S_f \subseteq S \subseteq P_\alpha$, one has that $z(w, x, p^{f, w, x}) \in P_{\mathfrak{G}(f)}$. Thus $T_f = \{z(w, x, p^{f, w, x}) : (w, x) \in S_f\}$ is a subset of $P_{\mathfrak{G}(f)}$ and is Σ_2^1 . Since $\varphi_{\mathfrak{G}(f)}$ is Σ_2^1 bounded, let γ_f be the least ordinal below δ_α so that $\varphi_{\mathfrak{G}(f)}[T_f] \subseteq \gamma_f$.

For each $\gamma < \delta_\alpha$, let $B_\gamma = \{f \in \text{surj}_\alpha : \gamma_f = \gamma\}$ and $B_{<\gamma} = \{f \in \text{surj}_\alpha : \gamma_f < \gamma\}$. The claim is that there is a γ^* so that $B_{<\gamma^*}$ is comeager. Suppose not, then for all γ , $B_{<\gamma}$ is not comeager. Since $\text{surj}_\alpha \setminus B_{<\gamma} = \bigcup_{\gamma' \geq \gamma} B_{\gamma'}$ and wellordered union of meager sets are meager, one must have some $\gamma' \geq \gamma$ such that $B_{\gamma'}$ is nonmeager. Thus it has been shown that for all $\gamma < \delta_\alpha$, there exists a $\gamma' > \gamma$ so that $\gamma < \gamma' < \delta_\alpha$ and $B_{\gamma'}$ is nonmeager. Let ϵ_0 be the least ordinal $\epsilon > 0$ so that B_ϵ is nonmeager. Suppose $\beta < \omega_1$ and for all $\alpha < \beta$, ϵ_α has been defined. Let ϵ_β be the least ordinal ϵ greater than or equal to $\sup\{\epsilon_\alpha : \alpha < \beta\}$ so that B_ϵ is nonmeager. This defines a sequence $\langle B_\epsilon : \epsilon < \omega_1 \rangle$ of disjoint nonmeager subsets of ${}^\omega\alpha$. Since AD implies all sets of reals have the Baire property, the existence of this sequence contradicts the countable chain condition of the topology on ${}^\omega\alpha$. Thus it has been shown that there is a γ^* so that $B_{<\gamma^*}$ is comeager.

Now fix a $(w, x) \in S$. Since $K_{w, x}$ and $B_{<\gamma^*}$ are comeager, $K_{w, x} \cap B_{<\gamma^*} \neq \emptyset$. Let $f \in K_{w, x} \cap B_{<\gamma^*}$.

$$\varphi_\alpha((w, x)) = \beta_{w, x} = \varphi_{\mathfrak{G}(f(w, x, p^{f, w, x}))}(z(w, x, p^{f, w, x})) = \varphi_{\mathfrak{G}(f)}(z(w, x, p^{f, w, x})) < \gamma_f < \gamma^*$$

The second equation follows from the definition of $(w, x) \in P_\alpha$. The third equation comes from the fact that $f \in K_{w, x}$ and the definition of $p^{f, w, x}$. The first inequality follows from the fact that $z(w, x, p^{f, w, x}) \in T_f$. The second inequality is obtained from $f \in B_{<\gamma^*}$. Thus it has been shown that $\varphi_\alpha[S] \subseteq \gamma^*$, and so φ_α satisfies Σ_2^1 bounding. \square

4. GENERAL RESULT FOR ALL ORDINALS

The following is the good coding system for functions from $\epsilon < \omega_1$ into ω_1 . Such coding system are used to prove partition properties.

Definition 4.1. (Martin, [2] Fact 4.9) Let $\epsilon < \omega_1$. Fix $w^* \in \text{WO}$ so that w^* codes a wellordering with domain ω of ordertype ϵ (assuming without loss of generality that $\omega \leq \epsilon$). For $n \in \omega$, let $\text{ot}(w^*, n) \in \epsilon$ denote the rank of n in the wellordering w^* . For each $\alpha < \epsilon$, let $\text{num}(w^*, \alpha) \in \omega$ denote the integer n so that $\text{ot}(w^*, n) = \alpha$. Define $\text{decode} : \mathbb{R} \rightarrow \mathcal{P}(\epsilon \times \omega_1)$ by $\text{decode}(x)(\beta, \gamma)$ if and only if $x^{\text{num}(w^*, \beta)} \in \text{WO}_\gamma$. Define $\text{GC}_{\beta, \gamma} \subseteq \mathbb{R}$ by $x \in \text{GC}_{\beta, \gamma}$ if and only if $\text{decode}(x)(\beta, \gamma)$. Observe that if $x \in \text{GC}_{\beta, \gamma}$, then for any $\xi < \omega_1$, if $\text{decode}(x)(\beta, \xi)$ holds, then one must have that $\gamma = \xi$. By $\text{AC}_\mathbb{R}^\omega$, for any $f : \epsilon \rightarrow \omega_1$, there is some $x \in \mathbb{R}$ so that $\text{decode}(x)$ is the graph of f .

This defines a good coding system $(\Sigma_1^1, \text{decode}, \text{GC}_{\beta, \gamma} : \beta < \epsilon, \gamma < \omega_1)$ for ${}^\epsilon\omega_1$. Let $\text{GC} = \bigcap_{\beta < \epsilon} \bigcup_{\gamma < \omega_1} \text{GC}_{\beta, \gamma}$. Thus for all $x \in \text{GC}$, $\text{decode}(x)$ is the graph of a function $f : \epsilon \rightarrow \omega_1$.

Fact 4.2. ([1]) Fix $\epsilon < \omega_1$. Let $(\Sigma_1^1, \text{decode}, \text{GC}_{\beta, \gamma} : \beta < \epsilon, \gamma < \omega_1)$ be a fixed good coding system for ${}^\epsilon\omega_1$. For any club $D \subseteq \omega_1$, there is a club $C \subseteq D$ so that $\text{INC}^\epsilon(C)$, which is the set of $x \in \text{GC}$ so that $\text{decode}(x) \in [C]^\epsilon$, is Π_1^1 .

If $\epsilon < \omega_1$ and $f : \omega \cdot \epsilon \rightarrow \omega_1$, let $\text{block}(f) : \epsilon \rightarrow \omega_1$ be defined by $\text{block}(f)(\alpha) = \sup\{f(\omega \cdot \alpha + n) : n \in \omega\}$.

Fact 4.3. ([2] Theorem 3.8; [1]) (Almost Everywhere Good Code Uniformization) Assume ZF + AD. Let $\epsilon < \omega_1$ and $(\Sigma_1^1, \text{decode}, \text{GC}_{\beta, \gamma} : \beta < \omega \cdot \epsilon, \gamma < \omega_1)$ be a good coding system for ${}^{\omega \cdot \epsilon}\omega_1$. Suppose $R \subseteq [\omega_1]_*^\epsilon \times \mathbb{R}$.

Then there is a club $C \subseteq \omega_1$ and a Lipschitz continuous function $\Xi : \mathbb{R} \rightarrow \mathbb{R}$ so that for all $x \in \text{INC}^{\omega^\epsilon}(C)$, $R(\text{block}(\text{decode}(x)), \Xi(x))$.

Theorem 4.4. Assume $\text{ZF} + \text{AD} + \text{DC}_{\mathbb{R}}$. There is no injection of $[\omega_1]^{<\omega_1} \rightarrow {}^\omega\text{ON}$.

Proof. Suppose there is an injection $\Phi : [\omega_1]^{<\omega_1} \rightarrow {}^\omega\text{ON}$. By the Moschovakis coding lemma, there is a surjection $\pi : \mathbb{R} \rightarrow [\omega_1]^{<\omega_1}$. Define $\Psi : \mathbb{R} \times \omega \rightarrow \text{ON}$ by $\Psi(r, n) = \Phi(\pi(r))(n)$. Thus $\Psi[\mathbb{R} \times \omega]$ is a surjective image of \mathbb{R} . Thus there is a $\delta < \Theta$ so that $\Psi[\mathbb{R} \times \omega]$ is in bijection with δ . This implies that there is an injection $\Phi' : [\omega_1]^{<\omega_1} \rightarrow {}^\omega\delta$ where $\delta < \Theta$. Thus it suffices to show that there is no injection $\Phi : [\omega_1]^{<\omega_1} \rightarrow {}^\omega\delta$ where $\delta < \Theta$. For the sake of contradiction, fix such an injection Φ . For each $\epsilon < \omega_1$ and $n \in \omega$, let $\Phi_n^\epsilon : [\omega_1]^\epsilon \rightarrow \delta$ be defined by $\Phi_n^\epsilon(f) = \Phi(f)(n)$.

Claim: For each $\epsilon < \omega_1$ and $n \in \omega$, there is a club $C \subseteq \omega_1$ so that $|\Phi_n^\epsilon[[C]_*^\epsilon]| \leq \omega_1$.

Given the claim, the theorem follows: If C is a club, let $A_C^\epsilon = \{\xi < \delta : (\exists f \in [C]_*^\epsilon)(\exists n \in \omega)(\Phi(f)(n) = \xi)\}$. Fixing an $\epsilon < \omega_1$, one can use $\text{AC}_{\omega}^{\mathbb{R}}$ and the claim to show that there is a sequence $\langle C_n : n \in \omega \rangle$ so that for all $n \in \omega$, $|\Phi_n^\epsilon[[C_n]_*^\epsilon]| = \omega_1$. Letting $C = \bigcap_{n \in \omega} C_n$, one has that $A_C^\epsilon \subseteq \bigcup_{n \in \omega} \Phi_n^\epsilon[[C_n]_*^\epsilon]$ and hence $|A_C^\epsilon| \leq \omega_1$.

It has been shown that for all $\epsilon < \omega_1$, there is a club C so that $|A_C^\epsilon| \leq \omega_1$. By Fact 2.7, there is a sequence $\langle C_\epsilon : \epsilon < \omega_1 \rangle$ so that for all $\epsilon < \omega_1$, $|A_{C_\epsilon}^\epsilon| \leq \omega_1$. Let $T = \bigcup_{\epsilon < \omega_1} A_{C_\epsilon}^\epsilon$ and note that $|T| = \omega_1$. Observe that if $f \in \bigcup_{\epsilon < \omega_1} [C_\epsilon]_*^\epsilon$, then $\Phi(f) \in {}^\omega T$. Since Φ is an injection, $|\bigcup_{\epsilon < \omega_1} [C_\epsilon]_*^\epsilon| = |[\omega_1]^{<\omega_1}|$, and $|T| = \omega_1$, one has that Φ induces an injection of $[\omega_1]^{<\omega_1}$ into ${}^\omega\omega_1$. This is impossible by Fact 2.6.

Thus it remains to show the claim: Now fix $\epsilon < \omega_1$ and $n \in \omega$. Let $S_{-1} = \{\delta\}$. Suppose $k \in \{-1\} \cup \omega$ and $S_k \subseteq \delta + 1$ has been defined with $|S_k| \leq \omega_1$. Let $S_k^0 = \{\gamma \in S_k : \text{cof}(\gamma) \leq \omega_1\}$ and let $S_k^1 = \{\gamma \in S_k : \text{cof}(\gamma) > \omega_1\}$. Let $\iota_0 : \omega_1 \rightarrow S_k^0$ and $\iota_1 : \omega_1 \rightarrow S_k^1$ be surjections.

Let $\nu_\xi = \iota_0(\xi)$. Applying Fact 3.3 to $\langle \nu_\xi : \xi < \omega_1 \rangle$, there is a sequence $\langle K_\xi : \xi < \omega_1 \rangle$ so that for all $\xi < \omega_1$, $K_\xi \subseteq \nu_\xi$, $|K_\xi| \leq \omega_1$, and $\sup K_\xi = \nu_\xi$.

Let $\delta_\xi = \iota_1(\xi)$. Applying Fact 3.10 to $\langle \delta_\xi : \xi < \omega_1 \rangle$, there is a sequence $\langle (P_\xi, \preceq_\xi) : \xi < \omega_1 \rangle$ of prewellorderings so that for each ξ , (P_ξ, \preceq_ξ) is a Σ_2^1 bounded prewellordering of length δ_ξ with $\varphi_\xi : P_\xi \rightarrow \delta_\xi$ being its associated surjective norm.

Fix an $f \in [\omega_1]_*^\epsilon$. Consider the relation $S_f \subseteq \text{WO} \times \mathbb{R}$ defined by $S_f(w, x)$ if and only if $(x \in P_{\text{ot}(w)}) \wedge (\Phi_n^\epsilon(f) < \varphi_{\text{ot}(w)}(x))$. Note that $w \in \text{dom}(S_f)$ if and only if $\Phi_n^\epsilon(f) < \delta_{\text{ot}(w)}$. Fix a Σ_2^1 set $U \subseteq \mathbb{R} \times \mathbb{R}^2$ which is universal for Σ_2^1 subsets of \mathbb{R}^2 . By the Moschovakis coding lemma, there is a $z \in \mathbb{R}$ so that $U_z \subseteq S_f$ and for all $\xi < \omega_1$, $U_z \cap (\text{WO}_\xi \times \mathbb{R}) \neq \emptyset$ if and only if $S_f \cap (\text{WO}_\xi \times \mathbb{R}) \neq \emptyset$. Such a $z \in \mathbb{R}$ will be called an f -selector.

Define $T \subseteq [\omega_1]_*^\epsilon \times \mathbb{R}$ by $T(f, z)$ if and only if z is an f -selector. Note that $\text{dom}(T) = [\omega_1]_*^\epsilon$. After fixing a good coding system $(\Sigma_1^1, \text{decode}, \text{GC}_{\beta, \gamma} : \beta < \omega \cdot \epsilon, \gamma < \omega_1)$ for ${}^{\omega^\epsilon}\omega_1$ and using Fact 4.3, there is a club $D \subseteq \omega_1$ and a Lipschitz function $\Xi : \mathbb{R} \rightarrow \mathbb{R}$ so that for all $x \in \text{Inc}^{\omega^\epsilon}(D)$, $T(\text{block}(\text{decode}(x)), \Xi(x))$. By Fact 4.2, there is a $C' \subseteq D$ so that $\text{Inc}^{\omega^\epsilon}(C')$ is Π_1^1 . Thus $\Xi[\text{Inc}^{\omega^\epsilon}(C')]$ is a Σ_2^1 set. Let $C \subseteq C'$ be the collection of limit points of C' . Fix a $\xi < \omega_1$. Let

$$V_\xi = \{x : (\exists z)(\exists w \in \text{WO}_\xi)(z \in \Xi[\text{Inc}^{\omega^\epsilon}(C')] \wedge U_z(w, x))\}.$$

V_ξ is Σ_2^1 and $V_\xi \subseteq P_\xi$ since all elements of $\Xi[\text{Inc}^{\omega^\epsilon}(C')]$ are f -selectors. Since P_ξ is Σ_2^1 bounded, there is a $\gamma < \delta_\xi$ so that $\varphi_\xi[V_\xi] \subseteq \gamma$. Note that if $f \in [C]_*^\epsilon$, then there is a $u \in \text{Inc}^{\omega^\epsilon}(C')$ so that $\text{block}(\text{decode}(u)) = f$. Then $\Xi(u)$ is an f -selector. If $\Phi_n^\epsilon(f) < \delta_\xi$, then there is some $w \in \text{WO}_\xi$ and $x \in P_\xi$ so that $U_{\Xi(u)}(w, x)$. Hence $x \in V_\xi$ and therefore $\Phi_n^\epsilon(f) < \varphi_\xi(x) < \gamma < \delta_\xi$.

Thus it has been shown there is a club C so that for all $\xi < \omega_1$, there is an ordinal $\gamma < \delta_\xi$ so that $\Phi_n^\epsilon(f) < \gamma$ or $\Phi_n^\epsilon(f) \geq \delta_\xi$ for all $f \in [C]_*^\epsilon$. For each $\xi < \omega_1$, let δ_ξ^* be the least such ordinal $\gamma < \delta_\xi$ so that there is a club C so that for all $f \in [C]_*^\epsilon$, $\Phi_n^\epsilon(f) < \gamma$ or $\Phi_n^\epsilon(f) \geq \delta_\xi^*$.

Now let $S_{k+1} = (\bigcup_{\xi < \omega_1} K_\xi) \cup \{\delta_\xi^* : \xi < \omega_1\}$. Note that S_{k+1} has the following property.

- (1) $S_{k+1} \subseteq \delta$ and $|S_{k+1}| = \omega_1$.
- (2) If $\xi \in S_k$ and $\text{cof}(\xi) \leq \omega_1$, then $\sup(S_{k+1} \cap \xi) = \xi$.
- (3) There is a club C so that for all $\xi \in S_k$ with $\text{cof}(\xi) > \omega_1$, there is a $\xi^* < \xi$ with $\xi^* \in S_{k+1}$ so that for all $f \in [C]_*^\epsilon$, $\Phi_n^\epsilon(f) < \xi^*$ or $\Phi_n^\epsilon(f) \geq \xi$.

The construction of S_{k+1} depends on $S_k \subseteq \delta$ and the surjections ι_0 and ι_1 . Since $\delta < \Theta$, there is a surjection of \mathbb{R} onto $\mathcal{P}(\delta)$. Thus $\text{DC}_{\mathbb{R}}$ is sufficient to create a sequence $\langle S_k : k \in \omega \rangle$ so that the relation between S_k and S_{k+1} is as specified above. Let $S = \bigcup_{k < \omega} S_k$ and observe that $|S| = \omega_1$.

Using $\text{AC}_\omega^\mathbb{R}$, there is a sequence $\langle C_k : k \in \omega \rangle$ which witnesses (3) for S_k (recall $S_{-1} = \{\delta\}$). Let $C = \bigcap_{k \in \omega} C_k$, and one will show that $\Phi_n^\epsilon[[C]_*^\epsilon] \subseteq S$. So suppose otherwise that there is an $f \in [C]_*^\epsilon$ so that $\Phi_n^\epsilon(f) \notin S$. Let $\nu = \Phi_n^\epsilon(f)$. Let $\nu_{-1} = \delta$. Suppose $\nu_k \in S_k$ has been defined with $\nu < \nu_k$. If $\text{cof}(\nu_k) \leq \omega_1$, then (2) implies there is some ordinal $\zeta \in S_{k+1}$ such that $\nu < \zeta < \nu_k$. If $\text{cof}(\nu_k) > \omega_1$, then by (3) for S_k , there is a $\nu_k^* \in S_{k+1}$ so that $\Phi_n^\epsilon(f) = \nu < \nu_k^* < \nu_k$. Thus in either case, let ν_{k+1} be the least ordinal $\zeta \in S_{k+1}$ so that $\nu < \zeta < \nu_k$. Then $\langle \nu_k : k \in \omega \rangle$ is an infinite descending sequence of ordinals which is impossible.

It has been shown that there is a club $C \subseteq \omega_1$ so that $\Phi_n^\epsilon[[C]_*^\epsilon] \subseteq S$. Since $|S| \leq \omega_1$, $|\Phi_n^\epsilon[[C]_*^\epsilon]| \leq \omega_1$. This proves the claim. \square

The next result is the almost everywhere continuity property for functions of the form $\Phi : [\omega_1]^{\omega_1} \rightarrow \omega_1$. It follows from the almost everywhere short length club uniformization for relations of the form $R \subseteq [\omega_1]_*^{<\omega_1} \times \text{club}_{\omega_1}$ which are \subseteq -downward closed in the club_{ω_1} coordinate ([4] Theorem 3.10) proved under AD. Unlike Fact 2.7, the everywhere version of this uniformization fails in $L(\mathbb{R}) \models \text{AD}$ ([4] Fact 3.9); however, the everywhere version does hold under $\text{ZF} + \text{AD}_\mathbb{R}$ ([4] Theorem 3.7).

Fact 4.5. ([4] Theorem 4.5) *Assume $\text{ZF} + \text{AD}$. Let $\Phi : [\omega_1]^{\omega_1} \rightarrow \omega_1$. Then there is a club $C \subseteq \omega_1$ so that $\Phi \upharpoonright [C]_*^{\omega_1}$ is continuous. That is, for all $f \in [C]_*^{\omega_1}$, there is an $\alpha < \omega_1$ so that for all $g \in [C]_*^{\omega_1}$, if $f \upharpoonright \alpha = g \upharpoonright \alpha$, then $\Phi(f) = \Phi(g)$.*

Fact 4.6. ([4] Theorem 4.6) *Assume $\text{ZF} + \text{AD}$. For all functions $\Phi : [\omega_1]_*^{\omega_1} \rightarrow \omega_1$, there is an $\alpha < \omega_1$ so that $|\Phi^{-1}[\{\alpha\}]| = |[\omega_1]_*^{\omega_1}|$.*

Proof. By Fact 4.5, there is a club so that $\Phi \upharpoonright [C]_*^{\omega_1}$ is continuous. Pick any $f \in [C]_*^{\omega_1}$ and let $\beta = \Phi(f)$. By continuity, there is an $\alpha < \omega_1$ so that for all $g \in [C]_*^{\omega_1}$ with $f \upharpoonright \alpha = g \upharpoonright \alpha$, $\Phi(g) = \Phi(f) = \beta$. Let $N_{f \upharpoonright \alpha}^C = \{g \in [C]_*^{\omega_1} : g \upharpoonright \alpha = f \upharpoonright \alpha\}$. Note that $|N_{f \upharpoonright \alpha}^C| = |[\omega_1]_*^{\omega_1}|$ and $N_{f \upharpoonright \alpha}^C \subseteq \Phi^{-1}[\{\beta\}]$. Thus $|\Phi^{-1}[\{\beta\}]| = |[\omega_1]_*^{\omega_1}|$. \square

The above argument is inefficient since it uses the almost everywhere continuity property (Fact 4.5) which follows from a suitable almost everywhere club uniformization property. The proof of this club uniformization property at a cardinal δ requires more than δ being a strong partition cardinal and even more than the existence of a good coding system for ${}^\delta\delta$. [1] isolates a strengthening of the good coding system sufficient to prove the necessary club uniformization and the almost everywhere continuity property. However, Fact 4.6 can be proved by purely combinatorial arguments using only the strong partition property. In particular, [1] shows that under ZF, if δ is a cardinal so that $\delta \rightarrow_* (\delta)_2^\delta$, $\kappa \in \text{ON}$, and $\Phi : [\delta]_*^\delta \rightarrow \kappa$, then there exists an $\alpha < \kappa$ so that $|\Phi^{-1}[\{\alpha\}]| = |[\delta]_*^\delta|$.

Theorem 4.7. *Assume $\text{ZF} + \text{AD} + \text{DC}_\mathbb{R}$. There is no injection of $[\omega_1]^{\omega_1}$ into ${}^{<\omega_1}\text{ON}$.*

Assuming just $\text{ZF} + \text{AD}$, $\neg(|[\omega_1]^{\omega_1}| \leq |{}^{<\omega_1}(\omega_\omega)|)$.

Proof. Since \mathbb{R} surjects onto $[\omega_1]^{\omega_1}$ by the Moschovakis coding lemma, any injection $\Upsilon : [\omega_1]^{\omega_1} \rightarrow {}^{<\omega_1}\text{ON}$ induces an injection $\Phi : [\omega_1]^{\omega_1} \rightarrow {}^{<\omega_1}\delta$ for some $\delta < \Theta$. If $\sigma \in {}^{<\omega_1}\delta$, then let $\text{length}(\sigma) = |\sigma|$. Let $\Psi : [\omega_1]^{\omega_1} \rightarrow \omega_1$ be defined by $\Psi = \text{length} \circ \Phi$. By Fact 4.6 and the fact that $|[\omega_1]_*^{\omega_1}| = |[\omega_1]^{\omega_1}|$, there is an $\epsilon < \omega_1$ so that $|\Psi^{-1}[\{\epsilon\}]| = |[\omega_1]^{\omega_1}|$. So $\Phi : \Psi^{-1}[\{\epsilon\}] \rightarrow {}^\epsilon\delta$ induces an injection from $[\omega_1]^{<\omega_1}$ into $\omega\delta$ since $|[\omega_1]^{<\omega_1}| < |[\omega_1]^{\omega_1}|$ and $|\epsilon\delta| = |\omega\delta|$. However this is impossible by Theorem 4.4. The second result is the same argument using Theorem 2.9. \square

REFERENCES

1. William Chan, *Definable combinatorics of strong partition cardinals*, In preparation.
2. ———, *An introduction to combinatorics of determinacy*, Trends in Set Theory, vol. 752, 2020, pp. 21–75.
3. William Chan and Stephen Jackson, *Applications of infinity-borel codes to definability and definable cardinals*, Submitted.
4. William Chan and Stephen Jackson, *Definable combinatorics at the first uncountable cardinal*, Trans. Amer. Math. Soc. **374** (2021), no. 3, 2035–2056.
5. William Chan, Stephen Jackson, and Nam Trang, *More definable combinatorics around the first and second uncountable cardinal*, Submitted.
6. Steve Jackson, *Structural consequences of AD*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 1753–1876. MR 2768700
7. Alexander S. Kechris, *AD and projective ordinals*, Cabal Seminar 76–77 (Proc. Caltech-UCLA Logic Sem., 1976–77), Lecture Notes in Math., vol. 689, Springer, Berlin, 1978, pp. 91–132. MR 526915

8. Alexander S. Kechris, Robert M. Solovay, and John R. Steel, *The axiom of determinacy and the prewellordering property*, Cabal Seminar 77–79 (Proc. Caltech-UCLA Logic Sem., 1977–79), Lecture Notes in Math., vol. 839, Springer, Berlin-New York, 1981, pp. 101–125. MR 611169
9. Alexander S. Kechris and W. Hugh Woodin, *Generic codes for uncountable ordinals, partition properties, and elementary embeddings*, Games, scales, and Suslin cardinals. The Cabal Seminar. Vol. I, Lect. Notes Log., vol. 31, Assoc. Symbol. Logic, Chicago, IL, 2008, pp. 379–397. MR 2463619
10. John R. Steel, *Closure properties of pointclasses*, Cabal Seminar 77–79 (Proc. Caltech-UCLA Logic Sem., 1977–79), Lecture Notes in Math., vol. 839, Springer, Berlin-New York, 1981, pp. 147–163. MR 611171
11. W. Hugh Woodin, *The cardinals below $|\omega_1|^{<\omega_1}$* , Ann. Pure Appl. Logic **140** (2006), no. 1-3, 161–232. MR 2224057
12. ———, *The axiom of determinacy, forcing axioms, and the nonstationary ideal*, revised ed., De Gruyter Series in Logic and its Applications, vol. 1, Walter de Gruyter GmbH & Co. KG, Berlin, 2010. MR 2723878

DEPARTMENT OF MATHEMATICS, CARNEGIE MELLON UNIVERSITY, PITTSBURGH, PA 15213
Email address: `wchan3@andrew.cmu.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH TEXAS, DENTON, TX 76203
Email address: `Stephen.Jackson@unt.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH TEXAS, DENTON, TX 76203
Email address: `Nam.Trang@unt.edu`