

IRREDUCIBILITY CRITERIA FOR LOCAL AND GLOBAL REPRESENTATIONS

HIRO-AKI NARITA, AMEYA PITALE, AND RALF SCHMIDT

(Communicated by Kathrin Bringmann)

ABSTRACT. It is proved that certain types of modular cusp forms generate irreducible automorphic representations of the underlying algebraic group. Analogous Archimedean and non-Archimedean local statements are also given.

INTRODUCTION

One of the motivations for this paper was to show that full level cuspidal Siegel eigenforms generate irreducible, automorphic representations of the adelic symplectic similitude group. Such a result is well known for the case of classical elliptic modular forms. Given an elliptic cusp form f (of some weight and some level), an adelic function Φ_f can be constructed (see [Ge], §5), which is a cuspidal automorphic form on the adelic group $\mathrm{GL}(2, \mathbb{A})$; here, \mathbb{A} denotes the ring of adèles of \mathbb{Q} . Let V_f be the space of automorphic forms generated by all right translates of Φ_f . In this classical situation it turns out that V_f is irreducible precisely when f is an eigenform for the Hecke operators T_p for almost all primes p . The proof uses the strong multiplicity one property for cuspidal automorphic representations of $\mathrm{GL}(2)$.

For most other types of modular forms, strong multiplicity one, or even weak multiplicity one, is not available. The goal of this paper is to show that, under certain circumstances, the automorphic representation V_f is still irreducible, even if multiplicity one is not known. Loosely speaking, this is the case whenever f is a holomorphic type of modular form and is an eigenfunction for *all* Hecke operators. See Corollary 3.2 for a precise statement. We stress that this result does not prove multiplicity one for full level automorphic forms; for example, under our current state of knowledge, it is still conceivable that two holomorphic Siegel cusp forms of degree $n > 1$ have the same weight and the same Hecke eigenvalues for all primes p , yet are linearly independent.

While our results are applicable mainly in a reductive setting, we keep the definitions general enough to include certain non-reductive situations, such as Jacobi forms. For completeness we also include analogous local Archimedean and non-Archimedean irreducibility criteria. All of this is well known to experts, but for

Received by the editors June 6, 2011.

2010 *Mathematics Subject Classification*. Primary 11F46, 11F50, 11F70; Secondary 22E50, 22E55.

Key words and phrases. Representation over local fields, automorphic representation, Hecke eigenvector.

The first author was partly supported by Grant-in-Aid for Young Scientists (B) 21740025, the Ministry of Education, Culture, Sports, Science and Technology, Japan.

©2012 American Mathematical Society
Reverts to public domain 28 years from publication

lack of a good reference we found it useful to collect the relevant results in this paper.

1. LOCAL NON-ARCHIMEDEAN THEORY

Let G be a group of td-type, as in [Ca]. We fix a left-invariant Haar measure on G . The Hecke algebra $\mathcal{H}(G)$ consists of the locally constant, compactly supported functions $f : G \rightarrow \mathbb{C}$, with product given by

$$(f_1 * f_2)(g) = \int_G f_1(h) f_2(h^{-1}g) dh.$$

Given an open-compact subgroup K of G , the symbol $\mathcal{H}(G, K)$ denotes the subalgebra of $\mathcal{H}(G)$ consisting of left and right K -invariant functions. If A is a subset of G , let I_A be the characteristic function of A . Then $\text{vol}(K)^{-1}I_K$ is an identity element of $\mathcal{H}(G, K)$. As usual, a representation π of G on a complex vector space V is called *smooth* if every $v \in V$ is stabilized by some open-compact K . In this case $\mathcal{H}(G)$ acts on V via

$$\pi(f)v = \int_G f(g)\pi(g)v dg.$$

The algebra $\mathcal{H}(G, K)$ acts on the space V^K of K -invariant vectors. We say that $v \in V^K$ is an eigenvector for $\mathcal{H}(G, K)$ if for all $f \in \mathcal{H}(G, K)$ there exists a scalar $\lambda(f)$ such that $\pi(f)v = \lambda(f)v$.

Lemma 1.1. *Let (π, V) be a smooth representation of G . Let K be an open-compact subgroup of G . Then, for $v \in V$ and $g \in G$,*

$$\int_K \int_K \pi(k_1 g k_2) v dk_2 dk_1 = \frac{\text{vol}(K)^2}{\text{vol}(KgK)} \pi(I_{KgK})v.$$

Proof. Let $KgK = \bigsqcup_i g_i K$, a finite disjoint union. Then

$$\pi(I_{KgK})v = \sum_i \int_{g_i K} \pi(h)v dh = \sum_i \int_K \pi(g_i k_2)v dk_2.$$

Applying $\pi(k_1)$ to both sides and integrating over K , we obtain

$$\text{vol}(K)\pi(I_{KgK})v = \sum_i \int_K \int_K \pi(k_1 g_i k_2)v dk_2 dk_1 = \sum_i \int_K \int_K \pi(k_1 g k_2)v dk_2 dk_1.$$

The assertion follows. □

Proposition 1.2. *Let (π, V) be a smooth representation of G such that every G -invariant subspace has a G -invariant complement.¹ Let K be an open-compact subgroup of G , and let $v_0 \in V^K$ be a non-zero vector with the following properties.*

- i) *The vectors $\pi(g)v_0$, where g runs through G , span V .*
- ii) *v_0 is an eigenvector for $\mathcal{H}(G, K)$.*

Then the representation π is irreducible.

¹For example, this is satisfied if π is unitarizable.

Proof. We first show that V^K is spanned by v_0 . Let $v \in V^K$. By hypothesis i), we can write

$$v = \sum_{i=1}^n c_i \pi(g_i)v_0 \quad \text{for some } c_i \in \mathbb{C}, g_i \in G.$$

Since v_0 is K -invariant,

$$v = \frac{1}{\text{vol}(K)} \sum_{i=1}^n c_i \int_K \pi(g_i k_2)v_0 dk_2.$$

Applying $\pi(k_1)$ and integrating over K , we obtain

$$v = \frac{1}{\text{vol}(K)^2} \sum_{i=1}^n c_i \int_K \int_K \pi(k_1 g_i k_2)v_0 dk_2 dk_1.$$

By Lemma 1.1 and hypothesis ii), the right hand side is a multiple of v_0 . This proves our claim that V^K is spanned by v_0 .

Now let W be a G -invariant subspace of V , and let W' be a G -invariant complement. Write

$$v_0 = w + w', \quad w \in W, w' \in W'.$$

Since v_0 is K -invariant, the same must be true for w and w' . By what we already proved, w and w' are both multiples of v_0 . It follows from the first hypothesis that V is entirely contained in W or in W' . This proves that V is irreducible. \square

Remark. Let F be a p -adic field with ring of integers \mathfrak{o} . Let $G = \text{GL}(2, F)$ and $K = \text{GL}(2, \mathfrak{o})$. Let $V = ||^{1/2} \times ||^{-1/2}$ be the parabolically induced representation of G which has the Steinberg representation as its unique subrepresentation and the trivial representation as its unique quotient. Then $\dim V^K = 1$, and the translates of a non-zero $v_0 \in V^K$ span all of V , yet V is reducible. This example shows that the hypothesis about invariant complements in Proposition 1.2 is not indispensable.

2. LOCAL ARCHIMEDEAN THEORY

Let G be a real Lie group and K a subgroup. We assume that K is of the form center times compact subgroup.² An equivalence class of irreducible (finite-dimensional) representations of K will be called a K -type. With \mathfrak{g} denoting the Lie algebra of G , we have the following simple irreducibility criterion.

Lemma 2.1. *Let (π, V) be a (\mathfrak{g}, K) -module such that every invariant subspace has an invariant complement (for example, this is satisfied if π is unitarizable). Let τ be a K -type with the following properties:*

- i) V is spanned by the vectors $\pi(X)v$, where X runs through $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ and v runs through the τ -isotypical component of V .
- ii) V contains the K -type τ exactly once.

Then π is irreducible.

Proof. Let W be an invariant subspace of V . Let W' be an invariant complement. By ii), exactly one of W or W' contains the K -type τ . Then i) implies that either $V = W$ or $V = W'$. \square

²The reason for not assuming that K itself is compact will become apparent once we consider the Jacobi forms example below.

As above let \mathfrak{g} be the Lie algebra of G , and let \mathfrak{k} be the Lie algebra of K . Let $\mathfrak{g}_{\mathbb{C}}$ and $\mathfrak{k}_{\mathbb{C}}$ be their complexifications. We will assume that K is connected, so that K -types are in one-to-one correspondence with the equivalence classes of irreducible representations of $\mathfrak{k}_{\mathbb{C}}$. We will assume that $\mathfrak{g}_{\mathbb{C}}$ admits a direct sum decomposition

$$(1) \quad \mathfrak{g}_{\mathbb{C}} = \mathfrak{p}_{\mathbb{C}}^{\pm} + \mathfrak{k}_{\mathbb{C}} + \mathfrak{p}_{\mathbb{C}}^{-}$$

with Lie subalgebras $\mathfrak{p}_{\mathbb{C}}^{\pm}$ and $\mathfrak{p}_{\mathbb{C}}^{-}$ such that $[\mathfrak{k}_{\mathbb{C}}, \mathfrak{p}_{\mathbb{C}}^{\pm}] \subset \mathfrak{p}_{\mathbb{C}}^{\pm}$. Given a K -type τ , we extend τ to a representation of $\mathfrak{k}_{\mathbb{C}} + \mathfrak{p}_{\mathbb{C}}^{-}$ on which $\mathfrak{p}_{\mathbb{C}}^{-}$ acts trivially and define the $\mathfrak{g}_{\mathbb{C}}$ -module

$$(2) \quad L_{\tau} := \mathcal{U}(\mathfrak{g}_{\mathbb{C}}) \otimes_{\mathcal{U}(\mathfrak{k}_{\mathbb{C}} + \mathfrak{p}_{\mathbb{C}}^{-})} \tau.$$

Proposition 2.2. *Let G and K be as above. Let (π, V) be a (\mathfrak{g}, K) -module such that every invariant subspace has an invariant complement. Let v_0 be a vector in V with the following properties:*

- i) V is spanned by the vectors $\pi(X)v_0$, where X runs through $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$.
- ii) The span of the vectors $\pi(k)v_0$, where k runs through K , is an irreducible representation τ of K .
- iii) The K -type τ occurs in L_{τ} exactly once.
- iv) $\pi(\mathfrak{p}_{\mathbb{C}}^{-})v_0 = 0$.

Then π is irreducible.

Proof. Let V_{τ} be the span of the vectors $\pi(k)v_0$, where k runs through K . By iv), $\mathfrak{p}_{\mathbb{C}}^{-}$ annihilates all of V_{τ} . Therefore, there exists a map

$$(3) \quad L_{\tau} \cong \mathcal{U}(\mathfrak{g}_{\mathbb{C}}) \otimes_{\mathcal{U}(\mathfrak{k}_{\mathbb{C}} + \mathfrak{p}_{\mathbb{C}}^{-})} V_{\tau} \longrightarrow V$$

given by $X \otimes v \mapsto \pi(X)v$. By i), this map is surjective. Hypothesis iii) implies that V contains the K -type τ only once. Hence, we can apply Lemma 2.1 to see that π is irreducible. \square

Remark. Hypothesis iii) is satisfied if G is linear, simple and connected, of Hermitian type, K is maximal compact, $\mathfrak{p}_{\mathbb{C}}^{\pm}$ (resp. $\mathfrak{p}_{\mathbb{C}}^{-}$) denotes the space spanned by the root vectors for the non-compact positive (resp. negative) roots, and τ is equivalent to the minimal K -type in a holomorphic discrete series representation.

As an application of Proposition 2.2, we consider holomorphic Jacobi forms on $\mathcal{H} \times \mathbb{C}$, as in [EZ]. Let $G^J = \mathrm{SL}(2, \mathbb{R}) \ltimes H(\mathbb{R})$ be the real Jacobi group, where H is a three-dimensional Heisenberg group; see [BS], Sect. 1.1, for the precise definition. Let K be the subgroup $\mathrm{SO}(2) \times Z$, where $Z \cong \mathbb{R}$ is the center of the Heisenberg group. Then, by [BS], Sect. 1.4, the complexified Lie algebra $\mathfrak{g}_{\mathbb{C}}$ of G^J admits a decomposition of the type (1).

Now let f be a Jacobi form of weight k and index m . We associate to f the function $\phi_f : G^J \rightarrow \mathbb{C}$ given by $\phi_f(g) = (f|_{k,m}g)(i, 0)$, where the slash action is the same as in Definition 4.1.1 of [BS]. Then the holomorphy of f is equivalent to $R(\mathfrak{p}_{\mathbb{C}}^{-})\phi_f = 0$, where R denotes right translation; see Proposition 4.1.2 of [BS]. Moreover, ϕ transforms by scalars under right translation by elements of K . Condition iii) of Proposition 2.2 is also satisfied, as is apparent from the description of the lowest weight representation $\pi_{k,m}^+$ of G^J in [BS], Sect. 3.1. Therefore, we obtain the following corollary (which could be generalized to higher degree Jacobi forms).

Corollary 2.3. *Let f be a holomorphic Jacobi form on $\mathcal{H} \times \mathbb{C}$ of weight k and index m . Then, under right translation, the associated function $\phi_f : G^J \rightarrow \mathbb{C}$ generates an irreducible (\mathfrak{g}, K) -module.*

We also seek an irreducibility criterion for non-connected Lie groups; this is important for certain global applications, which we will give below. Hence assume that G is a Lie group with connected component G_0 such that G_0 has finite index in G . Let K be a compact subgroup of G whose connected component K_0 has finite index in K . If (π, V) is a $\mathfrak{g}_{\mathbb{C}}$ -module and $\sigma \in K$ is an element, let (π^σ, V) be the $\mathfrak{g}_{\mathbb{C}}$ -module with underlying space V and action given by

$$(4) \quad \pi^\sigma(X)v = \pi(\text{Ad}(\sigma)X)v, \quad X \in \mathfrak{g}_{\mathbb{C}}, v \in V.$$

Again we assume that $\mathfrak{g}_{\mathbb{C}}$ admits a decomposition as in (1).

Proposition 2.4. *Let G and K be as above. Let (π, V) be a (\mathfrak{g}, K) -module such that every invariant subspace has an invariant complement. Let v_0 be a vector in V with the following properties:*

- i) V is spanned by the vectors $\pi(X)\pi(k)v_0$, where X runs through $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ and k runs through K .
- ii) The span of the vectors $\pi(k)v_0$, where k runs through K_0 , is an irreducible representation τ of K_0 .
- iii) The K_0 -type τ occurs in L_τ (defined as in (2)) exactly once.
- iv) For $\sigma \in K$, $\sigma \notin K_0$, the $\mathfrak{g}_{\mathbb{C}}$ -modules L_τ and $(L_\tau)^\sigma$ have no K_0 -type in common.
- v) $\pi(\mathfrak{p}_{\mathbb{C}}^-)v_0 = 0$.

Then π is irreducible.

Proof. By i),

$$V = \sum_{\sigma \in K/K_0} \pi(\sigma)\pi(\mathcal{U}(\mathfrak{g}_{\mathbb{C}}))v_0.$$

As in the proof of Proposition 2.2, there is a surjective map $L_\tau \rightarrow \pi(\mathcal{U}(\mathfrak{g}_{\mathbb{C}}))v_0$. Hence, by iii), the K_0 -type τ occurs in $\pi(\mathcal{U}(\mathfrak{g}_{\mathbb{C}}))v_0$ exactly once. Using iv), it follows that the K_0 -type τ occurs in all of V exactly once.

Now let $W \subset V$ be an invariant (\mathfrak{g}, K) -submodule, and let W' be an invariant complement. Then exactly one of W or W' contains the K_0 -type τ , and hence the vector v_0 . It follows from i) that either $V = W$ or $V = W'$. \square

In this section we have kept our hypotheses general in order to accommodate examples such as that of Jacobi forms. For background on highest weight modules and irreducibility in the semisimple setting, see Sect. 7 of [Ya].

3. GLOBAL THEORY

Let F be a number field, and let G be an algebraic group defined over F . Let Z be the center of G . Let \mathbb{A} be the ring of adeles of F , and let \mathbb{A}_∞ and \mathbb{A}_f be the subrings of Archimedean and finite adeles, respectively. Then we have the adelicized groups $G(\mathbb{A})$, $G(\mathbb{A}_\infty)$ and $G(\mathbb{A}_f)$. Let K_∞ be a fixed maximal compact subgroup of $G(\mathbb{A}_\infty)$. For present purposes, we call a function $\Phi : G(\mathbb{A}) \rightarrow \mathbb{C}$ an automorphic form on $G(\mathbb{A})$ if the following conditions are satisfied.

- i) Φ is smooth; i.e., Φ is right invariant under some open-compact subgroup K_f of $G(\mathbb{A}_f)$, and for any fixed $h \in G(\mathbb{A})$ the function $G(\mathbb{A}_\infty) \ni g \mapsto \Phi(gh)$ is C^∞ .
- ii) Φ is K_∞ -finite; i.e., the space spanned by the functions $G(\mathbb{A}) \ni g \mapsto \Phi(g\kappa)$, where κ runs through K_∞ , is finite-dimensional.
- iii) $\Phi(\gamma g) = \Phi(g)$ for all $g \in G(\mathbb{A})$ and $\gamma \in G(F)$.
- iv) There exists a unitary character χ of $Z(\mathbb{A})$ such that $\Phi(zg) = \chi(z)\Phi(g)$ for all $g \in G(\mathbb{A})$ and $z \in Z(\mathbb{A})$.

The usual definition of automorphic forms includes conditions such as slow growth, but for our purposes it is irrelevant whether we include such additional conditions or not. The important fact is that the space of automorphic forms is preserved by right translation R by elements of $G(\mathbb{A}_f)$ and of K_∞ . It is also preserved by the action of \mathfrak{g} , the Lie algebra of $G(\mathbb{A}_\infty)$, given by

$$(R(X)\Phi)(g) = \left. \frac{d}{dt} \right|_0 \Phi(g \exp(tX)).$$

This action extends to an action of $\mathfrak{g}_\mathbb{C}$ and the universal enveloping algebra $\mathcal{U}(\mathfrak{g}_\mathbb{C})$. We call any subspace of the space of automorphic forms invariant under right translation by $G(\mathbb{A}_f)$, K_∞ and \mathfrak{g} , irreducible or not, an *automorphic representation* of $G(\mathbb{A})$ (even though it might not be a representation of $G(\mathbb{A})$ at all). If Φ is an automorphic form, then the automorphic representation generated by Φ is by definition the linear span of all functions obtained by the right action of $G(\mathbb{A}_f)$, \mathfrak{g} and K_∞ on Φ . The automorphic form Φ is called square-integrable if

$$\int_{G(F)Z(\mathbb{A}) \backslash G(\mathbb{A})} |\Phi(g)|^2 dg < \infty.$$

Here, the integration is with respect to a Haar measure on $G(\mathbb{A})$. The space of square-integrable automorphic forms with respect to a fixed unitary character of the center $Z(\mathbb{A})$ is invariant under right translation and carries an obvious inner product.

Theorem 3.1. *For each finite place v , let K_v be an open-compact subgroup of $G(F_v)$; we assume that almost all, but not necessarily all, K_v are maximal compact in G_v . Let Φ be a square-integrable automorphic form on $G(\mathbb{A})$ satisfying the following properties:*

- i) *The (\mathfrak{g}, K_∞) -module spanned by the functions $R(X)R(k)\Phi$, where X runs through $\mathfrak{g}_\mathbb{C}$ and k runs through K_∞ , is irreducible.*
- ii) *For each finite place v , the function Φ is right invariant under K_v and is an eigenvector under the action of the local Hecke algebra $\mathcal{H}(G(F_v), K_v)$.*

Then the automorphic representation of $G(\mathbb{A})$ generated by Φ is irreducible.

Proof. Let V be the automorphic representation generated by Φ . Let F be a non-zero function in V which is right invariant under K_v for all finite v . We can write

$$(5) \quad F = \sum_{i=1}^n R(X_i)R(\kappa_i)R(h_i)\Phi, \quad X_i \in \mathcal{U}(\mathfrak{g}_\mathbb{C}), \kappa_i \in K_\infty, h_i \in G(\mathbb{A}_f).$$

Since Φ is right invariant under almost all local maximal compact subgroups, we may assume that there exists a finite set S of finite places such that each h_i has

trivial components outside of S . For $v \in S$, since F and Φ are both K_v -invariant,

$$F = \frac{1}{\text{vol}(K_v)^2} \sum_{i=1}^n R(X_i)R(\kappa_i) \int_{K_v} \int_{K_v} R(k_1 h_i k_2) \Phi dk_1 dk_2.$$

By Lemma 1.1, the double integration coincides, up to a multiple, with the action of the double coset $K_v(h_i)_v K_v$ on Φ . By hypothesis ii), this action reproduces Φ up to a multiple. We may therefore assume that all of the h_i 's in (5) have trivial v -component. Continuing through all places $v \in S$, we may assume that the elements h_i are not present at all, so that

$$(6) \quad F = \sum_{i=1}^n R(X_i)R(\kappa_i)\Phi, \quad X_i \in \mathcal{U}(\mathfrak{g}_{\mathbb{C}}), \kappa_i \in K_{\infty}.$$

Hence F lies in the $(\mathfrak{g}, K_{\infty})$ -module generated by Φ , which is irreducible by hypothesis i). It follows that

$$(7) \quad \Phi = \sum_{i=1}^m R(Y_i)R(\sigma_i)F, \quad Y_i \in \mathcal{U}(\mathfrak{g}_{\mathbb{C}}), \sigma_i \in K_{\infty}.$$

We proved that whenever F is non-zero and right invariant under all K_v , then a relation of the form (7) holds.

Now let W be an invariant subspace of V . Since V consists entirely of square-integrable functions, there exists an invariant complement W' to W inside V . Write

$$\Phi = F + F', \quad F \in W, F' \in W'.$$

Then, evidently, F and F' are both invariant under K_v for all finite v . By what we proved, Φ can be expressed in the form (7) if F is non-zero. A similar relation holds if F' is non-zero. It follows that Φ is contained in W or in W' . This proves the asserted irreducibility of V . □

It is desirable to have a more practical criterion for the irreducibility expressed in i) of Theorem 3.1. This is provided by the next result. We assume that $G(\mathbb{A}_{\infty})$ is of the form $Z'G'$, where Z' lies in the center of $G(\mathbb{A}_{\infty})$, and where G' is a Lie group whose connected component G'_0 is of finite index in G' . We assume that $K_{\infty} = (Z' \cap K_{\infty})K'$ with a maximal compact subgroup of G' whose connected component K'_0 has finite index in K' . We also assume that $\mathfrak{g}'_{\mathbb{C}}$, the complexification of the Lie algebra \mathfrak{g}' of G' , admits a direct sum decomposition as in (1), with \mathfrak{k} being the Lie algebra of K' . For a K'_0 -type τ , we define

$$(8) \quad L_{\tau} := \mathcal{U}(\mathfrak{g}'_{\mathbb{C}}) \otimes_{\mathcal{U}(\mathfrak{k}_{\mathbb{C}} + \mathfrak{p}_{\mathbb{C}}^{-})} \tau.$$

If (π, V) is a $\mathfrak{g}'_{\mathbb{C}}$ -module and $\sigma \in K'$ is an element, we define the twisted $\mathfrak{g}'_{\mathbb{C}}$ -module (π^{σ}, V) as in (4).

Corollary 3.2. *For each finite place v , let K_v be an open-compact subgroup of $G(F_v)$; we assume that almost all, but not necessarily all, K_v are maximal compact in G_v . Let Φ be a square-integrable automorphic form on $G(\mathbb{A})$ satisfying the following properties:*

- i) *The span of the vectors $\pi(k)\Phi$, where k runs through K'_0 , is an irreducible representation τ of K'_0 .*
- ii) *The K'_0 -type τ occurs in L_{τ} exactly once.*

- iii) For $\sigma \in K'$, $\sigma \notin K'_0$, the $\mathfrak{g}_{\mathbb{C}}$ -modules L_{τ} and $(L_{\tau})^{\sigma}$ have no K'_0 -type in common.
- iv) $R(\mathfrak{p}_{\mathbb{C}}^-)\Phi = 0$.
- v) For each finite place v , the function Φ is right invariant under K_v and is an eigenvector under the action of the local Hecke algebra $\mathcal{H}(G(F_v), K_v)$.

Then the automorphic representation of $G(\mathbb{A})$ generated by Φ is irreducible.

Proof. By Theorem 3.1, it suffices to show that the (\mathfrak{g}, K) -module generated by Φ is irreducible. Since Φ transforms under central elements by a scalar, this is equivalent to saying that the (\mathfrak{g}', K') -module generated by Φ is irreducible. But this is immediate from Proposition 2.4. \square

As an application of Corollary 3.2, consider a Siegel modular cusp form F of degree n with respect to the full modular group. We allow the vector-valued case; i.e., F takes values in an irreducible, holomorphic (polynomial) representation τ of $\mathrm{GL}(n, \mathbb{C})$. As in [AS], we associate with F an adelic function $\Phi : G(\mathbb{A}) \rightarrow \mathbb{C}$, where $G = \mathrm{GSp}(2n)$ and \mathbb{A} is the ring of adèles of \mathbb{Q} . Note that Φ is complex-valued, even if τ is more than one-dimensional; in this case the function Φ is not canonical, but the space of right translates of Φ by elements of $K'_0 \cong U(n)$ is, and is isomorphic to τ .

Corollary 3.3. *Let F be a vector-valued holomorphic Siegel modular cusp form of degree n and full level. Let $\Phi : G(\mathbb{A}) \rightarrow \mathbb{C}$ be an adelic function attached to F , where $G = \mathrm{GSp}(2n)$. If F is an eigenform under all Hecke operators, then Φ generates an irreducible, automorphic representation of $G(\mathbb{A})$.*

Proof. We apply Corollary 3.2 with $G' = \mathrm{Sp}(2n, \mathbb{R})^{\pm}$, the group of all $g \in \mathrm{GSp}(2n, \mathbb{R})$ with multiplier ± 1 . The identity component $G'_0 = \mathrm{Sp}(2n, \mathbb{R})$ has a maximal compact subgroup K'_0 isomorphic to $U(n)$. By our remarks above, condition i) of Corollary 3.2 is satisfied. Considering weights, it is easy to see that conditions ii) and iii) in Corollary 3.2 are satisfied. Condition iv) is equivalent to F being holomorphic; see [AS]. Condition v) is satisfied since F is a Hecke eigenform. \square

Corollary 3.4. *Let F be a vector-valued holomorphic Siegel modular cusp form of degree 2 and full level. Let $\Phi : G(\mathbb{A}) \rightarrow \mathbb{C}$ be an adelic function attached to F , where $G = \mathrm{GSp}(4)$. If F is an eigenform under almost all Hecke operators, then Φ generates an irreducible, automorphic representation of $G(\mathbb{A})$.*

Proof. By the argument in [Ve] (see also Corollary 3.1.7 of [Sc], where the same argument is used), F is automatically an eigenform for *all* Hecke operators. Hence the result follows from Corollary 3.3. \square

REFERENCES

- [AS] Asgari, M., Schmidt, R., *Siegel modular forms and representations*. Manuscripta Math. **104** (2001), 173–200. MR1821182 (2002a:11044)
- [BS] Berndt, R., Schmidt, R., *Elements of the representation theory of the Jacobi group*. Progress in Mathematics, 163. Birkhäuser, Basel, 1998. MR1634977 (99i:11030)
- [Ca] Cartier, P., *Representations of p -adic groups, A survey*. Proc. Symp. Pure Math., 33, Amer. Math. Soc., Providence, RI, 1979, part 1, 111–155. MR546593 (81e:22029)
- [EZ] Eichler, M., Zagier, D., *The theory of Jacobi forms*. Progress in Mathematics, 55. Birkhäuser, Boston, 1985. MR781735 (86j:11043)

- [Ge] Gelbart, S., *Automorphic Forms on Adèle Groups*. Annals of Mathematics Studies, 83. Princeton University Press, 1975. MR0379375 (52:280)
- [Sc] Schmidt, R., *Iwahori-spherical representations of $GSp(4)$ and Siegel modular forms of degree 2 with square-free level*. J. Math. Soc. Japan **57** (2005), 259–293. MR2114732 (2005i:11065)
- [Ve] Veenstra, T., *Siegel modular forms, L -functions, and Satake parameters*. J. Number Theory **87** (2001), 15–30. MR1816034 (2001m:11071)
- [Ya] Yamashita, H., *Multiplicity one theorems for generalized Gelfand-Graev representations of semisimple Lie groups and Whittaker models for the discrete series*. Representations of Lie groups, Kyoto, Hiroshima, 1986, 31–121, Adv. Stud. Pure Math., 14, Academic Press, Boston, 1988. MR1039835 (91e:22023)

DEPARTMENT OF MATHEMATICS, KUMAMOTO UNIVERSITY, KUROKAMI, KUMAMOTO 860-8555,
JAPAN

E-mail address: `narita@sci.kumamoto-u.ac.jp`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OKLAHOMA 73019

E-mail address: `apitale@math.ou.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OKLAHOMA 73019

E-mail address: `rschmidt@math.ou.edu`