Sept. 30, 1999

Measures and Dimensions in Conformal Dynamics

Mariusz Urbański[1]

Department of Mathematics, University of North Texas, Denton, TX 76203-5118, USA
web site: http://www.math.unt.edu/~urbanski
e-mail: urbanski@unt.edu

Abstract: In this survey we collect basic results concerning fractal and ergodic properties of Julia sets of rational functions of the Riemann sphere. We deal with Hausdorff and packing measures and dimensions, we recall and discuss conformal measures and invariant measures absolutely continuous with respect to them. We provide some constructions and proofs characteristic for the surveyed theories. We begin with facts concerning all rational functions. Then we devote lots of time to present the well-developed theory of parabolic maps, and in Section 3 we deal with the class of rational functions whose all critical points contained in the Julia set are non recurrent. This class comprises in particular hyperbolic, parabolic and subhyperbolic maps. Our last section collects some recent results about other subclasses of rational functions, e.g. Collet-Eckmann maps and Fibonacci maps. We attach at the end of this article two appendices which are only loosely related with Sections 1-4 but are close to the heart of the author. These contain a short description of tame mappings and the theory of equilibrium states and Perron-Frobenius operator associated with Hölder continuous potentials.

Contents:

§1. Dimensions of Julia sets.
§2. Parabolic (and hyperbolic) rational functions.
§3. Rational functions whose all critical points contained
in the Julia set are non recurrent.
§4. Other maps.
Appendix 1; Tame functions.
Appendix 2; Equilibrium states and Perron-Frobenius operator.

[1]Research partially supported by NSF Grant DMS 9801583.

§**1. Dimensions of Julia sets.** One of our leading problems will be to determine various fractal dimensions of a Julia set. We mean here first of all Hausdorff dimension, packing dimension and box-counting dimension. We will be also interested in the question of when the Hausdorff measures and the packing measures are positive and finite. Below we provide detailed description of these concepts. Given a subset A of a compact metric space (X, d), a countable family $\{B(x_i, r_i)\}_{i=1}^{\infty}$ of open balls centered at points of A is said to be a packing of A if and only if for any pair $i \neq j$

$$d(x_i, x_j) \geq r_i + r_j .$$

Given a nondecreasing function $g : (0, \varepsilon) \to (0, \infty)$ for some $\varepsilon > 0$, the g-dimensional outer Hausdorff measure $\mathrm{H}_g(A)$ of the set A is defined as

$$\mathrm{H}_g(A) = \sup_{\varepsilon > 0} \inf\Big\{\sum_{i=1}^{\infty} g\big(\mathrm{diam}(A_i)\big)\Big\},$$

where the infimum is taken over all countable covers $\{A_i : i \geq 1\}$ of A by arbitrary sets whose diameters do not exceed ε. If g is of the form x^t instead of writing H_{x^t} we write H_t and speak about t-dimensional outer Hausdorff measure. In this case one will get comparable numbers (in the sense that ratios are bounded away from zero and infinity) if instead of covering A by arbitrary sets one considers only open balls centered at points of A.

The g–dimensional outer packing measure $\Pi_g(A)$ of the set A is defined as

$$\Pi_g(A) = \inf_{\cup A_i = A} \Big\{\sum_i \Pi_g^*(A_i)\Big\}$$

(A_i are arbitrary subsets of A), where Π_g^*, the g-packing premeasure is given by:

$$\Pi_g^*(A) = \inf_{\varepsilon > 0} \sup\Big\{\sum_{i=1}^{\infty} g(2r_i)\Big\}.$$

Here the supremum is taken over all packings $\{B(x_i, r_i)\}_{i=1}^{\infty}$ of the set A by open balls centered at points of A with radii which do not exceed ε. Similarly as in the case of Hausdorff measures if g is of the form x^t instead of writing Π_{x^t} we write Π_t and speak about t-dimensional outer packing measure. These two outer measures H_g and Π_g define countable additive measures on the Borel σ-algebra of X. For additional properties of packing measures and a comprehensive discussion of these and related notions the reader is referred to the paper [TT] and the books [Mat] and [PU1].

The definitions of the Hausdorff dimension $\mathrm{HD}(A)$ of A and packing dimension $\mathrm{PD}(A)$ are the following

$$\mathrm{HD}(A) = \inf\{t : \mathrm{H}_t(A) = 0\} = \sup\{t : \mathrm{H}_t(A) = \infty\}$$

and

$$\mathrm{PD}(A) = \inf\{t : \Pi_t(A) = 0\} = \sup\{t : \Pi_t(A) = \infty\}.$$

Moreover we shall deal with lower box dimension $\underline{\mathrm{BD}}(A)$ and upper box dimension $\overline{\mathrm{BD}}(A)$ which are respectively defined as follows.

$$\underline{\mathrm{BD}}(A) = \liminf_{r\to 0} \frac{\log N(A,r)}{-\log r} \quad \text{and} \quad \overline{\mathrm{BD}}(A) = \limsup_{r\to 0} \frac{\log N(A,r)}{-\log r},$$

where $N(A,r)$ is the minimal number of balls with radii $\leq r$ needed to cover A. If the lower and upper box dimensions coincide, then their common value $\mathrm{BD}(A) = \underline{\mathrm{BD}}(A) = \overline{\mathrm{BD}}(A)$ is called the box dimension of A. In the literature all box dimensions are also called, perhaps more properly, box counting dimensions; we will keep here the shorter name. We recall (see [Fa] for example) that $\mathrm{HD}(A) \leq \mathrm{PD}(A) \leq \overline{\mathrm{BD}}(A)$ and $\mathrm{HD}(A) \leq \underline{\mathrm{BD}}(A) = \underline{\mathrm{BD}}(\overline{A}) \leq \overline{\mathrm{BD}}(A) = \overline{\mathrm{BD}}(\overline{A})$.

Let ν be a Borel probability measure on X and let $t \geq 0$ be a real number. Define the function $\rho = \rho_t(\nu) : X \times (0,\infty) \to (0,\infty)$ by

$$\rho(x,r) = \frac{\nu(B(x,r))}{r^t}.$$

The following two theorems are for our aims the key facts from geometric measure theory. The proof of the first one follows from the results obtained in [RT] and the proof of the second one is contained in [TT].

Theorem 1.3. Assume that X is a compact subspace of a d-dimensional Euclidean space. Then for every $t \geq 0$ there exist constants $h_1(t)$ and $h_2(t)$ with the following properties: If A is a Borel subset of X and $C > 0$ is a constant such that

(1) for all (but countably many) $x \in A$

$$\limsup_{r\to 0} \rho(x,r) \geq C^{-1},$$

then for every Borel subset $E \subset A$ we have $\mathrm{H}_t(E) \leq h_1(t) C \nu(E)$ and, in particular, $\mathrm{H}_t(A) < \infty$.

or

(2) for all $x \in A$

$$\limsup_{r\to 0} \rho(x,r) \leq C^{-1},$$

then for every Borel subset $E \subset A$ we have $\mathrm{H}_t(E) \geq C h_2(t) \nu(E)$.

Theorem 1.4. Assume that X is a compact subspace of a d-dimensional Euclidean space. Then there exist constants $p_1(t)$ and $p_2(t)$ with the following properties: If A is a Borel subset of X and $C > 0$ is a constant such that

(1) for all $x \in A$

$$\liminf_{r\to 0} \rho(x,r) \leq C^{-1},$$

then for every Borel subset $E \subset A$ we have $\Pi_t(E) \geq C p_1(t) \nu(E)$,

or

(2) for all $x \in A$

$$\liminf_{r \to 0} \rho(x, r) \geq C^{-1},$$

then for every Borel subset $E \subset A$ we have $\Pi_t(E) \leq p_2(t) C \nu(E)$ and, consequently, $\Pi_t(A) < \infty$.

(1') If ν is non–atomic then (1) holds under the weaker assumption that the hypothesis of part (1) is satisfied on the complement of a countable set.

Finally to end this short review over some facts from geometric measure theory we recall the following.

Definition 1.5. Given a Borel probability measure μ on a compact metric space X by $\mathrm{HD}(\mu)$, the Hausdorff dimension of measure μ, we understand the number

$$\inf\{\mathrm{HD}(Y) : \mu(Y) = 1\}.$$

Our main tool to understand fractal properties of a Julia set, i.e. its dimensions and measures described above will be the concept of conformal measure introduced in the case of Fuchsian groups by S. J. Patterson in [Pa] and adopted to the case of rational functions by D. Sullivan in [Su1]. Its definition follows below.

From now on let $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ be a rational function of the Riemann sphere $\overline{\mathbb{C}}$. We always assume that the degree of f is at least 2. The main object of our iterest will be the Julia set $J(f)$ and the dynamics of $f|_{J(f)}$. We recall that the Fatou set $\mathcal{F}(f)$ is defined as the set of those points $z \in \overline{\mathbb{C}}$ that admit an open neighbourhood U such that the family of iterates $\{f^n|_U\}_{n \geq 1}$ is normal on the sense of Montel. The Julia set $J(f)$ is then defined as $\overline{\mathbb{C}} \setminus \mathcal{F}$). The following basic properties of Julia sets can be found in [Bea] or [CG] for example.

(1.a) $J(f)$ is a non-empty compact subset of $\overline{\mathbb{C}}$.

(1.b) $J(f)$ is totally invariant, meaning that $f^{-1}(J(f)) = f(J(f)) = J(f)$.

(1.c) Either $J(f) = \overline{\mathbb{C}}$ or $J(f)$ is a nowhere dense subst of $\overline{\mathbb{C}}$.

The property (1.b) allows us to consider the dynamical system $f : J(f) \to J(f)$. The following property completes our list of basic elementary properties of Julia sets $J(f)$.

(1.b) The dynamical system $f : J(f) \to J(f)$ is topologically exact, i.e. for every non-empty open (in the relative topology) set $U \subset J(f)$ there exists an integer $n \geq 0$ such that $f^n(U) = J(f)$.

Definition 1.6. Let $t \geq 0$. A measure m on $J(f)$ is said to be t-conformal for $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$, if $m(J(f)) = 1$ and

$$m(f(A)) = \int_A |f'|^t \, dm$$

for every Borel set $A \subset J(f)$ such that $f|_A$ is injective. A t-conformal measure for some $t \geq 0$ will be called a conformal measure. Since these measures are concentrated on the set $J(f)$, they also will be called conformal for $f : J(f) \to J(f)$ or conformal for f.

In [Su2] Sullivan showed that for every rational mapping f there exists a conformal measure. We briefly sketch his construction below. First, if $J(f) = \overline{\mathbb{C}}$, this measure is the 2-dimensional Lebesgue measure on $\overline{\mathbb{C}}$. Then the exponent will be 2. Otherwise, if $J(f) \neq \overline{\mathbb{C}}$, take a point z in the Fatou set of f, the complement of the Julia set, but not in a Siegel disk nor in a Herman ring. Such a point must exist since f^q restricted to those domains is of degree 1, where q denotes their period. We may also require z to be the center of a disk D such that $2D$ is disjoint from

$$\mathrm{PCV}(f) = \bigcup_{n \geq 1} f^n(Crit(f)).$$

Since for every such a disk D sufficiently small, the sets $f^{-n}(2D)$ are mutually disjoint (this is why we did not want z to be located in a Siegel disk or a Herman ring), it follows from Koebe's distortion theorem that

$$\sum_{n \geq 1} \sum_{x \in f^{-n}(z)} |(f^n)'(x)|^{-2} < +\infty.$$

Let now $\delta \geq 0$ be the infimum of all the exponents $t \geq 0$ for which the series

$$M_t = \sum_{n \geq 1} \sum_{x \in f^{-n}(z)} |(f^n)'(x)|^{-t}$$

converges. Of course $\delta \leq 2$. For every $s > \delta$ let us consider the following atomic probability measures

$$m_s = M_s^{-1} \sum_{n \geq 1} \sum_{x \in f^{-n}(z)} |(f^n)'(x)|^{-s} \delta_x$$

To conclude the construction of conformal measures consider any limit point of measures $\{m_s\}$ if $s \searrow \delta$ in the weak topology of probability measures on the sphere $\overline{\mathbb{C}}$. Denote this limit by m. This is our candidate for a conformal measure and we want to discuss the problem of whether m really fulfills the requirements imposed on conformal measures. First, is m concentrated on the Julia set? One can easily notice that if the series M_δ converges this is not the case. Then simply $m = M_\delta^{-1} \sum_{n \geq 1} \sum_{x \in f^{-n}(z)} |(f^n)'(x)|^{-\delta} \delta_x$ which gives measure 0 to the Julia set. There is however a way of overcoming this difficulty by modifying the the series M_t and the sequence m_s a little bit. Namely, one employs so called slowly varying factors. To be more precise there exists a function $h : (0, \varepsilon) \to (0, +\infty)$ such that $\lim_{t \to 0}(h(at)/h(t)) = 1$ for every $a > 0$ and the series

$$\sum_{n \geq 1} \sum_{x \in f^{-n}(z)} h(|(f^n)'(x)|^{-1}) |(f^n)'(x)|^{-\delta}$$

diverges. We then multiply each factor $|(f^n)'(x)|^{-t}$ appearing in the definition of the series M_t by $h(|(f^n)'(x)|^{-t})$ and each factor $|(f^n)'(x)|^{-s}$ appearing in the definition of the measure m_s by $h(|(f^n)'(x)|^{-s})$. Then, having in mind how the point z was chosen, we conclude that $m(J(f)) = 1$. It can be verified by an explicit computation that the measure m is conformal.

This construction does not tell us too much about the exponent of conformal measure. In order to get such information we proposed in [DU1] a general scheme of constructing generalizations of conformal measures and in [DU2] we have applied it to the case of (Sullivan's) conformal measures. Our approach is a modification of Patterson's and Sullivan's and the main difference is that we start with a point in the Julia set and not in the Fatou set. We will not go into the details here. The interested reader is referred to [DU2] (comp. also [PU1, Chapters 8,10] and [Mc]). Instead, we will describe the precise statements of results from [DU2] and some selected results from Chapters 8-10 of [PU1] that are interesting for us at the moment .

We begin with recalling the notion of entropy of a measure-theoretic dynamical system. So, suppose that $T : X \to X$ is a continuous map of a compact metric space (X, d) and that μ is a Borel T-invariant probability measure on X. T-invariance means that if A is a Borel subset of X then $\mu(f^{-1}(A)) = \mu(A)$. We call μ (or T) ergodic if the only Borel invariant subsets of T (i.e. satisfying $T^{-1}(A) = A$) are either of measure 0 or 1. Given $n \geq 0$ we define the metric d_n on X by setting

$$d_n(x, y) = \max\{d(T^i(x), T^i(y)) : 0 \leq i \leq n\}.$$

Denote by $B_n(x, r)$ the open ball in the metric d_n centered in x and with radius r. If the measure μ is ergodic, then (see [BK]) for μ-a.e. point $x \in X$ the limit

$$\lim_{r\to 0} \lim_{n\to\infty} -\frac{\log \mu(B_n(x, r))}{\log n}$$

exists, is independent of x, and this limit is called the entropy of the system T with respect to the measure μ. This entropy is denoted by $\mathrm{h}_\mu(T)$. Usually a different, more classical approach is undertaken to define the entropy $\mathrm{h}_\mu(T)$ (see [BK], [Wa1], and [HK] for example), the one chosen here is probably the fastest and, at the same time, it reflects in a better way, the nature of entropy.

If μ is a Borel probability measure on the Julia set of a rational function $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$, then $\int \log |f'| d\mu \leq \log ||f'|| < \infty$ (we mean here the derivative calculated with respect to the spherical metric) and, as it was proved in [Pr1], $\int \log |f'| d\mu \geq 0$. We call the integral $\int \log |f'| d\mu$ the Lyapunov (characteristic) exponent of f with respect to the measure μ and we denote it by $\chi_\mu(f)$. The following theorem was proved in [Ma1] (comp. also [Pr2] and Chapters 8-10 of [PU1]).

Theorem 1.7. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a rational function and μ is a Borel probability ergodic invariant measure on $\overline{\mathbb{C}}$ such that $\chi_\mu(f) > 0$, then

$$\mathrm{HD}(\mu) = \frac{\mathrm{h}_\mu(f)}{\chi_\mu(f)}.$$

We would like to point out that ergodicity of f and the fact that the Lyapunov exponent is positive imply that μ is supported on the Julia set. We would also like to add that due to Ruelle's inequality $\mathrm{h}_\mu(f) \leq 2\chi_\mu(f)$, the inequality $\chi_\mu(f) > 0$ is implied by the inequality $\mathrm{h}_\mu(f) > 0$. Following [DU2] we will need the following.

Definition 1.8. The dynamical dimension $\mathrm{DD}(J(f))$ is defined as

$$\mathrm{DD}(J(f)) = \sup\{\mathrm{HD}(\mu)\},$$

where the supremum is taken over all ergodic invariant measure of positive entropy.

We call a compact forward invariant subset $X \subset J(f)$ (i.e. satisfying property $f(X) \subset X$) hyperbolic if there exists $n \geq 1$ such that

$$|(f^n)'(x)| > 1$$

for every $x \in X$ and f^n is topologically conjugate to a subshift of finite type. If only condition $|(f^n)'(x)| > 1$ is satisfied we call the map $f|_X$ expanding. We will also need the following (see [Sh], comp. [PU1]).

Definition 1.9. The hyperbolic dimension $\mathrm{hD}(J(f))$ is defined to be the supremum of Hausdorff dimensions of all hyperbolic subsets of $J(f)$.

and

Definition 1.10. $\delta(f)$ is the minimal exponent for which a conformal measure exists.

We are now in the position to formulate the main result of this section.

Theorem 1.11. For all rational functions $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ it holds

$$\mathrm{DD}(J(f)) = \mathrm{hD}(J(f)) = \delta(f).$$

The equality of the numbers $\mathrm{DD}(J(f))$ and $\delta(f)$ was essentially proved in [DU2]. The proof became complete after F. Przytycki demonstrated in [Pr1] that each compact forward invariant subset of $J(f)$ contains points z for which $\limsup_{n\to\infty} |(f^n)'(z)| = \infty$. The reader may also consult [Mc1] for the proof. The equality $\mathrm{DD}(J(f)) = \mathrm{hD}(J(f))$ has been shown in Chapter 8 of [PU1]. The general scheme of the proof of Theorem 1.11 is the following: First, it follows easily from the classical Bowen' type formula formula (see [Bo1], comp Chapter 6 of [PU1]) that $\mathrm{hD}(J(f)) \leq \mathrm{DD}(J(f))$. A generalization of Bowen's formula will be discussed in Section 2 which is devoted to parabolic and hyperbolic rational functions. Second, using Pesin's theory type and Katok's type considerations (see Chapter [8] of [PU1]) one shows that $\mathrm{hD}(J(f)) \geq \mathrm{DD}(J(f))$. The easiest part (modulo Pesin's and Katok-Strelcyn type results (see Chapter [8] of [PU1])) is to show that $\mathrm{hD}(J(f)) \leq \delta(f)$.

One uses here the fact that each ergodic invariant measure of positive entropy gives rise to a subset of the set of conical points of full measure. The set of conical points will be discussed in greater detail soon. Finally, developing the technique of so called compact invariant $K(V)$-sets and defining on them classical topological pressure of the functions $-t\log|f'|$, $t\in\mathbb{R}$, one constructs a conformal measure with exponent $\leq \mathrm{DD}(J(f))$. This proves the inequality $\delta(f)\leq \mathrm{DD}(J(f))$ and finishes the proof of Theorem 1.11.

A natural byproduct of the proof of Theorem 1.11 is the appearance of the concept of conical points. They correspond to conical points in the theory of Kleinian groups and were introduced in [U1] (comp. also [DMNU] and [Mc1]). Here is their definition.

Definition 1.12. We call a point $z\in J(f)$ a conical point of f if there exist $\theta>0$ and an infinite increasing sequence $n_k\geq 1$ of positive integers such that for each k there exists $f_z^{-n_k}$, a holomorphic inverse branch of f^{n_k}, which is defined on the disk $B(f^{n_k}(z),\theta)$ and sends the point $f^{n_k}(z)$ to z. The set of all conical points of f will be denoted by $J_c(f)$.

In [DMNU] we have proved the following.

Theorem 1.13. There exists at most one value of t for which a t-conformal measure exists and is supported on the set of conical points of f. Additionally, for such a t there is exactly one t-conformal measure supported on the set of conical points of f.

We would like to end up the discussion concentrated around Theorem 1.11 with two problems, some partial answers to these will be given in the following sections.

Problem 1.14. Is the Hausdorff dimension of the Julia set always equal to its dynamical dimension?

Problem 1.15. Does there always exist a conformal measure supported on the set of conical points?

We shall end this section with an estimate from below of the Hausdorff dimension of the Julia set which is caused by the existence of so called parabolic points. First we need the following.

Definition 1.16. A periodic point z of f, say of period $q\geq 1$, is called parabolic if the derivative $(f^q)'(z)$ is a root of unity.

In the literature parabolic points are frequently called rationally indifferent or neutral periodic points. A Fatou's theorem says that (see [Bea] and [CG] for ex.) that the set Ω of all parabolic points is finite and contained in the Julia set. Suppose now that $\omega\in\Omega$ is a fixed point of f. Looking at the Taylor series expansion of f around ω and at Fatou's flower theorem (see [Bea] and [CG] for ex.) one can deduce (see [ADU]) that there exists $\sigma>0$ such that for every sufficiently small $\delta>0$ and every point $z\in B(\omega,\delta)\setminus\{\omega\}$ all continuous inverse branches $f_\omega^{-n}: B(z,\sigma|z-\omega|)\to\mathbb{C}$, $n\geq 1$, of f^n moving point z towards

ω are well defined. Looking again at the Taylor series expansion of f around ω one can conclude (see [ADU]) that for every $x \in B(z, \sigma|z-\omega|)$

$$|(f_\omega^{-n})'(x)| \asymp n^{-(p+1)/p} \tag{1.1}$$

and

$$|f_\omega^{-n}(x) - \omega| \asymp n^{-1/p}, \tag{1.2}$$

where $p = p(\omega)$ is the number of petals associated with the parabolic point ω and where the comparability constants depend only on the distance of z to ω. Relying on (1.1) and following the proof of Theorem 8.5 in [ADU] (see also [Mc1]) we shall provide a simple argument for the following.

Theorem 1.17. If a rational function $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ has a parabolic point with p petals than $\mathrm{HD}(J(f)) > p/(p+1)$.

Proof. Let ω be a parabolic point of f. Passing to a sufficiently high iterate we may assume that f fixes ω. Consider now a point $z \in J(f) \cap B(\omega, \delta) \setminus \{\omega\})$ and let $V \subset B(z, \sigma|z-\omega|)$ be a neighbourhood of z which is chosen so small that all the inverse images $f_\omega^{-n}(V)$, $n \geq 0$, are mutually disjoint. Let m be an arbitrary conformal measure for $f : J(f) \to J(f)$ and let $h = \mathrm{HD}(J(f))$. Since $z \in J(f)$, $m(V) > 0$. Using (1.1) and Koebe's distortion theorem we obtain the following

$$1 \geq m\left(\bigcup_{n\geq 1} f_\omega^{-n}(V)\right) = \sum_{n\geq 1} m(f_\omega^{-n}(V)) \asymp \sum_{n\geq 1} n^{\frac{-(p+1)h}{p}} m(V).$$

Thus $h > p/(p+1)$ and the proof is complete. ■

We would like to conclude this section with the remark that McMullen's paper [Mc3] contains an algorithm for computing the Hausdorff dimension of any conformal hyperbolic set up to any desired accuracy. Also, in this paper lots of various examples are treated.

§**2. Parabolic (and hyperbolic) rational functions.** We call a continuous map $T : X \to X$ from a compact metric space (X, d) into itself expansive if there exists $\delta > 0$ such that for all pairs of points $x, y \in X$, $x \neq y$, there exists $n \geq 0$ such that $d_n(x, y) \geq \delta$. In the sequel, any number δ satisfying this property will be called an expansive constant. In [DU3] we have proved the following topological characterization of expansive rational functions $f : J(f) \to J(f)$.

Theorem 2.1. A rational function $f : J(f) \to J(f)$ is expansive if and only if the Julia set $J(f)$ contains no critical points of f.

One direction of this equivalence is simple. Namely, if the Julia set $J(f)$ contains a critical point, say c, then $f : J(f) \to J(f)$ is not expansive. To see this, notice that by perfectness of $J(f)$ we can find a sequence x_n, $n \geq 1$, of points in $J(f)$ converging to c. Since c is a critical point of f, for each point x_n (sufficiently close to c) there exists a different point $y_n \neq x_n$ such that

$$f(y_n) = f(x_n). \tag{2.1}$$

Moreover we may require that $\lim_{n\to\infty} y_n = c$ and, in particular, that $\lim_{n\to\infty} |y_n - x_n| = 0$. This together with (2.1) contradict expansiveness. The other direction is more subtle and requires a good control of the inverse branches.

We call a rational function $f : J(f) \to J(f)$ hyperbolic if there exists $n \geq 1$ such that $\inf\{|(f^n)'(z)| : z \in J(f)\} > 1$. Of course, each hyperbolic function is expansive. The following topological characterization of hyperbolicity is known (see [Bea] and [CG] for ex. and the references therein).

Theorem 2.2. A rational function $f : J(f) \to J(f)$ is hyperbolic if and only if

$$\overline{\mathrm{PCV}} \cap J(f) = \emptyset.$$

It follows from Theorems 2.1 and 2.2 that a rational function is expansive but not hyperbolic if and only if the Julia set contains no critical points of f but intersect the ω-limit set of critical points. Now, if we have a rational function whose Julia set $J(f)$ contains no critical points of f, then it follows from the classification theorem of connected components of Fatou's set (the complement of the Julia set) and from the fact that the boundaries of Siegel disks and Herman rings intersect the ω-limit set of critical points (see [Bea] and [CG] for ex.) that if f is not hyperbolic, then there must be a parabolic point in the Julia set. On the other hand, the existence of such a point obviously rules out hyperbolicity. Thus, we obtain the following.

Theorem 2.3. A rational function $f : J(f) \to J(f)$ is expansive but not hyperbolic if and only if the Julia set $J(f)$ contains no critical points of f but contains at least one parabolic point.

In order to say more about the geometry and dynamics of expansive rational functions we need the concept of topological pressure (see [Ru], [Wa], [Bo2] and [PU1] for ex.). In order to introduce it, we choose one of the fastest methods. For alternative approaches better suited to derive various properties of topological pressure see also the items quoted above. So, consider a continuous mapping $T : X \to X$ of a compact metric space (X, d) and a continuous function $\phi : X \to \mathbb{R}$, called, following physical tradition, potential. Given $n \geq 0$ and $\varepsilon > 0$, we say that a subset F of X is (n, ε)-separated if it is separated with respect to the metric d_n, which means that if x and y are two distinct points of X, then $d_n(x, y) \geq \varepsilon$. Fixing now $\varepsilon > 0$ we consider an arbitrary sequence $F_n(\varepsilon)$, $n \geq 1$, of maximal (in the sense of inclusion) (n, ε)-separated sets. We then define the topological pressure of

the function ϕ with respect to the mapping T as follows

$$\mathrm{P}(T,\phi)=\lim_{\varepsilon\to 0}\limsup_{n\to\infty}\frac{1}{n}\log\left(\sum_{x\in F_n(\varepsilon)}\exp\sum_{j=0}^{n-1}\phi\circ T^j(x)\right).$$

Observe that in the case of expansive maps it is not necessary to take the limit for $\varepsilon\to 0$; it is then enough to take ε as an expansive constant of T. In case when $\phi=0$, the topological pressure is called topological entropy. It has been proved (see [Gr] and [Ly2]) that topological entropy of any rational function is equal to the logarithm of its degree. Topological pressure belongs to topological dynamics, whereas metric entropy is a notion in ergodic theory. The link joining them is given by the following formula called variational principle (see [Ru], [Wa], [Bo2] and [PU1] for ex.).

$$\mathrm{P}(T,\phi)=\sup\{\mathrm{h}_\mu(T)+\int\phi d\mu\},$$

where the supremum is taken over all Borel probability T-invariant (ergodic) measures of X.

Since the Julia set of an expansive rational function $f: J(f)\to J(f)$ contains no critical points, we may define for a given $t\in\mathbb{R}$ the function

$$\mathrm{P}(t)=\mathrm{P}(f,-t\log|f'|),$$

which is called the pressure function and which is well-defined. The basic properties of this function are collected in the following.

Theorem 2.4. Suppose that $f: J(f)\to J(f)$ is expansive. Then

(a) The pressure function is convex and therefore continuous.

(b) The pressure function is non-increasing.

(c) If the mapping f is hyperbolic, then the pressure function is strictly decreasing and $\lim_{t\to+\infty}\mathrm{P}(t)=-\infty$.

(d) If the mapping f is parabolic then there exists a number $s(f)>0$ such $\mathrm{P}(t)=0$ for all $t\geq s(f)$ and $\mathrm{P}|_{[0,s(f)]}$ is strictly decreasing.

The part (a) of this theorem is classical, the part concerning parabolic maps ((b) and (d)) has been proved in [DU3]. The proof makes an extensive use of the variational principle and the non-negativeness of Lyapunov exponents. The fact that in the parabolic case for all t, $P\big(t\big)\geq 0$ is caused by the existence of invariant probability measures supported on (periodic) orbits of parabolic points. In case of hyperbolic maps we denote the only zero of $\mathrm{P}(t)$ also by $s(f)$. The following theorem, which is classical in the hyperbolic case and called Bowen-Manning-McCluskey formula (see [Bo1], comp. [MC]), has been proved in the parabolic case in [DU3]. It can be considered as a strengthening of Theorem 1.11 and partial answer to the Problem 1.14.

Theorem 2.5. If a rational function $f : J(f) \to J(f)$ is expansive, then we have the following equalities

$$\mathrm{DD}(J(f)) = \mathrm{hD}(J(f)) = \delta(f) = s(f) = \mathrm{HD}(J(f)).$$

Using Theorem 1.7 and the variational principle, it is not difficult to see that $s(f) \leq \mathrm{DD}(J(f))$. So, a priori, the largest number in Theorem 2.5 is $\mathrm{HD}(J(f))$, the Hausdorff dimension of the Julia set $J(f)$. We would like to provide the sketch of the proof that $\mathrm{HD}(J(f)) \leq \delta(f)$. In order to do this and later on to be able to discuss the so called uniform formula for conformal measure, we need the concept of hyperbolic zoom which appeared in [U2], which was extensively used in author's papers with M. Denker and which got its name in [SU1]. We start with the remark that in the expansive case the ω-limit set of critical points is contained in forward orbits of parabolic points and attracting periodic points. Therefore, given $e > 0$, an expansive constant of $f : J(f) \to J(f)$, there exists a positive constant $\theta > 0$ such that for every $z \in J(f) \setminus B(\Omega, e)$ the ball $B(z, \theta)$ is disjoint from PCV. Consequently all continuous inverse branches of all iterates f^n, $n \geq 1$, are well-defined on all the balls $B(z, \theta)$, where $z \in J(f) \setminus B(\Omega, e)$. It now follows from expansiveness of $f : J(f) \to J(f)$ that if $x \in J(f) \setminus \bigcup_{n \geq 0} f^{-n}(\Omega)$, then there exists an infinite increasing sequence $n_j(x)$, $j \geq 1$, such that for every $j \geq 1$ we have

$$f^{n_j(x)}(x) \notin B(\Omega, e).$$

In the sequel, we let $\{n_j(x)\}$ denote the maximal sequence satisfying the above property. Consequently we get the following

Theorem 2.6. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a parabolic rational function, then the set of conical points $J_c(f)$ coincides with the set $J(f) \setminus \bigcup_{n \geq 0} f^{-n}(\Omega)$.

The sequence of numbers

$$r_j(x) = |(f^{n_j(x)})'(x)|^{-1} \tag{2.2}$$

is called the hyperbolic zoom at the point x. Applying Koebe's distortion theorem and conformality property, it is easy to observe that if m is a t-conformal measure for f, then there exists a constant $B(m) \geq 1$ such that for all $x \in J_c(f)$

$$B(m)^{-1} \leq \frac{m(B(x, r_j(x)))}{r_j(x)^t} \leq B(m). \tag{2.3}$$

We are now in position to provide the proof taken from [U2] (comp. [DU3]) of the following.

Theorem 2.7. If H_t is the t-dimensional Hausdorff measure on $J(f)$ and m is a t-conformal measure for $f : J(f) \to J(f)$, then H_t is absolutely continuous with respect to m such that the Radon-Nikodym derivative is bounded from above. Consequently, $t \geq \mathrm{HD}(J(f))$ and there is no t-conformal measure for $t < \mathrm{HD}(J(f))$.

Proof. Let $F \subset J(f)$ be any Borel set. Put $E = J_c(f) \cap F = F \setminus \bigcup_{n=0}^{\infty} f^{-n}(\Omega)$. Since the set $\bigcup_{n=0}^{\infty} T^{-n}(\Omega)$ is at most countable, $\mathrm{H}_t(E) = \mathrm{H}_t(F)$. Fix $\gamma, \epsilon > 0$. Since m is regular we can find a countable cover $\{B(x_i, r(x_i))\}_{i=1}^{\infty}$ of E of multiplicity bounded by a universal constant $C \geq 1$ (coming from the Besicovic covering theorem) such that $x_i \in E$, $0 < r(x_i) < \gamma$ is of the form $r_j(x_i)$ for every $i = 1, 2, \ldots$ (defined in (2.2)) and such that

$$m(\bigcup_{i=1}^{\infty} B(x_i, r(x_i)) \setminus E) < \epsilon$$

Hence, applying (2.3) to the measure m, we obtain

$$\sum_{i=1}^{\infty} r(x_i)^t \leq B(m) \sum_{i=1}^{\infty} m(B(x_i, r(x_i))) \leq B(m) C m(\bigcup_{i=1}^{\infty} B(x_i, r(x_i))) \leq CB(m)(\epsilon + m(E))$$

Letting $\varepsilon \searrow 0$ and then $\gamma \searrow 0$, we get $\mathrm{H}_t(F) = \mathrm{H}_t(E) \leq CB(m)m(E) \leq CB(m)m(F)$. The proof is complete. ■

As already announced, it follows immediately from this theorem that $\mathrm{HD}(J(f)) \leq \delta(f)$. Thus the proof of Theorem 2.5 is complete. ■

Theorem 2.5 implies in particular that there are no t-conformal measures with exponents $t < \mathrm{HD}(J(f))$. The following theorem describes the structure of all conformal measures.

Theorem 2.8. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a parabolic map, then the following 3 statements hold:

(a) There is no t-conformal measure with $t < \mathrm{HD}(J(f))$.

(b) There exists exactly one $\mathrm{HD}(J(f))$-conformal measure. This measure is atomless.

(c) For every $t > \mathrm{HD}(J(f))$ there are t-conformal measures. Each such a measure is a convex combination of purely atomic measure supported on $\bigcup_{n \geq 0} f^{-n}(\Omega)$.

As was already mentioned, item (a) of this theorem is already contained in Theorem 2.5. The existence of $\mathrm{HD}(J(f))$-conformal measures is also given by Theorem 2.5. The existence of an atomless $\mathrm{HD}(J(f))$-conformal measure is proved in Theorem 8.7 of [ADU] and its uniqueness is demonstrated in Theorem 4.6 of [DU4]. Finally, item (c) is proved in Theorem 13 of [DU3].

Recently (see Section 3 and especially Section 5) there has been very strong interest in the Hausdorff dimension of various Julia sets. Particularly popular is the problem of whether the Hausdorff dimension of a Julia set is equal to 2 or less than 2. Up to our knowledge the first result in this direction which goes beyond the hyperbolic case has been obtained in [ADU], where the following theorem has been proved.

Theorem 2.9. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a parabolic map, then $\mathrm{HD}(J(f)) < 2$.

Below we provide the proof of this theorem. We give this prove since the method employed there has turned out to be applicable also in other settings.

Proof of Theorem 2.9. First, using the concept of hyperbolic zooms, the distortion property on the balls $B(x, r_j(x))$ and Lebesgue's density theorem, one demonstrates as in [Ly1] and [DH1] (cf. also [DU3]) that the (2-dimensional) Lebesgue measure of the Julia set $J(f)$ is equal to 0. We now follow the proof of Theorem 8.8 in [ADU]. Suppose on the contrary that $\mathrm{HD}(J(f)) = 2$ and let m be a 2-conformal measure existing by Theorem 2.5. For every $x \in J_c(f)$ let $\{r_j(x)\}_{j=1}^{\infty}$ be the hyperbolic zoom at the point x introduced by (2.2). Fix $\varepsilon > 0$ and denote by λ the (2-dimensional) Lebesgue measure on $\mathbb{C}$. Since $\lambda(J(f)) = 0$ and since $\lim_{j\to\infty} r_j(x) = 0$, for every $x \in J_c$ there exists a radius $r(x)$ being of the form $r_j(x)$ such that

$$\lambda\Big(\bigcup_{x\in J_c} B(x, r(x))\Big) < \varepsilon.$$

Now by the Besicovic covering theorem (see [Gu], comp. [PU1]) we can choose a countable subcover $\{B(x_i, r(x_i))\}_{i=1}^{\infty}$ of the cover $\{B(x, r(x))\}_{x\in J_c(f)}$ of $J_c(f)$, of multiplicity bounded by some universal constant $C \geq 1$. Therefore, using (2.3) with $t = 2$, we obtain

$$\begin{aligned} m(J_c) &\leq \sum_{i=1}^{\infty} m(B(x_i, r(x_i))) \leq B(m) \sum_{i=1}^{\infty} r(x_i)^2 = B\pi^{-1} \sum_{i=1}^{\infty} \lambda(B(x_i, r(x_i))) \\ &\leq B\pi^{-1} C \lambda\Big(\bigcup_{i=1}^{\infty} B(x_i, r(x_i))\Big) \leq BC\pi^{-1}\varepsilon. \end{aligned}$$

Thus, letting $\varepsilon \to 0$, we obtain $m(J_c) = 0$. Since, by Theorem 2.8(b), m is atomless, it implies that $m(J(f)) = 0$. This contradiction finishes the proof. ■

There is another nice way, not using conformal measures, to show that $\mathrm{HD}(J(f)) < 2$ for parabolic maps, which shows even more, that the upper box-counting dimension of these Julia sets is less than 2. Namely, a bounded subset X of a Euclidean space is said to be porous if there exists a positive constant $c > 0$ such that each open ball centered at a point of X and of an arbitrary radius $0 < r \leq 1$ contains an open ball of radius cr disjoint from X. The notion of porosity has appeared in several contexts and for a short survey and some bibliographical references the reader may see the paper [YR]. By a simple counting argument it is straightforward to prove the following.

Theorem 2.10. If X is a porous subset of R^d, then $\overline{\mathrm{BD}}(X) < d$.

Since L. Geyer has proved in [Ge] that

Theorem 2.11. The Julia set of a parabolic rational function is porous.

these two latter theorems imply

Theorem 2.12. The upper box-counting dimension of the Julia set of a parabolic rational function is less than 2.

Notice however that in [DU5], using the jump transformation (see below), we have proved the following.

Theorem 2.13. If f is a parabolic mapping, then $\overline{\mathrm{BD}}(J(f)) = \mathrm{HD}(J(f))$.

In particular combining this theorem and Theorem 2.9, we get Theorem 2.12.

For parabolic maps our interests went deeper to study finer structure of the Julia sets. We were interested in which cases Hausdorff and packing measures are finite and positive. The complete answer is given by the the following theorem proven in [DU6].

Theorem 2.14. If f is a parabolic mapping and $h := \mathrm{HD}(J(f))$, then

(a) $\mathrm{H}_h(J(f)) < \infty$ and $\Pi_h(J(f)) > 0$.
(b) $\mathrm{H}_h(J(f)) = 0$ if and only if $h < 1$.
(c) $\Pi_h(J(T)) = \infty$ if and only if $h > 1$.

The proof of this theorem makes an extensive use of the h-conformal measure m. The fact $\mathrm{H}_h(J(f)) < \infty$ follows immediately from the first part of Theorem 2.7; the fact $\Pi_h(J(f)) > 0$ can be proved similarly. The results $\mathrm{H}_h(J(f)) = 0$ if $h < 1$ and $\Pi_h(J(T)) = \infty$ if $h > 1$ are not too difficult and their proofs use closer and closer visits of forward iterates of points from J_c to Ω. The hardest parts are to prove that $\mathrm{H}_h(J(f)) > 0$ if $h \geq 1$ and $\Pi_h(J(f)) < \infty$ if $h \leq 1$. To see this we need to have a closer look at the ratios $\frac{m(B(x,r))}{r^h}$, where m is the unique h-conformal measure. The first rather implicit investigations of these ratios have been made in [DU6], more explicit and extensive in [DU5] and eventually in [SU1] they took on the final form resulting in the following.

Theorem 2.15. Let f be a parabolic rational map with Julia set $J(T)$ of Hausdorff dimension h. Let m denote the associated h-conformal measure supported on $J(T)$. Then there exists a function $\phi : J(f) \times \mathbb{R}^+ \to \mathbb{R}^+$ such that for each $z \in J(f)$ and for every positive $r < \mathrm{diam}(J(f))$ we have that

$$m(B(z,r)) \asymp r^h \cdot \phi(z,r).$$

The function ϕ is called the conformal fluctuation function. For $r < \mathrm{diam}(J(f))$, the values of ϕ are determined as follows. For $z \in J_c(f)$, assume that r relates to the hyperbolic zoom at z such that $r_{j+1}(z) \leq r < r_j(z)$ and such that $f^k(z) \in B(\omega, e)$, for all $k \in (n_j(z), n_{j+1}(z)]$ and for some $\omega \in \Omega$. In this situation we have that

$$\phi(z,r) \asymp \begin{cases} \left(\frac{r}{r_j(z)}\right)^{(h-1)p(\omega)} & \text{for } r \geq r_j(z)\left(\frac{r_{j+1}(z)}{r_j(z)}\right)^{1/(1+p(\omega))} \\ \left(\frac{r_{j+1}(z)}{r}\right)^{(h-1)} & \text{for } r \leq r_j(z)\left(\frac{r_{j+1}(z)}{r_j(z)}\right)^{1/(1+p(\omega))} \end{cases}$$

If $z \in J_p$, then for every $r > 0$ small enough

$$\phi(z, r) \asymp r^{(h-1)p(\omega)}.$$

Let

$$p_{\max} = \max\{p(\omega) : \omega \in \Omega\}$$

A finer analysis of the conformal fluctuation function ϕ done in [SU1] leads to the following theorem proven in [SU1] as Corollary 4.4.

Theorem 2.16. For the conformal fluctuation function ϕ of the h-conformal measure m associated to a parabolic rational map f the following holds.

(a) If $h = 1$, then for all $z \in J_c(f)$ and all $0 < r < \operatorname{diam}(J(T))$

$$\phi(z, r) \asymp 1.$$

(b) If $h < 1$, then for m-almost every $z \in J(f)$

$$\limsup_{r \to 0} \frac{\log \phi(z, r)}{\log \log \frac{1}{r}} = \frac{(1-h)p_{\max}}{h + (h-1)p_{\max}}.$$

(c) If $h > 1$, then for m-almost every $z \in J(f)$

$$\liminf_{r \to 0} \frac{\log \phi(z, r)}{\log \log \frac{1}{r}} = \frac{(1-h)p_{\max}}{h + (h-1)p_{\max}}.$$

The main ingredient in the proof of this result was so called Khintchin Limit Law for parabolic rational maps proven in [SU1] (Theorem 4.3) with the help of the jump transformation and invariant measures absolutely continuous with respect to conformal measure which will be discussed more extensively later in this section. This Khintchin law reads as follows.

Theorem 2.17. (The Khintchin Limit Law for parabolic rational maps) The hyperbolic zoom at m-almost every $z \in J(f)$ has the property that

$$\limsup_{j \to \infty} \frac{\log \left(r_j(z) \,/\, r_{j+1}(\xi) \right)}{\log \log \frac{1}{r_j(z)}} = \frac{1 + p_{\max}}{h + (h-1)p_{\max}}.$$

Note that this Khintchin law is the analog of the logarimic law of geodesics for limit sets of Kleinian groups (see [SV]). Given $\rho \in \mathbb{R}$ let

$$\psi_\rho(r) := r^h \left(\log \frac{1}{r} \right)^{(1+\rho)(1-h)p_{\max}/(h+(h-1)p_{\max})}$$

be a gauge function and let H_{ψ_ρ} and Π_{ψ_ρ} be the corresponding Hausdorff and packing measure. As an immediate consequence of our finer analysis of the conformal fluctuation function (Theorem 2.16) we get the following result proven in [SU1] (Corollary 4.5).

Theorem 2.18. Let f be a parabolic rational map with Julia set $J(f)$ of Hausdorff dimension h and with h-conformal measure m. We then have

(a) If $\rho > 0$ and $h < 1$, then $m << \mathrm{H}_{\psi_\rho}$ and $\mathrm{H}_{\psi_\rho}(J(f)) = \infty$

(b) If $\rho > 0$ and $h > 1$, then there exists a Borel set E_ρ such that $m(E_\rho) = 1$ and $\Pi_{\psi_\rho}(E_\rho) = 0$

(c) If $\rho \leq 0$ and $h < 1$, then then there exists a Borel set F_ρ such that $m(F_\rho) = 1$ and $\mathrm{H}_{\psi_\rho}(F_\rho) = 0$

(d) If $\rho \leq 0$ and $h > 1$, then $m << \Pi_{\psi_\rho}$ and $\Pi_{\psi_\rho}(J(f)) = \infty$.

We would like to remark that for $h < 1$ similar results were obtained in [DR].

Another direction of exploring finer fractal structure of parabolic Julia sets is motivated by multifractal analysis and Diophantine approximation (see [Bes] and [Ja]) of irrational numbers and the limit sets of Kleinian groups (see an extended discussion of this motivation in [SU2]). In order to define briefly the setting let us consider an expansive rational function $f : J(f) \to J(f)$ and a Hölder potential $g : J(f) \to \mathbb{R}$ satisfying $g \geq \log |f'|$. Fix $x \in J(f)$. Following [HV] we then define the set

$$\mathcal{J}_{g,x}(f) = \Big\{ z \in J(f) : z \in B\big(y, \exp\big(-\sum_{j=0}^{n-1} g \circ f^j(y)\big)\big) \text{ for infinitely many pairs } (y,n) \text{ such that } f^n(y) = x \Big\}.$$

In [HV] Hill and Velani have proved the following.

Theorem 2.19. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a hyperbolic rational function, then for every $x \in J(f)$, $\mathrm{HD}(\mathcal{J}_{g,x}(f)) = s(g)$, where $s(g)$ is the only zero of the pressure function $t \mapsto \mathrm{P}(-tg)$.

It turns out that in parabolic case the situation is much more complicated. In [SU2] we have considered the function $g = (1+\sigma)\log|f'|$, $(\sigma \geq 0)$ and we have proved the following, where $\mathcal{J}_x^\sigma(f) = \mathcal{J}_{(1+\sigma)\log|f'|,x}(f)$.

Theorem 2.20. Suppose that $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a parabolic rational function. Fix an element $x \in f^{-1}(\Omega) \setminus \Omega$. Then

(a) If $h < 1$, then $\mathrm{HD}(\mathcal{J}_x^\sigma(f)) = \frac{h}{1+\sigma}$.

(b) If $h \geq 1$, then

$$\mathrm{HD}(\mathcal{J}_x^\sigma(f)) = \begin{cases} \frac{h}{1+\sigma} & \text{for } \sigma \geq h-1 \\ \frac{h+\sigma p(\omega)}{1+\sigma(1+p(\omega))} & \text{for } \sigma < h-1. \end{cases}$$

Given $\omega \in \Omega$, $y \in f^{-n}(f^{-1}(\omega) \setminus \{\omega\})$ for some $n \geq 1$ and $\alpha > 0$ we define $r(y) = |(f^n)'(y)|^{-1}$ and $r_{y,\alpha} = \alpha^{1/(1+p(\omega))} r(y)^{p(\omega)/(1+p(\omega))}$. Our approach heavily relied on the following Dirichlet cover type result and developed the methods worked out originally to tackle similar problems in the theory of Kleinian groups (see for ex. [St1] and [St2]).

Theorem 2.21. Suppose that $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a parabolic rational function. Fix an element $\omega \in \Omega$. Then there exist universal constants $\kappa_c, \kappa_p, \alpha_0 > 0$, depending only on f, such that for each $\omega \in \Omega$ and for each $0 < \alpha < \alpha_0$ the following holds.

(a) The family $\{B(y, \kappa_p r_{y,\alpha}) : r(y) \geq \alpha\}$ provides a packing of $J(f)$.
(b) The family $\{B(y, \kappa_c r_{y,\alpha}) : r(y) \geq \alpha\}$ provides a covering of $J(f)$.

We would like to draw reader's attention especially to the case $h > 1$ and $\sigma < h - 1$ in Theorem 2.20, since it shows that the situation in parabolic setting is much more complex than in the hyperbolic case and the most natural extension of Theorem 2.19 (requiring now $s(g)$ to be the least zero of the pressure function $\mathrm{P}(-tg)$) to the parabolic case fails since, due to Theorem 2.5 $s\big((1+\sigma)\log|f'|\big) = \frac{h}{1+\sigma}$. In order to formulate our results in terms multifractal analysis let us introduce the following notation. Let ν denote a Borel probability measure on $\mathbb{R}^n$. For $\theta > 0$, we define the following sets.

$$\mathcal{I}^{\theta}(\nu) := \{z \in \mathrm{supp}(\nu) : \liminf_{r\to 0} \frac{\log \nu(B(z,r))}{\log r} \leq \theta\},$$

$$\mathcal{I}_{\theta}(\nu) := \{z \in \mathrm{supp}(\nu) : \liminf_{r\to 0} \frac{\log \nu(B(z,r))}{\log r} \geq \theta\},$$

$$\mathcal{S}^{\theta}(\nu) := \{z \in \mathrm{supp}(\nu) : \limsup_{r\to 0} \frac{\log \nu(B(z,r))}{\log r} \leq \theta\},$$

$$\mathcal{S}_{\theta}(\nu) := \{z \in \mathrm{supp}(\nu) : \limsup_{r\to 0} \frac{\log \nu(B(z,r))}{\log r} \geq \theta\}.$$

The collections of Hausdorff dimensions of these sets, for $\theta > 0$, are referred to as the weak singularity spectra of ν (see [St2]). As a consequence of Theorem 2.20 we have proved in [SU2] the following.

Theorem 2.22. The weak singularity spectra of the h-conformal measure m of a parabolic rational map with Julia set of Hausdorff dimension h are determined by the following, where, let us recall, $p_{mx} := \max_{\omega\in\Omega} p(\omega)$.

(a) If $h = 1$, then the weak singularity spectra of m are trivial. Namely, in this case we have for all $z \in J(T)$ that

$$\lim_{r\to 0} \frac{\log m(B(z,r))}{\log r} = h.$$

(b) If $h < 1$, then

$$\mathrm{HD}(\mathcal{I}^{\theta}(m)) = \begin{cases} 0 & \text{for } 0 < \theta \leq h + (h-1)p_{mx} \\ \frac{h(\theta-(h+(h-1)p_{mx}))}{(1-h)p_{mx}} & \text{for } h + (h-1)p_{mx} < \theta < h \\ h & \text{for } \theta \geq h \end{cases}.$$

(c) If $h > 1$, then

$$\mathrm{HD}(\mathcal{S}_{\theta}(m)) =$$

$$\begin{cases} h & \text{for } 0 < \theta \leq h \\ \frac{(h-1)(h+(h-1)p_{mx})}{(\theta-1)p_{mx}} - \frac{h-p_{mx}}{p_{mx}} & \text{for } h < \theta < \frac{h(h+(h-1)p_{mx})-(h-1)p_{mx}}{h} \\ \frac{h(h+(h-1)p_{mx}-\theta)}{(h-1)p_{mx}} & \text{for } \frac{h(h+(h-1)p_{mx})-(h-1)p_{mx}}{h} \leq \theta < h + (h-1)p_{mx} \\ 0 & \text{for } \theta \geq h + (h-1)p_{mx}. \end{cases}$$

$$\mathrm{HD}(\mathcal{I}^{\theta}(m)) = \begin{cases} 0 & \text{for } 0 < \theta < h \\ h & \text{for } \theta \geq h. \end{cases}$$

(d) In both cases $h < 1$ and $h > 1$, we have that

$$\mathrm{HD}(\mathcal{I}_{\theta}(m)) = \begin{cases} h & \text{for } 0 < \theta \leq h \\ 0 & \text{for } \theta > h. \end{cases}$$

$$\mathrm{HD}(\mathcal{S}^{\theta}(m)) = \begin{cases} 0 & \text{for } 0 < \theta < h \\ h & \text{for } \theta \geq h. \end{cases}$$

We would like to devote the last part of this section to discuss ergodic theory of h-conformal measure m. This theory has been developed in [DU4] and [ADU]. Recall (see [Ru, 7.29, p.146]) that a family $\Re = \{R_1, \ldots, R_s\}$ is said to be a Markov partition for a continuous map $T : X \to X$ of a compact metric space X if the following conditions are satisfied: ($\mathrm{Int}(A)$ will denote the interior of the set A)

$$\begin{aligned} &R_i = \overline{\mathrm{Int}(R_i)} \qquad \text{for every } R_i \in \Re. \\ &\text{If } i \neq j \text{ then } \mathrm{Int}(R_i) \cap \mathrm{Int}(R_j) = \emptyset. \\ &\text{Each set } T(R_i) \text{ is a union of sets } R_j \in \Re. \end{aligned}$$

We proved in Theorem 2.2 of [DU4] that each continuous expansive mapping of a compact metric space admits a compatible with topology metric with respect to which this mapping is expanding. Since each open continuous expanding map admits Markov partitions of arbitrarily small diameters (see Theorem 2.4 and Lemma 2.3 from [DU4], comp. [Ru] and [Bo2]) (with boundaries of measure zero for a fixed atomless probability measure), we obtain the following

Theorem 2.23. If $f : J(f) \to J(f)$ is a parabolic rational map and if μ is an atomless probability measure on $J(f)$, then $f : J(f) \to J(f)$ admits Markov partitions $\Re = \{R_1, \ldots, R_s\}$ of arbitrarily small diameters such that $\mu(\partial R_1 \cup \ldots \cup \partial R_s) = 0$ (∂A denotes the boundary of the set A).

Having a Markov partition $\Re$ with a sufficiently small diameter we define the associated jump transformation as follows. We denote by $R(\Omega)$ the union of all elements of $\Re$ that have non-empty intersection with Ω and given $z \in J(f) \setminus \Omega$ we define $n(z) \geq 0$ to be the least integer $n \geq 0$ such that $f^n(z) \notin R(\Omega)$. We then define the jump transformation $f^* : J(f) \setminus \Omega \to J(f)$ (depending on the Markov partition) by setting

$$f^*(z) = f^{n(z)+1}(z).$$

The idea of the jump transformation is taken from [Sc1] (cf. also [Sc2]). Using this mapping and adapting the Schweiger formalism developed in [Sc1] to our context, we proved in [DU4] the following two theorems.

Theorem 2.24. If $f : J(f) \to J(f)$ is a parabolic rational map and m is the only h-conformal measure, then for every jump transformation f^* there exists a unique probability f^*-invariant measure μ^* absolutely continuous with respect to m. Moreover μ^* is ergodic and equivalent with m.

Theorem 2.25. If $f : J(f) \to J(f)$ is a parabolic rational map and m is the only h-conformal measure, then there exists a unique up to a multiplicative constant σ-finite f-invariant measure μ absolutely continuous with respect to m. Moreover μ is ergodic, conservative and equivalent with m.

We developed the concept of Schweiger's formalism further in [ADU] under the name of Markov fibered systems. We also heavily relied on Theorem 2.24 and additional properties of the jump transformation in [SU1].

Definition 2.26. Suppose that μ is a σ-finite measure on a metric space X. We call $x \in X$ a point of infinite condensation of μ if and only if $\mu(U) = \infty$ for every open neighbourhood of x. Otherwise we call x a point of finite condensation.

In [DU4] we prove as Theorem 4.9 the following.

Theorem 2.27. If $f : J(f) \to J(f)$ is a parabolic rational map, then the set of points of infinite condensation of invariant measure μ produced in Theorem 2.25 is contained in Ω.

In [ADU] we have dealt with the problem of when the σ-finite invariant measure μ is finite. We gave there in Theorem 9.8 the following complete answer to this question.

Theorem 2.28. If $f : J(f) \to J(f)$ is a parabolic rational map, then the σ-finite invariant measure μ is finite if and only if

$$h = \mathrm{HD}(J(f)) > 2 \max \left\{ \frac{p(\omega)}{p(\omega)+1} \right\}.$$

More precisely, we have the following.

Theorem 2.29. If $f : J(f) \to J(f)$ is a parabolic rational map and $\omega \in \Omega$, then μ is of infinite condensation at ω if and only if $h \leq \frac{2p(\omega)}{p(\omega)+1}$.

As an immediate consequence of Theorem 2.28 we get the following.

Corollary 2.30. If $h = \mathrm{HD}(J(f)) \leq 1$, then the measure μ is infinite.

Corollary 2.31. If $h > 1$ and $p(\omega) = 1$ for every $\omega \in \Omega$, then μ is finite.

Observe that if $B : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a Blaschke product, then B is parabolic and $J(B) \subset J(f)$. So, as an immediate consequence of Corollary 2.30 we get the following.

Corollary 2.32. If $B : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a Blaschke product, then the σ-finite invariant measure μ is infinite.

Since it is proven (see [U2] and [Zd]) that the Hausdorff dimension of the Julia set of the quadratic polynomial $z \mapsto z^2 + 1/4$ is greater than 1, as an immediate consequence of Corollary 2.31 we get the following.

Corollary 2.33. The σ-finite invariant measure μ for the quadratic polynomial $z \mapsto z^2 + 1/4$ is finite.

Definition 2.34. Suppose that $T : X \to X$ is a continuous mapping of a compact metric space and that $g : X \to \mathbb{R}$ is a continuous potential. A Borel probability T-invariant measure μ on X is called an equilibrium state of g (and T) if $\mathrm{P}(T, g) = \mathrm{h}_\mu(T) + \int g d\mu$ (See also the variational principle discussed in the beginning of this section).

It turns out that Theorem 2.28 can be completed as follows.

Theorem 2.35. If $f : J(f) \to J(f)$ is a parabolic rational map, then the σ-finite invariant measure μ is finite if and only if one of the following two condition holds.

(a) The potential $g = -h \log |f'|$ has at least one atomless equilibrium state.

(b) The topological pressure function $t \mapsto \mathrm{P}(t)$ is not differentiable at the point $t = h$.

We would like to conclude this section with the remark that recently parabolic bifurcations have been extensively investigated and lots of interesting dimension results have been obtained (see for ex. [BZ], [Mc1], [Sh], [DSZ], [HS], [Zi1] and [Zi2]).

§3. Rational functions whose all critical points contained in the Julia set are non recurrent.

We recall that if $T : X \to X$ is a continuous map of a topological space X, then for every point $x \in X$, the ω-limit set of x denoted by $\omega(x)$ is defined to be the set of all limit points of the sequence $\{T^n(x)\}_{n \geq 0}$. We call a point x recurrent if $x \in \omega(x)$; otherwise x is called non-recurrent.

Definition 3.1. A rational function $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is called an NCP map if all critical points contained in the Julia set $J(f)$ are non-recurrent.

The class of NCP maps obviously contains all expanding and parabolic maps. It also comprises the important class of so called subexpanding maps which are defined by the requirement that $f|_{\omega(\mathrm{Crit}(f)) \cap J(f)}$ is expanding and the class of geometrically finite maps defined by the property that the forward trajectory of each critical point contained in the Julia set is finite and disjoint from Ω. The class of subexpanding maps was explored in [DU7] from the point of view of the fractal properties of the Julia set. We have dealt with NCP maps in [U3] and [U4] exploring their fractal and ergodic properties. Their

geometric properties have been studied in [CJY] and the rigidity type results have been obtained in [PU2]. In [Mc1] the reader may find some properties of geometrically finite maps proven earlier in [U3] for NCP maps. The paper [Ha] contains some ergodic properties of geometrically finite maps.

We start our description of fractal and ergodic properties of NCP maps with the following characterization of conical points obtained in [U3] as Proposition 6.1.

Theorem 3.2. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is an NCP map, then

$$J_c(f) = \overline{\mathbb{C}} \setminus \bigcup_{n\geq 0} f^{-n}\big(\Omega(f) \cup \mathrm{Crit}(J(f))\big).$$

As a sort of complementary result to this theorem we would like to state the following rather straightforward consequence of topological exactness of all rational functions restricted to their Julia sets which was stated in [U3] (Lemma 5.2).

Theorem 3.3. The set $\omega(\mathrm{Crit}(J(f)))$ is nowhere dense in $J(f)$.

Let, as before, $h = \mathrm{HD}(J(f))$ be the Hausdorff dimension of the Julia set $J(f)$. We proved in [U3] that there exists exactly one h-conformal measure m and that this measure is atomless. As Lemma 7.3 we have proved there the following.

Theorem 3.4. The conformal measure m is supported on the set of transitive points, i.e. those points whose ω-limit set is the whole Julia set.

In Proposition 7.4 of [U3] it is shown that Theorem 2.7 remains true for NCP maps. This, the existence of an h-conformal measure, and Theorem 11 immediately imply the following partial answer to the Problem 1.14.

Theorem 3.5. If $f : J(f) \to J(f)$ is an NCP rational function, then we have the following equalities

$$\mathrm{DD}(J(f)) = \mathrm{hD}(J(f)) = \delta(f) = s(f) = \mathrm{HD}(J(f)).$$

The complete characterization of geometric measures on the Julia sets of NCP maps is provided by the following theorem proven in [U3] as Theorem 1.1.

Theorem 3.6. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is an NCP rational function, then

(a) $\mathrm{H}_h(J(f)) < \infty$ and $\Pi_h(J(f)) > 0$.
(b) $\mathrm{H}_h(J(f)) = 0$ if and only if $h < 1$ and $\Omega(f) \neq \emptyset$.
(c) $\Pi_h(J(f)) = \infty$ if and only if $h > 1$ and $\Omega(f) \neq \emptyset$.

Since it is straightforward to check that both Hausdorff and packing measures appearing in the above theorem are conformal when finite and positive, it follows from this theorem and uniqueness of h-conformal measure that the conformal measure has always a geometrical

characterization as a normalized Hausdorff or packing measure. Dealing still with geometry let me add the following two results.

Theorem 3.7. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is an NCP rational function, then either $J(f) = \overline{\mathbb{C}}$ or $\mathrm{HD}(J(f)) < 2$.

Theorem 3.8. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is an NCP rational function such that $J(f) \neq \overline{\mathbb{C}}$, then the Julia set $J(f)$ is porous and consequently the upper box dimension $\overline{\mathrm{BD}}(J(f)) < 2$.

The first of these two theorems has been stated in [U3] and the second one is proved in [PU3]. Let me add that in [U4] we have provided a number of sufficient conditions for $\mathrm{HD}(J(f)) = \overline{\mathrm{BD}}(J(f))$. The proof of Theorem 3.7 relieses on the following three results which form a basic toolbox for the class of NCP maps and Collet-Eckmann type maps (see Definition 4.6).

Theorem 3.9. There exist $\theta > 0$ and $M \geq 0$ such that for every $x \in J(f) \setminus B(\Omega, \delta)$ every integer $n \geq 0$, every connected component V of $f^{-n}(B(x, \theta)$ is simply connected and the restriction $f^n|_V$ has at most M critical points (counted with multiplicities).

Theorem 3.10. For each $\varepsilon > 0$ and $D < \infty$ there are constants C_1 and C_2 such that the following holds for all rational maps $F : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$, all $x \in \overline{\mathbb{C}}$, all $1/2 \leq r < 1$ and all $0 < \gamma \leq 1/2$:
Assume that V (resp. V') is a simply connected component of $F^{-1}(B(x, \gamma))$ (resp. $F^{-1}(B(x, \frac{1}{2}\gamma))$ with $V \supset V'$. Assume further that $\overline{\mathbb{C}} \setminus V$ has diameter at least ε and F has at most D critical points (counted with multiplicity) in V. Then

(a) $$|F'(y)|\mathrm{diam}(V') \leq C_1(1-r)^{-C_2}\delta$$

for all $y \in V'$. Furthermore, if $r = 1/2$ and $0 < \tau < 1/2$, let $B'' = B(z, \tau\gamma)$ be any disk contained in $B(x, \gamma/2)$ and let V'' be a component of $F^{-1}(B'')$ contained in V'. Then

(b) $$\mathrm{diam}(V'') \leq C_3 \mathrm{diam}(V')$$

with $C_3 = C_3(\tau, \varepsilon, D)$ and $\lim_{\tau \to 0} C_3(\tau, \varepsilon, D) = 0$, and

(c) V'' contains a disk of radius greater than or equal to $C_4\mathrm{diam}(V')$

around every preimage of $F^{-1}(z)$ that is contained in V''. Here $C_4 = C_4(\tau, \varepsilon, D) > 0$.

Theorem 3.11. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a rational function and if $\Omega \neq \emptyset$ then for every $\rho > 0$ there exists $c = c(\rho) > 0$ such that for each $x \in J(f)$ and each r such that $\rho\mathrm{dist}(x, \Omega) \leq r \leq 1$ there exists an open ball $B \subset B(z, r) \setminus J(f)$ with radius cr.

Theorem 3.9 follows from [Ma2, Theorem II]. See also [CJY, Theorem 2.1] or [U3, Lemma 2.12 and Lemma 5.1] where this theorem or its variants have been proved and substantially

used. A crucial step in the proof of Theorem 3.9 is that given an arbitrary $\varepsilon > 0$ there exists θ so that the diameters of all the above components are less than ε. Theorem 3.10 is a part of the "bounded distortion" lemma that has been proved in [Pr3] and [PR1, Lemma 2.1]. Finally Theorem 3.11 is a straightforward consequence of Fatou's flower theorem.

We would like to start now a discussion of ergodic theory of NCP maps. It will be based on [U4] and [Ha]. The starting point is the following result proved in [U4] as Theorem 4.2

Theorem 3.12. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is an NCP rational function, then up to a multiplicative constant there exists exactly one f-invariant σ-finite measure μ absolutely continuous with respect to h-conformal measure m. Moreover μ is equivalent with m, conservative and ergodic.

The idea of the proof of Theorem 3.12 is to apply a general sufficient condition for the existence of σ-finite absolutely continuous invariant measure proven in [Mar]. In order to formulate this condition suppose that X is a σ-compact metric space, m is a Borel probability measure on X, positive on open sets, and that a measurable map $T : X \to X$ is given with respect to which measure m is quasi-invariant, i.e. $m \circ T^{-1} << m$. Moreover we assume the existence of a countable partition $\alpha = \{A_n : n \geq 0\}$ of subsets of X which are all σ-compact and of positive measure m. We also assume that $m(X \setminus \bigcup_{n\geq 0} A_n) = 0$, and if additionally for all $m, n \geq 1$ there exists $k \geq 0$ such that

$$m(T^{-k}(A_m) \cap A_n) > 0,$$

then the partition α is called irreducible. Martens' result comprising Proposition 2.7 and Theorem 2.10 of [Mar] reads the following.

Theorem 3.13. Suppose that $\alpha = \{A_n : n \geq 0\}$ is an irreducible partition for $T : X \to X$. Suppose that T is conservative and ergodic with respect to the measure m. If for every $n \geq 1$ there exists $K_n \geq 1$ such that for all $k \geq 0$ and all Borel subsets A of A_n

$$K_n^{-1} \frac{m(A)}{m(A_n)} \leq \frac{m(T^{-k}(A))}{m(T^{-k}(A_n))} \leq K_n \frac{m(A)}{m(A_n)}, \tag{3.1}$$

then T has a σ-finite T-invariant measure μ absolutely continuous with respect to m. Additionally μ is equivalent with m, conservative and ergodic, and unique up to a multiplicative constant.

Proof of Theorem 3.12. Let us first sketch the proof of Theorem 3.13 given in [Mar] and then, as its consequence, a short proof of Theorem 3.12 taken from [U4]. So, one fixes an element $A \in \alpha = \{A_n : n \geq 0\}$ and then one considers the following sequences of measures

$$S_k(m) = \sum_{i=0}^{k-1} m \circ f^{-i} \quad \text{and} \quad Q_k(m) = \frac{S_k(m)}{S_k(m)(A)}.$$

It is proven in [Mar] that each weak limit μ of the sequence $Q_k(m)$ (a sequence $\{\nu_k : k \geq 1\}$ of measures on X is said to converge weakly if for all $n \geq 1$ the measures ν_k converge weakly on all compact subsets of A_n) is a required invariant measure and

$$\mu(F) = \lim_{n\to\infty} Q_k(m)(F)$$

for every Borel set $F \subset X$.

In order to complete the proof of Theorem 3.11 we only need to construct an irreducible partition α with property (3.1). Indeed, set $Y = J(f)\setminus(\mathrm{PCV}\cup\Omega)$. For every $y \in Y$ consider a ball $B(y, r(y))$ such that $r(y) > 0$, $m(\partial B(y, r(y))) = 0$, and $r(y) < (1/2)\mathrm{dist}(y, \mathrm{PCV}\cup\Omega)$. The balls $B(y, r(y))$, $y \in Y$, cover Y and since Y is a metric separable space, one can choose a countable cover, say $\{\tilde{A}_n : n \geq 0\}$, from them. We may additionally require that the family $\{\tilde{A}_n : n \geq 0\}$ is locally finite that is that each point $x \in Y$ has an open neighborhood intersecting only finitely many balls $\tilde{A}_n$, $n \geq 0$. We now define the family $\alpha = \{A_n : n \geq 0\}$ inductively setting

$$A_0 = \tilde{A}_0 \quad \text{and} \quad A_{n+1} = \tilde{A}_{n+1} \setminus \bigcup_{k=1}^{n} \overline{\tilde{A}_n}$$

(and throwing away empty sets). Obviously α is a disjoint family and

$$\bigcup_{n\geq 1} A_n \supset J(f) \setminus (\mathrm{PCV} \cup \Omega) \setminus \bigcup_{n\geq 0} \partial\tilde{A}_n$$

whence in view of [U3, Corollary 7.2], $m\big(\bigcup_{n\geq 0} A_n\big) = 1$. The distortion condition (3.1) follows now from Koebe's distortion theorem with all constants $K_n = K$, the constant corresponding to the scale $1/2$ in this theorem and irreducibility of partition α follows from openess of the sets A_n and topological exactness of the map $f : J(f) \to J(f)$.

The following results describing the structure of invariant measure μ have been proved in [U4].

Theorem 3.14. If $\Omega = \emptyset$, then there exists an f-invariant probability measure μ equivalent to m.

Theorem 3.15. If $\omega(\mathrm{Crit}(f)) \cap \Omega = \emptyset$, then the set of points of infinite condensation of μ is contained in Ω.

Theorem 3.16. If $h = \mathrm{HD}(J(f)) \geq 1$, then the set of points of infinite condensation of μ is contained in the set of parabolic points.

Theorem 3.17. If $\omega \in \Omega \setminus \mathrm{PCV}$, then μ is of infinite condensation at ω if and only if $h \leq \frac{2p(\omega)}{p(\omega)+1}$.

Theorem 3.18. If $\omega \in \Omega$ and $h \leq \frac{2p(\omega)}{p(\omega)+1}$, then μ has infinite condensation at ω.

Theorem 3.19. If $c \in J(f)$ is a critical point of f of order q, $\omega = f(c) \in \Omega$, and $h \leq \frac{2qp(\omega)}{p(\omega)+1}$, then ω is of infinite condensation of measure μ.

We would like to conclude this section with the result which is a combined consequence of Theorem 3.15 and Theorem 3.17 and which has been later proved as Theorem 1.2 in [Ha].

Theorem 3.20. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is geometrically finite then the invariant measure μ is finite if and only if $h > 2 \max\{\frac{p(\omega)}{p(\omega)+1} : \omega \in \Omega\}$.

§**4. Other maps.** The famous result of Bishop and Jones (see [BJ]) saying that the Hausdorff dimension of the limit set of a finitely generated but not geometrically finite Kleinian group is 2 has stimulated an extensive activity aiming to explain the situation for rational functions of the Riemann sphere $\overline{\mathbb{C}}$. It has been known (see Theorem 2.9 and Theorem 3.7) that the Hausdorff dimension of Julia sets ($\neq \overline{\mathbb{C}}$) of parabolic and NCP maps is less than 2. To the other pole of the spectrum belonged the following result of Shishikura [Sh] (see also [HS], [Zi], and [Mc1]).

Theorem 4.1. $\sup\{\mathrm{HD}(J(f_c))\} = 2$, where the supremum is taken over all parameters c such that the quadratic map $f_c(z) = z^2 + c$ is parabolic. The set of parameters c such that $\mathrm{HD}(J(f_c)) = 2$ is dense in the boundary of the Mandelbrot set.

The first result in the opposite direction, going beyond NCP maps, was due to F. Przytycki who proved in [Pr4] the following.

Theorem 4.2. There exists a set of positive Lebesgue measure parameters $c \in \mathbb{R}$ for which $\mathrm{HD}(J(f_c)) < 2$.

Since then a number of similar results have been obtained. Let us list some of them below.

Theorem 4.3. ([Mc2]) If $f(z) = \mathrm{e}^{2\pi i\theta} z + z^2$, where θ is a badly approximable irrational number (of bounded type) (so, f has a Siegel disk) then the Julia set $J(f)$ is porous and consequently $\mathrm{HD}(J(f)) \leq \overline{\mathrm{BD}}(J(f)) < 2$.

A quadratic map $f_c(z) = z^2 + c$ is a Fibonacci map if, loosely speaking, the closest returns of the critical point 0 occur precisely at the moments given by Fibonacci sequence $S_n = S_{n-1} + S_{n-2}$ and $S_0 = S_1 = 2$. Graczyk and Smirnov proved in [GS1] the following.

Theorem 4.4. If $f_c : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a real ($c \in \mathbb{R}$) quadratic Fibonacci map, then $\mathrm{HD}(J(f_c)) < 2$.

A simply connected domain $A \subset \overline{\mathbb{C}}$ is called a Hölder domain (with exponent $\alpha \in (0, 1]$) if the Riemann mapping $R : D^1 \to A$ ($D^1 = \{z \in \mathbb{C} : |z| < 1\}$) can be extended in a Hölder

continuous fashion (with exponent α) to the closed disk $\overline{D^1}$. P. Jones and N. Makarov proved in [JM] the following.

Theorem 4.5. There exists a constant $c > 0$ such that for every $\alpha \in (0, 1]$ and every Hölder domain with exponent α, $\mathrm{HD}(\partial A) \leq 2 - c\alpha$.

Definition 4.6. A rational function $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is called Collet-Eckmann if there are constants $C > 0$ and $\lambda > 1$ such that for every critical point in $J(f)$ whose forward orbit does not contain any other critical point, and every $n \geq 1$

$$|(f^n)'(f(c))| \geq C\lambda^n. \tag{4.1}$$

Graczyk and Smirnov proved in [GS2] the following.

Theorem 4.7. The basin of attraction to ∞ of any Collet-Eckmann polynomial is a Hölder domain. In particular, due to Theorem 4.5 the Hausdorff dimension of its Julia set is strictly less than 2.

In [PR] and [PU3] some extensions of the last part of this theorem have been proved. In fact a weaker form of porosity has been established there sufficient to conclude that the upper box counting dimension is less than 2. Namely

Definition 4.8. A set $X \subset \mathbb{R}^d$, $d \geq 1$, is called mean porous if there are constant $p_1 < \infty$ and $p_2 > 0$ such that for each $z \in E$ there exist an increasing sequence n_j of positive integers and a sequence of points $z_j \in \mathbb{R}^d$ with $||z - z_j|| \leq 2^{-n_j}$ such that $n_j < p_1 j$ and and $\mathrm{dist}(z_j, X) > p_2 2^{-n_j}$.

In [KR] the following extension of Theorem 2.10 has been proved.

Theorem 4.9. If X is a mean porous subset of a Euclidean space R^d, then $\overline{\mathrm{BD}}(X) < d$.

The main result of [PR] reads as follows.

Theorem 4.10. The Julia set of a Collet-Eckmann rational function with no parabolic points is mean porous. In particular, due to Theorem 4.9, its upper box-counting dimension is strictly less than 2.

In [PU3] an appropriate definition of Collet-Eckmann rational functions with parabolic points is provided (loosely speaking one requires (4.1) to take form $|(f^n)'(f(c))| \geq C\lambda^{\phi(n)}$ for an appropriately changed time $\phi(n) \leq n$) and the following theorem has been proved.

Theorem 4.11. The Julia set of a Collet-Eckmann rational function with parabolic points is mean porous. In particular, due to Theorem 4.9, its upper box-counting dimension is strictly less than 2.

Another direction of recent activity has been stimulated by Problem 1.14 (see also Theorem 1.11). In [Pr3] F. Przytycki has proved the following.

Theorem 4.12. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a Collet-Eckmann map (in fact it suffices here to assume that $|(f^n)'(f(c))| \geq Ce^{\theta\sqrt{n}}$), then $\underline{\mathrm{DD}}(J(f)) = \mathrm{HD}(J(f))$. If additionally f has no parabolic points, then $\mathrm{HD}(J(f)) = \overline{\mathrm{BD}}(J(f))$.

and

Theorem 4.12. If $f(z) = z^2 + c$ is a non-renormalizable quadratic polynomial with c out of the main cardioid in the Mandelbrot set, then $DD(J(f)) = \mathrm{HD}(J(f))$.

We would like to conclude this section with the remark that in the papers [Pr3], [Pr4], [PR] and [PU3] so called Rule II (see [DPU]) has been used extensively. Its formulation and the context surrounding it the reader may find in Appendix 2.

Appendix 1; Tame functions. Unless stated otherwise all the results formulated in this appendix are taken from [PU2]. As in the previous sections let $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ be a rational function of the Riemann sphere $\overline{\mathbb{C}}$. By $T(f)$ we denote the set of all points in $J(f)$ for which

$$\limsup_{n\to\infty} \mathrm{dist}(f^n(z), \mathrm{PCV}(f)) > 0.$$

Definition A1.1. We say that a rational function $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is tame if $\mathrm{PCV}(f) \cap J(f)$ is nowhere dense in $J(f)$ and there exists a t-conformal measure m (sometimes to be more specific denoted by m_f) such that $m(T(f)) = 1$.

Remark A1.2. The set $T(f)$ was already considered in [GPS] (called there the transverse limit set). In fact its origin goes back to [Ly].

Remark A1.3. Notice that if the singular set $\mathrm{PCV}(f)$ of an arbitrary rational function f is nowhere dense in $J(f)$, then $T(f)$ contains the set of all transitive points of f (i.e. points with dense forward orbit). Therefore for such a function to be tame it is sufficient to have a conformal measure m supported on the set of transitive points.

Clearly $T(f)$ is a subset of the set of conical points of f and therefore as an immediate consequence of Theorem 1.13 we get the following.

Theorem A1.4. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is tame, then there exists at most one value t for which a t-conformal measure exists and is supported on $T(f)$. Additionally, for such a t there exists exactly one t-conformal measure supported on $T(f)$ and this measure is ergodic.

Thus the measure $m = m_f$ and the exponent $t = t_f$ are determined uniquely. It is easy to see that m_f is non-atomic. We can say something more about the exponent $t = t_f$. Indeed,

with considerations similar to those in the proof of Theorem 2.7, we can demonstrate the following.

Theorem A1.5. If a rational function $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is tame, then the t_f-dimensional Hausdorff measure of $T(f)$ is finite. In particular $t_f \geq \mathrm{HD}(T(f))$ (HD abbreviates Hausdorff dimension).

Employing the method introduced by M. Martens in [Mar] and proceeding in the same way as in the proof of Theorem 4.2 in [U2] (see Theorem 3.11 and the discussion surrounding it), we can demonstrate the following.

Theorem A1.6. If a rational function $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is tame, then up to a multiplicative constant there exists exactly one f-invariant σ-finite measure $\mu = \mu_f$ absolutely continuous with respect to m. Moreover, μ is equivalent with m, conservative and ergodic.

Examples of tame functions are provided for instance by NCP maps. Indeed, Theorem 3.2 and Theorem 3.4 of this paper along with the paragraph preceding it, imply the following.

Theorem A1.7. Each NCP rational function f is tame and $t_f = \mathrm{HD}(J(f))$. Moreover, the t_f-conformal measure is supported on the set of transitive points.

The main result of the paper [PU2] is the following.

Theorem A1.8. Suppose that f and g are two tame rational maps. Let h be an invertible map from a full measure m_f subset of $J(f)$ onto a full measure m_g subset of $J(g)$, preserving the algebras of measurable sets for m_f and m_g and conjugating f to g, namely $h \circ f = g \circ h$. Then the following conditions (1)-(6) are equivalent.

(1) h extends to a Möbius conjugacy between $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ and $g : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$.
(2) h extends to a conformal homeomorphism conjugating f and g on neighbourhoods of $J(f)$ and $J(g)$ in $\overline{\mathbb{C}}$.
(3) h extends to a real-analytic diffeomorphism conjugating f and g on neighbourhoods of $J(f)$ and $J(g)$ in $\overline{\mathbb{C}}$.
(4) h extends to a homeomorphism from $J(f)$ to $J(g)$ such that h and h^{-1} are Lipschitz continuous.
(5) h extends to a homeomorphism from $J(f)$ to $J(g)$ such that for every periodic point x of f, say of period p, $|(f^p)'(x)| = |(g^p)'(h(x))|$.
(6) The measure class of m_f is transported under h to the measure class of m_g.

Here for the implication (6) $\Rightarrow$ (2), f and g are assumed not to be critically finite with parabolic orbifold (see [Th, Ch.13], [DH2, §9] or [Zd] for the definition) for which $J(f) = J(g) = \overline{\mathbb{C}}$. We call such maps exceptional.

Remark A1.9. The implication (5) $\Rightarrow$ (2) was in fact proved by E. Prado in [Pra] for all rational maps. The only missing point, the non-linearity (see [PU2] for the definition) caused by parabolic points turns out to be easy. Prado's proof of the only hard part (5) $\Rightarrow$ (2) was done by approximating $J(f)$ by forward invariant expanding repellers inside $J(f)$, where h extends conformally, see [Su3], [Pr5]. Our proof of the implication (5) $\Rightarrow$ (2) is different and goes via (6). This requires the assumptions that f is tame. Thus the main new result in our paper is the implication (6) $\Rightarrow$ (2) which extends Sullivan's result for non-linear repellers, see [Su3] and [Pr5]); we use the same scheme of proof. This implication is called rigidity. Note that the implication (2) $\Rightarrow$ (1) is straightforward. This holds even in a general, not just tame, situation. Any conformal conjugacy on neighbourhoods of Julia sets for any two rational functions extends to a conformal conjugacy (Möbius map) to the whole sphere. In particular by Theorem A1.9 the measure theoretic conjugacy class of a tame map coincides with its conformal conjugacy class. In other words one cannot perturb a tame map inside the measure theoretic conjugacy class changing its conformal conjugacy class. Equivalence classes of measures classify tame maps.

A rational function $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is said to be Julia real-analytic if its Julia set is contained in a finite union of pairwise disjoint real-analytic curves which will be denoted by $\Gamma = \Gamma_f$. Frequently in such a context one alternatively speaks about real analyticity of the Julia set $J(f)$. We conclude this section with two technical ingredients interesting themselves which have been proved in [PU2] as Corollary 2.4 and Proposition 2.3 respectively; the first being an immediate consequence of the second.

Theorem A1.10. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is tame, then the Jacobian of the map $f : J(f) \to J(f)$ with respect to the measure μ has a real-analytic extension on a neighbourhood of $J(f) \setminus \mathrm{PCV}$ in $\mathbb{C}$. If the map f is Julia real-analytic, then the Jacobian has a real-analytic extension on a neighbourhood of $J(f) \setminus \mathrm{PCV}$ in Γ.

Theorem A1.11. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a tame mapping, then the Radon-Nikodym derivative $\rho = d\mu/dm$ has a real-analytic real-valued extension on a neighbourhood of $J(f) \setminus \mathrm{PCV}$ in $\mathbb{C}$. If f is Julia real-analytic, then ρ has a real-analytic extension on a neighbourhood of $J(f) \setminus \mathrm{PCV}$ in Γ.

Appendix 2; Equilibrium states and Perron-Frobenius operator. In this appendix $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a rational function of the Riemann sphere $\overline{\mathbb{C}}$ (no additional assumptions imposed) and $\phi : J(f) \to \mathbb{R}$ is a Hölder continuous function satisfying the following condition

$$\mathrm{P}(\phi) = \mathrm{P}(\phi, f) > \sup(\phi), \tag{A2.1}$$

where, let us recall, $\mathrm{P}(\phi)$ is topological pressure defined between Theorem 2.3 and Theorem 2.4. We start with the following general result of Lyubich (see [Ly2]).

Theorem A2.1. If $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a rational function then the function $\mu \mapsto \mathrm{h}_\mu(f)$ is upper semi-continuous and consequently each continuous potential has an equilibrium state.

In fact in [Ly2] M. Lyubich proved more that each rational function is asymptotically h-expansive, the notion introduced by M. Misiurewicz in [Mi] in connection with his investigations of continuous maps of an interval. Recall from [DU1] that a normalized Borel measure ν on a Borel measure space $(X, \mathcal{F})$ is called g-conformal for the continuous mapping $T : X \to X$, where $g : X \to \mathbb{R}$ is a positive continuous function, if $\nu(T(A)) = \int_A g \, d\nu$ for every $A \in \mathcal{F}$ such that $T|A$ is invertible. The paper [DU1] contains some basic facts about this notion and, in particular, sufficient conditions for the existence of these measures. The central result of [DU8] is the following.

Theorem A2.2. Let $f : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ be a rational function and let $\phi : \overline{\mathbb{C}} \to \mathbb{R}$ be a Hölder continuous function satisfying (A2.1). Then:

(a) There exists exactly one $\exp(c - \phi)$-conformal probability measure m and necessarily $c = P(\phi)$.
(b) There exists exactly one equilibrium state μ for ϕ and f. The measures μ and m are equivalent and $\rho = \frac{d\mu}{dm}$ is a positive continuous function.
(c) The dynamical system $(J(f), f, \mu)$ is metrically exact and, consequently, its natural extension is a K-system.

Let us add that later in [Pr6] F. Przytycki proved the same results and that in the special case of measure of maximal entropy ($\phi = 0$) some partial results have been obtained in [FLM], [Ly2], and [Ma3]. In order to formulate the results concerning Perron-Frobenius operator (to be defined later) let us recall that a bounded linear operator $A : X \to X$ on a complex Banach space X is called almost periodic, if for every $x \in X$ the family $\{A^n(x)\}_{n=1}^{\infty}$ is relatively compact in the strong topology. The basic property of almost periodic operators proven in [Ly2] is the following.

Theorem A2.3. If a bounded linear operator $A : X \to X$ on a complex Banach space X is almost periodic then the Banach space X decomposes into a direct sum $X_u + X_0$ of two closed, A-invariant subspaces X_u and X_0 such that

(a) X_u is the closure of the linear span of the unitary eigenvectors of A. (An eigenvector is called unitary, if its corresponding eigenvalue is of modulus 1.)
(b) $X_0 = \{x \in X : \lim_{n\to\infty} A^n(x) = 0\}$.

The Perron-Frobenius operator $\mathcal{L} : C(J(f)) \to C(J(f))$ is defined by the following formula

$$\mathcal{L}(g)(x) = \sum_{y \in f^{-1}(x)} g(y) \exp(\phi(y) - \mathrm{P}(\phi)),$$

where every critical point $y \in f^{-1}(x)$ is counted with its multiplicity. The main result on spectral properties of the Perron-Frobenius operator is the following theorem proven in [DU8] as Theorem 35 and later in [Pr6] (see also [Ly2] for the special case $\phi = 0$).

Theorem A2.4. The Perron-Frobenius operator $\mathcal{L} : C(J(f)) \to C(J(f))$ is almost periodic. Let ρ denote the Radon-Nikodym derivative of the equilibrium state μ for ϕ and f

with respect to the $\exp(\mathrm{P}(\phi)-\phi)$-conformal measure m. Then

$$(C(J(f)))_u = \mathbb{C}\rho \text{ and } (C(J(f)))_0 = \{\phi : \int \phi\, d\nu = 0\}.$$

Moreover, if $g = g_u + g_0$ with $g_u \in (C(J(f)))_u$ and $g_0 \in (C(J(f))_0$, then $g_u = (\int g\, dm)\rho$.

In [DPU] we have proved the following results improving spectral properties of the Perron-Frobenius operator and our knowledge about the Radon-Nikodym derivative $d\mu/dm$.

Theorem A2.5. Let $\phi : J(f) \to \mathbb{R}$ be a Hölder continuous potential. Then there are constants $L > 0$ and $\theta > 0$ such that

$$||\mathcal{L}^n(\phi) - \rho \int \phi\, dm||_\infty \le L \exp\bigl(-\theta\sqrt{n}\bigr)$$

for all $n \ge 0$.

Theorem A2.6. The Radon-Nikodym derivative of the equilibrium measure μ of the potential ϕ with respect to the $\exp\bigl(P(\phi)-\phi\bigr)$-conformal measure m, is a Hölder continuous function.

Using Gordin's method (see [Go]), as an immediate consequence of Theorem A2.5, we obtained in [DPU] the following.

Theorem A2.7. Every Hölder continuous function $g : J(f) \to \mathbb{R}$ satisfies the central limit theorem. Namely there exists $\sigma \ge 0$ such that

$$\lim_{n\to\infty} \frac{1}{n} \int \left(\sum_{k=0}^{n-1} (g \circ f^k - \mu(g))\right)^2 d\mu = \sigma^2,$$

and if $\sigma^2 > 0$, then for any $t \in \mathbb{R}$

$$\mu\left(\{z \in J(f) : \frac{1}{\sqrt{n\sigma^2}} \sum_{k=0}^{n-1} \bigl(g(f^k(z)) - \mu(g)\bigr) \le t\}\right) \to \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{t} \exp[-u^2/2]\, du.$$

We would like to notice that in [DPU] we have proved the result, called Rule II, which played an important role there and in later activity in the field, especially the activity of F. Przytycki (see [Pr3], [Pr4], [PR] and [PU3]). In order to state this result, define for every $c \in \overline{\mathbb{C}}$ the function $k_c : \overline{\mathbb{C}} \to \{0, 1, 2, ...\} \cup \{\infty\}$ by setting

$$k_c(x) = \min\{n \ge 0 : x \notin B(c,\ ae^{-(n+1)})\}.$$

and $k_c(x) = \infty$ if $x = c$. Here a is an arbitrary positive number such that $a < \mathrm{diam}\overline{\mathbb{C}}$. The Rule II is the following.

Theorem A2.8. (Rule II) There exists a constant $Q > 0$ such that if $c \in J(f)$ is a critical point of f, $n \geq 1$ is an integer, and if $x \in J(f)$ satisfies

$$k_c(f^j(x)) \leq k_c(f^n(x)) \quad \text{for every} \quad j = 1, 2, ..., n-1,$$

then

$$\min\{k_c(x), k_c(f^n(x))\} + \sum_{j=1}^{n-1} k_c(f^j(x)) \leq Qn.$$

We would like to conclude the above considerations with the remark that recently N. Haydn has improved (see [Hay]) Theorem A2.5 substantially by demonstrating the following.

Theorem A2.9. Let $\phi : J(f) \to \mathbb{R}$ be a Hölder continuous potential. Then there are constants $L > 0$ and $\theta > 0$ such that

$$||\mathcal{L}^n(\phi) - \rho \int \phi \, dm||_\infty \leq L \exp\bigl(-\theta n\bigr)$$

for all $n \geq 0$.

In [Ma4] R. Mann'e has proved that a power of each dynamical system $(J(f), f, \mu_0)$ is isomorphic to a one-sided Bernoulli shift, where μ_0 is the measure of maximal entropy. D. Heicklen and Ch. Hoffman have recently proved in [HH] the following improvement of Mané's result.

Theorem A2.10. Every dynamical system $(J(f), f, \mu_0)$ is isomorphic to a one-sided Bernoulli shift, where μ_0 is the measure of maximal entropy.

Acknowledgment. The author is deeply grateful to Bernd Stratmann who has carefully read the first draft of this article and has made many helpful comments and corrections.

References

[ADU] J. Aaronson, M. Denker, M. Urbański, Ergodic theory for Markov fibered systems and parabolic rational maps, Transactions of A.M.S. 337 (1993), 495-548

[Bea] A.F. Beardon, Iteration of Rational Maps, Springer-Verlag, New York, 1991.

[Bes] A.S. Besicovitch, 'Sets of fractional dimension(IV): On rational approximation to real numbers', Jour. London Math. Soc. 9 (1934) page 126-131.

[BJ] C. Bishop, P. Jones, Hausdorff dimension and Kleinian groups, SUNY preprint 1994.

[BZ] O. Bodart, M. Zinsmeister, Quelques resultats sur la dimension de Hausdorff des ensembles polynomes quadratiques, Fund. Math. 151 (1996), 121-137.

[Bo1] R. Bowen, Hausdorff dimension of quasi-circles, Publ. Math. IHES, 50 (1980), 11-25.

[Bo2] R. Bowen, Equilibrium states and the ergodic theory for Anosov diffeomorphisms. Lect. Notes in Math. 470, Springer, 1975.

[BK] M. Brin, A. Katok, On local entropy, in Geometric Dynamics, Lect. Notes in Math. 1007 (1983), 30-38, Springer Verlag

[CG] L. Carleson, T.W. Gamelin, Complex Dynamics, Springer-Verlag, New York, 1993.

[CJY] L. Carleson, P. W. Jones, J.-Ch. Yoccoz, Julia and John, Bol. Soc. Bras. Mat. 25 (1994), 1-30.

[DMNU] M. Denker, D. Mauldin, Z. Nitecki, M. Urbański, Conformal measures for rational functions revisited, Fundamenta Math. 157 (1998), 161-173.

[DPU] M. Denker, F. Przytycki, M. Urbański, On the transfer operator for rational functions on the Riemann sphere, Ergod. Th. and Dynam. Sys. 16 (1996), 255-266

[DR] M. Denker, S. Rohde, On Hausdorff Measures and SBR Measures for Parabolic Rational Maps, Preprint.

[DU1] M. Denker, M. Urbański, On the existence of conformal measures, Trans. A.M.S. 328 (1991), 563-587

[DU2] M. Denker, M. Urbański, On Sullivan's conformal measures for rational maps of the Riemann sphere, Nonlinearity 4 (1991), 365 – 384

[DU3] M. Denker, M. Urbański, Hausdorff and conformal measures on Julia sets with a rationally indifferent periodic point, J. London Math. Soc. 43 (1991), 107-118

[DU4] M. Denker, M. Urbański, On absolutely continuous invariant measures for expansive rational maps with rationally indifferent periodic points, Forum Math. 3(1991), 561-579

[DU5] M. Denker, M. Urbański, The Capacity of parabolic Julia sets, Math. Zeitsch. 211, (1992), 73-86

[DU6] M. Denker, M. Urbański, Geometric measures for parabolic rational maps, Ergod. Th. and Dynam. Sys. 12 (1992), 53-66

[DU7] M. Denker, M. Urbański, On Hausdorff measures on Julia sets of subexpanding rational maps, Israel J. of Math. 76 (1991), 193-214

[DU8] M. Denker, M. Urbański, Ergodic theory of Equilibrium states for rational maps, Nonlinearity 4 (1991), 103-134

[DH1] A. Douady, J.H. Hubbard, Etude dynamique des polynomes complexes I,II, Publications mathematique d'Orsay 84-2, 85-4, 1985.

[DH2] A. Douady, J.H. Hubbard, A proof of Thurston's topological characterization of rational functions, Acta Math. 171.2 (1993), 263-297.

[DSZ] A. Douady, P. Sentenac, M. Zinsmeister, Implosion parabolique et dimension de Hausdorff, C. R. Acad. Sci, Paris, 325 (Serie 1) (1997), 765-772.

[Fa] K. Falconer, Fractal geometry, Mathematical Foundations and Applications, John Wiley & Sons, 1990.

[FLM] A. Freire; A. Lopes; R. Mañé: An invariant measure for rational maps. Bol. Soc. Bras. Mat. 14, (1983), 45-62.

[Ge] L. Geyer, Porosity of parabolic Julia sets. To appear in Complex Variables.

[Go] M.I. Gordin, The central limit theorem for stationary processes. Dokl. Akad. Nauk SSSR **188**, (1969), 1174-1176.

[GS1] J. Graczyk, S. Smirnov, The Fibonacci Julia set, conformal measures and Hausdorff dimension, Preprint.

[GS2] J. Graczyk, S. Smirnov, Collet, Eckmann, & Hölder, Preprint 1996.

[Gr] M. Gromov, On the entropy of holomorphic maps, Preprint IHES.

[GPS] P. Grzegorczyk, F. Przytycki, W. Szlenk, On iterations of Misiurewicz's rational maps on the Riemann sphere, Ann. Inst. Henri Poincaré, 53 (1990), 431-434.

[Gu] M. de Guzmán, Differentiation of integrals in $\mathbb{R}^n$. Lect. Notes in Math. 481, Springer Verlag.

[HK] B. Hasselblatt, A. Katok, Introduction to the modern theory of dynamical systems, Encyclopedia of Mathematics and its applications 54, Cambridge University Press, 1995.

[Ha] G. Havard, Mesures invariantes pour les fractions rationnelles geometriquement finies, Preprint 1998.

[Hay] N. Haydn, Convergence of the transfer operator for rational functions, Preprint.

[HH] D. Heicklen, C. Hoffman, Rational maps are 1-sided Bernoulli, Preprint 1998.

[HS] S. Heinemann, B. Stratmann, Julia sets with dimensions close to 2, Math. Goettingensis Preprint, Heft 1 (1998).

[HV] R. Hill, S. Velani, Metric Diophantine approximation in Julia sets of expanding rational maps, Publ. Math. 85 (1997), 193-216.

[Ja] V. Jarník, Diophantische Approximationen and Hausdorff Mass, Mathematicheskii Sbornik 36 (1929) page 371-382.

[JM] P. Jones, N. Makarov, Density properties of harmonic measure, Annals of Math.,

[KR] P. Koskela, S. Rohde, Hausdorff dimension and mean porosity. To appear in Math. Ann.

[Ly1] M. Lyubich, On a typical behaviour of trajectories for a rational map of sphere, Dokl. Ak. N. U.S.S.R. 268 (1982), 29-32.

[Ly2] M. Lyubich, Entropy properties of rational endomorphisms of the Riemann sphere. Ergod. Th. Dynam. Sys. 3 (1983), 351-386.

[Ma1] R. Mané, The Hausdorff dimension of invariant probabilities of rational maps, Dynamical Sysstems, Valparaiso 1986, Lect. Notes in Math. 1331, Springer-Verlag (1988), 86-117.

[Ma2] R. Mané, On a theorem of Fatou. Bol. Soc. Brasil. Mat. 24 (1993), 1-12.

[Ma3] R. Mané, On the uniqueness of the maximizing measure for rational maps, Bol. Soc. Bras. Mat. 14 (1983), 27-43.

[Ma4] R. Mané, On the Bernoulli property of rational maps. Ergod. Th. Dynam. Sys. 5 (1985), 71-88.

[MC] A. Manning, H. McCluskey, Hausdorff dimension for horseshoes. Ergod. Th. and Dynam. Sys. 3 (1983), 251–260.

[Mar] M. Martens, The existence of σ-finite invariant measures, Applications to real one-dimensional dynamics, Preprint SUNY Stony Brook IMS preprint 1992/1.

[Mat] P. Mattila, Geometry of sets and measures in euclidean spaces, Cambridge Studies in Advanced Mathematics 44, Cambridge University Press, 1995.

[Mc1] C. McMullen, Hausdorff dimension and conformal dynamics II: Geometrically finite rational maps, Preprint 1997.

[Mc2] C. McMullen, Self-similarity of Siegel disks and Hausdorff dimension of Julia sets. Preprint 1996.

[Mc3] C. McMullen, Hausdorff dimension and conformal dynamics III: Computation of dimension.

[Mi] M. Misiurewicz, Topological conditional entropy. Studia Math. 55, (1976), 175-200.

[Pa] S. J. Patterson, The limit set of a Fuchsian group, Acta Math. 136 (1976), 241-273.

[Pra] E. Prado, Teichmüller distance for some polynomial-like maps, SUNY Stony Brook IMS preprint 1996/2, revision 1997.

[Pr1] F. Przytycki, Lyapunov characteristic exponents are non-negative. Proc. Amer. Math. Soc. 119.1 (1993), 309–317.

[Pr2] F. Przytycki, Hausdorff dimension of harmonic measure on the boundary of an attractive basin for a holomorphic map, Invent. Math. 80 (1985), 169-171.

[Pr3] F. Przytycki, Iterations of holomorphic Collet-Eckmann maps: conformal and invariant measures. Trans. AMS 350.2 (1998), 717-742.

[Pr4] F. Przytycki, On measure and Hausdorff dimension of Julia sets for holomorphic Collet-Eckmann maps, International conference on dynamical systems, Montevideo 1995, Pitman Research Notes in Math. 362 (1996), 167-181.

[Pr5] F. Przytycki, Sullivan's classification of conformal expanding repellors, Preprint 1991, to appear in the book "Fractals in the plane - ergodic theory methods by F. Przytycki and M. Urbański.

[Pr6] F. Przytycki, On the Perron-Frobenius-Ruelle operator for rational maps on the Riemann sphere and for Hölder continuous functions. Bol. Soc. Bras. Mat. 20 (1990), 95-125.

[PR] F. Przytycki and S. Rohde, Porosity of Collet-Eckmann Julia sets. Fund. Math. 155 (1998), 189-199.

[PU1] F. Przytycki, M. Urbański, Fractals in the Plane - the Ergodic Theory Methods, available on the web:http://www.math.unt.edu/˜urbanski, to appear in Cambridge Univ. Press.

[PU2] F. Przytycki, M. Urbański, Rigidity of tame rational functions, Preprint 1998, to appear Bull. Pol. Acad. Sci.

[PU3] F. Przytycki, M. Urbański, Porosity of Julia sets of non-recurrent and parabolic Collet-Eckmann rational functions, Preprint 1999.

[RT] C. Rogers, S. Taylor, Functions continuous and singular with respect to a Hausdorff measure, Mathematika, 8 (1961), 1-31.

[Ru] D. Ruelle, Thermodynamic formalism, Encyclopedia of Math. and Appl., vol. 5, Addison - Wesley, Reading Mass., 1976

[Sc1] F. Schweiger: Number theoretical endomorphisms with σ-finite invariant measures. Isr. J. Math. 21 (1975), 308-318.

[Sc2] F. Schweiger, Ergodic Theory of Fibred Systems and Metric Number Theory, Oxford Sci. Publ., Oxford, 1995.

[Sh] M. Shishikura, The Hausdorff Dimension of the Boundary of the Mandelbrot Set and Julia Sets, Ann. of Math. 147 (1998), 225-267.

[St1] B. Stratmann, Fractal dimensions for Jarnik limit sets; the semi-classical approach', Ark. för Mat. 33 (1995), 385-403.

[St2] B. Stratmann, Weak singularity spectra of the Patterson measure for geometrically finite Kleinian groups with parabolic elements', Preprint 1998.

[SU1] B.O. Stratmann, M. Urbański, The Geometry of Conformal Measures for Parabolic Rational Maps, to appear in Math. Proc. Cambr. Phil. Soc., 127 (3) 1999.

[SU2] B.O. Stratmann, M. Urbański, Jarnik and Julia; a Diophantine analysis for parabolic rational maps, Preprint 1999.

[SV] B.O. Stratmann, S. Velani, The Patterson measure for geometrically finite groups with parabolic elements, new and old, Proc. London Math. Soc. 71 (1995) page 197 - 220.

[Su1] D. Sullivan, Seminar on conformal and hyperbolic geometry. Preprint IHES (1982).

[Su2] D. Sullivan, Conformal dynamical systems. In: Geometric dynamics, Lect. Notes in Math. 1007 (1983), 725-752, Springer Verlag.

[Su3] D. Sullivan, Quasiconformal homeomorphisms in dynamics, topology, and geometry, Proc. Internat. Congress of Math., Berkeley, Amer. Math. Soc., 1986, 1216-1228.

[TT] S. J. Taylor, C. Tricot, Packing measure, and its evaluation for a Brownian path, Trans. A.M.S. 288 (1985), 679 - 699.

[Th] W. Thurston Three-Dimensional Geometry and Topology, Lecture Notes, Princeton University, 1978-1979.

[U1] M. Urbański, On Some Aspects of Fractal Dimensions in Higher Dimensional Dynamics, in Proc. of the Göttingen Workshop Problems on Higher Dimensional Complex Dynamics, Mathematica Gottingensis 3 (1995) 18-25.

[U2] M. Urbański, On Hausdorff dimension of Julia set with a rationally indifferent periodic point, Studia Math. 97 (1991), 167 – 188.

[U3] M. Urbański, Rational functions with no recurrent critical points, Ergod. Th. and Dynam. Sys. 14 (1994), 391-414.

[U4] M. Urbański, Geometry and ergodic theory of conformal nonrecurrent dynamics, Ergod. Th. and Dynam. Sys. 17 (1997), 1449-1476.

[Wa1] P.Walters, An introduction to ergodic theory, Springer-Verlag, 1982.

[Wa2] P.Walters, A variational principle for the pressure of continuous transformations, Amer. J. Math. 97 (1975), 937 - 971.

[Zd] A. Zdunik, Parabolic orbifolds and the dimension of maximal measure for rational maps, Inv. Math. 99 (1990), 627-649.

[Zi1] M. Zinsmeister (after A. Douady), Basic parabolic implosions in five days, Preprint 1997/8.

[Zi2] M. Zinsmeister, Fleur de Leau-Fatou et dimension de Hausdorff, Preprint 1998.